

Management of Gray Mold of Iceberg Lettuce by Biological Control Agents and Chitosan Formulations

R.G.A.S.Rajapakse¹, M.P.T.Premarathna¹, Shyamalee Kohombange^{1*}, D.G.N.S.B. Jayasinghe¹, H.A.S. Rohana¹, H.G.Gunathilake¹, K.R.C.de Silva² and A.K. Rathnayake²

¹Horticultural Crops Research and Development Institute, Gannoruwa, Peradeniya, Sri Lanka

²Atomic Energy Board of Sri Lanka, Colombo

ABSTRACT

Gray mold caused by *Botrytis* is the major problem in iceberg lettuce cultivation in poly tunnels in Sri Lanka. Currently management of this disease of lettuce depends mainly on foliar application of fungicides. Continuous application of fungicides for control of gray mold could not be recommended as lettuce mainly consumed as fresh vegetables. Therefore, studies were conducted to identify the causal agent, varietal resistance to pathogen and effective chitosan formulation and biological control agents on control of gray mold diseases of iceberg lettuce in poly tunnels. Causal agent of this disease was identified as *Botrytis cinerea*. Commercially grown varieties Eden and Maruli were equally susceptible to the disease. Different isolates of pathogen were shown different virulence levels on iceberg lettuce variety Eden. In vitro test showed that, almost complete inhibition of mycelia growth of all *Botrytis* isolates at 600 ppm chitopower 2 and liquid formulation of *Trichoderma asperellum* (4×10^5 conidia/ml) but 600 ppm chitopower 1 or liquid formulation of fluorescent *Pseudomonas* (10^6 bacteria/ml) or fungicide- Dicloran 75WP (3000ppm) were suppressed some isolates only. Studies in poly tunnel showed that lowest disease severity index (3.6% DSI) of gray mold in iceberg lettuce plants treated with *Trichoderma asperellum* and highest DSI (77.7 %) in control treatment. Dicloran 75WP, chitopower 2 and fluorescent *Pseudomonas* were recorded 16.6%, 18.5 and 46.2% DSI respectively. Results revealed that *Trichoderma asperellum* bio control agent and chitopower 2 could be used as alternatives to synthetic fungicides in controlling of gray mold disease of iceberg lettuce grown in poly tunnels.

Key Words: Biological control, Chitosan, Gray mold, Fungicide

1. INTRODUCTION

Gray mold disease is become an epidemic level of iceberg lettuce cultivations especially in poly-tunnels in Sri Lanka. Currently management of ~~betrytis gray mold~~ disease of crops depends mainly on foliar application of fungicides. Several fungicides ~~like Azoxystrobin, Chlorothalonil, Mancozeb, Pyraclostrobin, Thiophenate Methyl, and Captan~~ are ~~currently~~ used in Sri Lanka to control the disease by farmers. ~~These are Azoxystrobin, Chlorothalonil, Mancozeb, Pyraclostrobin, Thiophenate Methyl, and Captan~~. Farmers, however, claimed that application of fungicides is not effective when symptoms have developed in cool and humid condition in poly tunnels (Personal communication with growers).

Rajapakse and Premarathna, 2015 reported that *Botrytis cinerea* is causal agent of gray mold of lettuce in poly tunnels. They also revealed that continuous foliar spraying of 4g/l concentrations of fungicide, Dicloran 75 WP were reduced the gray mold of lettuce ~~caused by Botrytis~~. However, continues application of fungicides for control of ~~Botrytis gray mold~~ disease could not be recommended as lettuce mainly consume as fresh vegetables. Chitosan a natural biodegradable de-acetylated form of chitin has been proven to control numerous pre and post-harvest disease on various horticultural commodities (Aziz *et al.*, 2005, Kareem *et al.*, 2006, Wang *et al.*, 2007, Bautista-Banos *et al.*, 2005; [Farouk S; B E A Belal; H H A EL-Sharkawy 2017 The role of some elicitors on the management of Roumy Ahmar grapevines downy mildew disease and its related to inducing growth and yield characters. Scientia Horticulturae 225:646-658. <http://dx.doi.org/10.1016/j.scienta.2017.07.054>](#)

[Gomah A A; M E Abdallah; Samia A Haroun; S Farouk; Huda H Badr 2015 Exploitation of biostimulants as an alternative strategy to control bacterial rot diseases of onion \(*Allium cepa* L.\). Egyptian J Phytopathology 43 \(1-2\): 65-77](#)

[Farouk S; Safaa A Youssef; Abeer A Ali 2012 Induction of systemic resistance in tomato against *Alternaria solani* by biostimulants and vitamins. Alex. J. Agric. Res. 57 \(1\): 117-129.](#)

[Farouk S; Safaa A Youssef; Abeer A Ali 2012 Exploitation of biostimulants and vitamins as an alternative strategy to control early blight of tomato plants. Asian Journal of Plant sciences, 11\(1\): 36-43. DOI:10.3923/ajps.201237370-AJPSANSI](#)). It has reported that chitosan induces a series of defense reactions in plants correlated with enzymatic activities and also it has a direct effect on microbes by fungistatic or fungicidal potential (Meng *et al.*, 2010, Raafat and Sahl, 2009; [Belal; H H A EL-Sharkawy 2017 The role of some elicitors on the management of Roumy Ahmar grapevines downy mildew disease and its related to inducing growth and yield characters. Scientia Horticulturae 225:646-658. <http://dx.doi.org/10.1016/j.scienta.2017.07.054>](#)). Efficacy of the different chitosan compounds on growth inhibition of fungal pathogen such as *Colletotrichum* and *Fusarium* species was demonstrated by in vitro test (Rajapakse *et al.*, 2014). Chenyan Hua *et al.*, 2009 reported that chitosan has a greater inhibitory effect on Kiwi fruit gray mold and Ben Shalon *et al.*, 2003 reported that almost complete inhibition

of *Botrytis* conidia germination (in vitro) and controlled the gray mould of cucumber caused by *B. cinerea* by chitosan. There were several research reports indicated that selected strains of biological control agents *Pseudomonas fluorescens* and *Trichoderma* species had significantly reduced lesion areas on whole lettuce plants caused by *Botrytis cinerea* (Card, 2005, Freeman *et al.*, 2004, Walker *et al.*, 2001). It has been reported that *Trichoderma harzianum* control *Botrytis cinerea* and many other fungi by acquiring induce systemic resistance of plants and antagonism of nutrient activated conidia against *Botrytis cinerea* (Elad, 2000, Freeman *et al.*, 2004, Hjelyord *et al.*, 2001, Meyer *et al.*, 1998). Similarly Walker *et al.*, 1996 and 2001 demonstrated that antifungal activity of *Pseudomonas antimicrobica* by inhibiting conidia germination and outgrowth of *Botrytis cinerea*. Application of agro-chemicals are the major plant protection method over decades even though they are associated with many disadvantages including their expensive applications, environmental pollution and human health hazards due to excessive usage. This has emerged a worldwide huge trend to explore other environmental friendly alternative methods for plant protection. Therefore, the objectives of this study were to identify the causal agent of grey mold disease, resistance varieties and effective chitosan formulations and biological control agents on control of gray mold of iceberg lettuce in poly tunnels.

2. MATERIALS AND METHODS

2.1 Pathogen isolation and identification

Disease samples of Iceberg lettuce were collected from poly tunnels in hill country Nuwara Eliya area. Pathogen was isolated from lesions of disease affected leaves of iceberg lettuce on Potato Dextrose Agar (PDA). Single spore isolates of pathogen were prepared from mycelia of single conidia cultures grown on PDA. Pathogen was identified on the basis of microscopic observations of size and shape of conidia, mycelia arrangements, culture morphology on PDA and comparison with published data (Card 2000, Ellis and Waller, 1974). Five isolates of *Botrytis* was selected for further studies and mycelia growth rate on PDA and conidial diameter of each isolate was measured at 7 days after inoculation. The isolates of pathogen were cultured on PDA slant and stored for further studies.

2.2 Pathogenicity of botrytis isolates on iceberg lettuce

Pathogenicity of all isolates was tested by wound inoculation with conidia suspension and subsequent symptoms development on leaves. Five isolates of *Botrytis* obtained from diseased plants and iceberg lettuce plants of variety Eden was used for the pathogenicity test. One month old lettuce plants grown in poly tunnel under 18-26°C temperature and 85-90% relative humidity were inoculated with isolates of *Botrytis* by pin prick methods using conidia suspension (10^6 conidia/ml) of different isolates of *Botrytis*. Five iceberg lettuce plants were used to inoculate each *Botrytis* isolate. Un-inoculated plants were served as control. Disease symptoms were observed after 5 days of inoculation and re-isolation of pathogen was done on PDA medium to perform Koch postulation (Card 2000, Fernandez-Ortuno *et al.*, 2011).

2.3 Response of iceberg lettuce varieties to infection of Botrytis isolates.

Experiment was conducted in poly tunnel (16-28°C, 80-90% relative humidity) at Dolosbage area (WU1) in Kandy District, Sri Lanka. Two commercially grown iceberg lettuce varieties Eden (V1) and Maruli (V2) and five *Botrytis* isolates (Br, Bs, Bl, Bp and Bt) were used for inoculation with lettuce varieties. Experimental design was completely Randomized design with two factor factorial. One month old ice berg lettuce plants of both varieties were inoculated with conidia suspension (10^6 conidia/ml) of different *Botrytis* isolates using pin prick method to observe subsequent symptoms development on plants. Disease Severity of gray mold in Iceberg lettuce varieties was measured using rating scale at harvest (Elad *et al.*, 1992). Symptoms of leaf tissue were evaluated according to Disease Severity Index of 0-4, where 0=no disease, 1=1-12% rot, 2=13-25%, 3=26-50%, 4=51-100%. The data obtained were tabulated and analyzed subjected to the analysis of variance procedure of SAS 9.1 software. Duncan's Multiple Range Test was performed to compare the differences among treatment means at $p=0.05$.

2.4 Effect of biological control agents, Chitosan-chitosan formulations and fungicide (Dicloran 75WP) on mycelia growth of Botrytis isolates (in vitro).

Studies (*in vitro*) were conducted with the treatments of two formulations of chitosan i.e. Chitopower 1 (600ppm), chitopower 2 (600ppm) developed by Sri Lanka Atomic Energy Board and two biological control agents developed by Horticultural Crops Research and Development Institute, Sri Lanka i.e. liquid formulation of *Trichoderma asperellum* (4×10^5 conidia/ml) and fluorescent *Pseudomonas* (1×10^6 bacteria/ml) and fungicide- Dicloran 75WP (3000ppm). Untreated plants were served as control. Five isolates of *Botrytis* were used for each study. Experimental procedure was Completely Randomized Design and each treatment replicated five times. Effectiveness of Chitopower 1, Chitopower 2 and fungicide Dicloran 75WP in controlling of mycelia growth of *Botrytis* isolates was measured by culturing them on amended PDA medium. Effectiveness of biological control agents in controlling mycelia growth of *Botrytis* isolates was measured by duel culture techniques on PDA medium. All PDA plats were incubated at 28 °C for 7 days. Percent inhibition of fungi (growth reduction over control) was calculated by the following equation.

$$I = \frac{100(C-T)}{C}$$

Where I is the % inhibition of mycelia growth, C the growth of fungus in control plate (mm), and T is the growth of fungus on the treated plate (mm).

2.5 Effect of biological control agents, chitopower 2 and fungicide Dicloran 75WP on gray mold of Iceberg lettuce grown in Poly tunnel.

Experiment was conducted in poly tunnels at Dolosbage area (WU1) in Kandy District, Sri Lanka. Mean day and night temperature in the location was 23 °C and 16 °C respectively. RH was over 90% throughout the day. Ice berg lettuce variety Eden was used for experiment.

Plants were established in pots and standard crop management practices were done throughout the study. Top of the areal part of each ice berg plants were artificially inoculated with conidial suspension (10^6 conidia/ml) of most virulent *Botrytis* isolate BI by pin prick method when plants were one month old. The treatments were selected based on the results of *in vitro* test (Table 3) and applied to the plants to cover whole aerial parts of the plant as an aqueous spray by using a knapsack sprayer one day after artificial inoculation of pathogen and continued 5 times with 7 days intervals. Fertilizer mixture of urea, super phosphate and MOP 6: 30: 30 was used as basal and top dressing fertilizer. The experiment was laid out in a Completely Randomized Design and five treatments were randomized in with ten replicates. Data were collected at harvesting stage. Measurements were taken on Disease Severity Index (%) (Elad *et al.*, 1992) and yield loss compared to most effective treatment. The data obtained were tabulated and analyzed subjected to the Analysis of Variance procedure of SAS 9.1 software. Duncan's Multiple Range Test was performed to compare the differences among treatment means at $p=0.05$.

Treatments were used as follows.

- T1- Foliar spraying of liquid formulation of *Trichoderma asperellum* (4.5×10^5 conidia/ml)
- T2- Foliar spraying of liquid formulation of fluorescent *Pseudomonas* (1×10^6 conidia/ml)
- T3- Foliar spraying of [Chitopower chitopower II](#) (600ppm)
- T4- Foliar spraying of [Fungicidefungicide](#)- Dicloran 75WP (3000ppm)
- T5- control

RESULTS AND DISCUSSION

Gray mold symptoms were observed in iceberg lettuce plants grown in poly tunnels at Nuwara Eliya area in Sri Lanka. The symptoms observed were water-soaked, brown or gray blight, and abundant mycelia with conidia were appeared on the infected leaves. Isolates of *Botrytis* were identified by comparison of their colony morphology on PDA, microscopically observations of size, shape of conidia and mycelia arrangements with published data (Ellis and Waller, 1974).

The fungus produced light gray to gray colonies with dark mycelium and abundant conidia on PDA at 26°C for 7 days. It was observed that significant variation of mycelia growth rate of different isolates on PDA. The conidia of all isolates were one-celled, ovoid in shape, dark brown, and average size of conidia of different isolates was varied from 12-14x9-11 µm. Identification of *Botrytis cinerea* was from its typical spore and conidiophores structure. Willets, 1997 reported that conidiophores are specialized structure

that incorporates a terminal cluster of synchronously produced conidia, borne on a well-developed conidiogenous hypha, which resembles a bunch of grapes. Primary cultures of isolates on PDA have a light grey at first, later becoming grey or grayish-brown (Table 1) with dark walled erect septate hyphae. The dark coloured mature conidiophores, which branch alternately. This conformed to the description of *Botrytis cinerea* (Ellis and Waller 1974).

To determine the pathogenicity test, conidial suspension (1×10^6 conidia/ml) was inoculated with iceberg lettuce plants variety Eden by pin prick methods and allow to grow in poly tunnel (16-28°C, 80-90% relative humidity). After 5 days, grey mold symptoms similar to the original symptoms were developed on inoculated plant leaves. The fungal pathogen was re-isolated from the disease lesions of the inoculated plants and the re-isolated pathogen exhibited the same morphological characteristics as those of the original isolates. Thus, the isolates of fungal pathogen fulfilled the criteria stipulated by the Koch's postulates and were identified as the causal agent of the gray mold on iceberg lettuce plants.

Table 1. Characteristic of fungal isolates of *Botrytis* collected from leaves of infected plants

Botrytis isolates	Colony colour on PDA at 7 DAI	Colony diameter on PDA at 7 DAI	Average size of conidia (µm)
Bs	Light gray at first, later Grey, reverse black,	5.6 ^c	12x9
Bp	Grey at first, later grayish-brown, reverse black	9.2 ^a	13x9
Bt	Grey at first, later grayish-brown, reverse black	9.2 ^a	14x11
Br	Light gray at first, later gray, reverse black	6.4 ^b	13x10
Bl	Grey at first, later grayish-brown, reverse black	9.1 ^a	13x9

Note: Means with the same letter(s) on the column are not significantly different at P= 0.05 [please indicate the LSD value](#)

Varietal response of iceberg lettuce to different isolates of *Botrytis* showed that there were significant variations of percent disease severity index (DSI%) among *Botrytis* isolates (Table 2). Both iceberg lettuce varieties Eden and Maruli were equally susceptible to all *Botrytis* isolates. However, *Botrytis*

isolate-BI could be considered as most virulent isolates as it had the highest DSI on both iceberg varieties compared other isolates.

Table 2: Mean Disease Severity Index (DSI %) of grey mold of lettuce varieties inoculated with different isolates of Botrytis.

Varieties	Disease Severity Index (%)					
	Br	Bs	B1	Bp	Bt	Mean
V1	17.0	20.7	98.5	8.9	11.1	31.0
V2	12.6	11.8	92.6	8.9	2.9	25.8
Mean	14.8 ^b	16.3 ^b	95.6 ^a	8.9 ^c	7.0 ^c	

Note: Means with the same letter(s) are not significantly different at P= 0.05 [please indicate the LSD value](#)

V1 & V2: lettuce variety Eden and Maruli

Br, Bs, B1, Bp&Bt: different Botrytis isolates

Study (*in vitro*) was conducted to identify the effect of two chitosan formulations ([Chitopower-chitopower 1](#) and Chitopower 2), two biological control agents (formulation of *Trichoderma asperellum* and fluorescent *Pseudomonas*) and fungicide- Dicloran 75WP (3000ppml) on growth of different isolates of *Botrytis* on PDA (Table 2).

Table 3:Effect of [Chitopower-chitopower 1](#), chitopower 2, *Trichoderma asperellum*, fluorescent *Pseudomonas* and fungicide- Dicloran 75WP on mycelia growth of *Botrytis* isolates (*in vitro* test).

Botrytis isolate	Growth reduction % of Botrytis isolates on PDA at 7 DAI				
	600ppm Chitopower 1	600ppm Chitopower 2	Dichloran 75WP (3000ppm)	<i>Trichoderma asperellum</i>	fluorescent <i>Pseudomonas</i>
Br	100	100	79.3	100	100
Bt	83.2	100	71.4	100	46.3
B1	54.6	100	100	100	100
Bs	94.4	100	56.9	100	23.3
Bp	100	100	47.3	100	100

Results revealed that chitopower 2 (600ppm) and *Trichoderma asperellum* bio-control agent completely suppressed mycelia growth of all tested isolates of *Botrytis* (in vitro) (Table 3). Biological control agent fluorescent *Pseudomonas* also completely suppressed most virulent *Botrytis* isolate i.e. B1 and other two isolates Br and Bp. Chitopower 1 (600ppm) completely suppressed only Br and Bp i.e. less virulent isolates on both iceberg lettuce varieties.

Ben Shalon *et al.*, (2003) reported that almost complete inhibition of *Botrytis* conidia germination was found, in vitro by chitosan molecule which is similar to chitopower 2 but not with chitosan oligomers which is similar to chitopower 1. They also reported that chitosan controlled the gray mould caused by *Botrytis cinerea* compared with control but chitin oligomers did not show any effect on disease control. They further concluded that although a dual mode of action was involved in the control of gray mould by chitosan, the antifungal activity of the compound was an essential factor. It was evident that the induction of the defense response without the antifungal activity was not enough to suppress the diseases. Therefore, chitopower 2 (600ppm), *Trichoderma asperellum* and fluorescent *Pseudomonas* was selected to identify the effective and environmental friendly disease control practice for gray mold of iceberg lettuce grown in poly tunnels.

Table 4: Mean Disease Severity Index % of treatments and percentage yield loss compared to most efficient treatment when iceberg lettuce plants (var. Eden) treated with chitopower 2, biological control agents *Trichoderma asperellum* and fluorescent *Pseudomonas* and fungicide Dichloran 75WP.

Treatments	Mean DSI (%) of gray mold	Percentage yield loss compared to most effective treatment (T1)
T1- <i>Trichoderma asperellum</i>	3.6 ^c	
T2- fluorescent <i>Pseudomonas</i>	46.2 ^{ab}	33
T3- Chitopower 2	18.5 ^{bc}	11
T4- Dichloran 75WP	16.6 ^{bc}	9
T5 – Control	77.7 ^a	85

Note: Means with the same letter(s) are not significantly different at P= 0.05 [please indicate the LSD value](#)

The lowest yield loss was observed in *Trichoderma asperellum* treated plots while highest yield loss was observed in control plots (Table 4). Benitez *et al.*, (2004) reported that *Trichoderma* strains exert bio control against fungal pathogens either indirectly by competing for nutrients and space, modifying the environmental conditions or promoting plant growth and plant defensive mechanisms and antibiosis or directly by mechanisms such as mycoparasitism. Chitopower 2 and fungicide Dichloran 75 WP treated plots also recorded comparatively lower DSI (%) and yield losses (Table 4). Chenyan Hua *et al.*, 2009 reported that low molecular weight chitosan has a greater inhibitory effect on Kiwi fruit gray mold due to induction of accumulation of higher levels of hydrogen peroxide, greater level of activity of several defense-related enzymes, greater ability to penetrate the cell walls in kiwifruit epidermal peel tissue and activate a greater defense response. Similarly the low molecular weight chitosan which was used to produce Chitopower 2 formulation showed remarkable disease control ability of gray mold of iceberg lettuce. Ben Shalonet *et al.*, 2003 reported that almost complete inhibition of *Botrytis* conidia germination (in vitro) and controlled the gray mould of cucumber caused by *B. cinerea* by chitosan. Results in DSI and yield loss data indicated that gray mold of iceberg lettuce was effectively controlled by spraying of liquid formulation of *Trichoderma asperellum* bio control agent as well as chitopower 2 starting from early crop growth stage and continuing up to harvesting stage with 7 days intervals.

CONCLUSION

Causal agent of gray mold of iceberg lettuce in poly tunnels was re-conformed as *Botrytis cinerea*. Commercially grown varieties Eden and Maruli were equally susceptible to disease. Almost complete inhibition of mycelia growth of all *Botrytis* isolates was found, in vitro, at 600 ppm chitopower 2 and liquid formulation of *Trichoderma asperellum*. Gray mold of iceberg lettuce caused by *Botrytis cinerea* in poly tunnels was effectively controlled by spraying of liquid formulation of *Trichoderma asperellum* bio control agent as well as Chitopower 2, starting from early crop growth stage and continuing up to harvesting stage with 7 days intervals.

REFERENCES

- Aziz Aziz, Patrica Trotal-Aziz, Laurent Dhuicq, Philippe Jeandet, Michel Couderchet and Guy Vernet. Chitosan oligomers and copper sulfate induce grapevine defense reaction and resistance to gray mold and downy mildew. *Histopathology*. 2006; 96(11):1188- 1189.
- Bautista-Banos S, AN Hernandez-Lauzardo, Velazquez-del Valle MG, Hernandez-Lopez M, Ait Barka E, Bosquez-Molina E, Wilson, C.L. Chitin as a potential natural compound to control pre and postharvest diseases of horticultural commodities. *Mexican J. Phytopathol*, 2005; 25(2):108-118.

Ben-shalom, N., R. Ardi, R. Pinto, C. Aki and Fallik, E. Controlling gray mould caused by *Botrytis cinerea* in cucumber plants by means of chitosan. *Crop Protection*, 2003; 22(2):285-296.

Benitez, T., A.M. Rincon, M.C. Limon and Codon, A. C. Bio control mechanisms of *Trichoderma* strains. *Int. Microbiol*, 2004, 7(4):249-260.

Card, S. D. Biological control of *Botrytis cinerea* in lettuce and strawberry crops. PhD thesis, Lincoln University, Canterbury, New Zealand. 2005.

Chenyan Hua, Yulin Li, Xiaofu Wang, Kai kai, Mengke Su, Weishi, Danfeng Zhang and Yongsheng Liu. The effect of low and high molecular weight chitosan on the control of gray mold (*Botrytis cinerea*) on kiwifruit and host response. *Scientia Horticulturae*, 2019, Vol 246, 27 Feb 2019, page 700-709

Eled, Y., Krishner, B. and Gotlib, Y. Attempts to control *Botrytis cinerea* of rose by pre- and post harvest treatments with biological and chemical agents. 1992, *Crop Protection*, 12: 69-73.

Elad, Y. *Trichoderma harzianum* T39, Preparation for bio control of plant diseases-control of *Botrytis cinerea*, *Sclerotinia sclerotiorum* & *Cladosporium fulvum*. *Bio control Science & Technology*, 2000, **10**: 499-507.

Ellis, M. B. and Waller, J.M. *Sclerotinia fuckeliana* (Conidial state: *Botrytis cinerea*). CMI description of pathogenic fungi and bacteria. 1974.

Freeman, S., Minz, O., Kolesnik, I., Barbul, O., Zveibil, A., Maymon, M., Nitzani, Y., Kirshner, B., Rav-David, D., Bilu, A., Dag, A., Shafir, S. and Elad, Y. *Trichoderma* bio control of *Colletotrichum acutatum* & *Botrytis cinerea* & survival in strawberry. *European Journal of Plant Pathology*. 2004, 110: 361-370.

Frenandez-ortuno, X. Li., W. Chai and Schnabel, G. First report of gray mold of strawberry caused by *Botrytis cinerea* in South Carolina. The American Phyto pathological Society. 2011. <https://doi.org/10.1094/PDIS-06-11-0529>.

Hjeljord, L. G., Stensvand, A. and Tronsmo, A. Antagonism of nutrient-activated conidia of *Trichoderma harzianum* (atroviride) PI against *Botrytis cinerea*. *Phytopathology*. 2011, 91: 1172-1180.

Kareem, A. E., Nehals, F., Mougy, E., Nadia, G., Gamal, E. and Fouth, Y. O. Use of chitin and chitosan against tomato root rot disease under greenhouse condition. *Research Journal of Agricultural and Biological Science*, 2006, 2:147 – 152.

Meng, Z., Yang, L., Kennedy, J. F. and Tian, S. Effects of chitosan and oligochitosan on growth of two fungal pathogens and physiological properties in pear fruit. *Carbohydrate Polymers*.2010, 81, 70–75.

Meyer, D. G., Bigirimana, J., Elad, Y. and Hofte, M. Induced systemic resistance in *Trichoderma harzianum* T39 biocontrol of *Botrytis cinerea*. *European Journal of Plant Pathology*.1998, 104: 279-286.

Rajapakse, R. G. A. S., Edirimanna, E.R.S.P., Prmaratne, P., Kahawatte, J., Kulathunga, S.S., Dissanayake, C.K. and De Silva, K.R.C. Effect of Chitosan on mycelia growth of two pathogenic fungi; *Colletotrichum gloeosporioides*, *Fusarium oxysporum* sp. *cubense*. Proceedings-SLCARP International Agriculture Symposium, Colombo, Sri Lanka.2014.

Rajapakshe, R.G.A.S. and Premarathna P. Management of Botrytis Leaf Blight of Lettuce with Fungicides. *Crop Life: Sri Lanka Plant Protection Industry Journal*. 2015, 8:6-11.

Raafat, D. and Sahl, H. G. Chitosan and its antimicrobial potential – a critical literature survey. *Microbial Biotechnology*, 2009.2:186 – 201.

Walker, R., Emslie, K. A. & Allan, E. J. Bioassay methods for the detection of antifungal activity by *Pseudomonas antimicrobica* against the grey mould pathogen *Botrytis cinerea*, *Journal of Applied Bacteriology*,1996,81: 346-348.

Walker, R., Innes, C. M. J. and Allan, E. J. The potential biocontrol agent *Pseudomonas antimicrobica* inhibits germination of conidia & outgrowth of *Botrytis cinerea*. *Letters in Applied Microbiology*, 2001. 32: 346-348.

Willems, H. J. Morphology, development & evolution of stromatalsclerotia & macro conidia of the *Sclerotiniaceae*. *Mycological Research*.1997, 101: 939-952.

Wang, J., Wang, B., Jiang, W. and Zhao., Y. Quality and shelf life of, mango (*Mangifera indica* L. cv. 'Tainong') coated by using chitosan and polyphenols. *Food Science and Technology International*, 2007.13: 317-322.