

Protein and mineral contents in some fish species available in the Brahmaputra River of Bangladesh

Deleted: r

ABSTRACT

An experiment was conducted in the laboratory of the Department of Agricultural Chemistry, Bangladesh Agricultural University (BAU), Mymensingh to determine protein and major mineral nutrients (viz. Ca, Mg, Na, K, P, S and Fe) in different available fish species of the Brahmaputra river of Bangladesh. Total 32 fish samples of 15 fish species were collected from three locations (viz. Bhabakhali bazar, BAU campus and Mymensingh town) of the Brahmaputra river during November 2017. The highest amount of Ca (2.00%), Mg (4.17%), Na (0.41%), K (3.24%), P (0.17%), S (0.129%) and Fe (226.9 mg kg⁻¹) were obtained from the *chela* (*Salmophasia bacaila*), *chanda* (*Chanda nama*), *chingri* (*Macrobrachium* sp.), *shingi* (*Heteropneustes fossilis*), *bele* (*Glossogobius giuris*), *baim* (*Macrognathus aculeatus*) and *mola* (*Amblypharyngodon mola*), respectively and the sequence of the mineral nutrients was K > Mg > Ca > Na > P > S > Fe. The study results revealed that out of 15 fish species, 3, 11 and 12 species can contribute 100% of Ca, Mg and K requirement for human by taking 100g fish flesh, respectively. The maximum content of N (3.88%) was obtained from *shingi* (*Heteropneustes fossilis*), while the minimum (2.81%) was recorded from *mola* (*Amblypharyngodon mola*). The protein content among the fish samples varied between 17.6-24.3% with a mean value of 21.2%. Finally, the study concluded that the common fishes available in the Brahmaputra river are a good source of protein and major mineral nutrients, which contributes in nutrition to the local people of the country.

Comment [h1]: Give name of species

Keywords: Protein, minerals, fish species, the Brahmaputra River, Bangladesh

Deleted: r

1. INTRODUCTION

Fish is commonly found in natural water bodies and well known for its superior nutritional quality with a very good supply of essential minerals [1-3]. Furthermore, fishes found in river are very popular in Bangladesh. A few years ago, fishes available in natural water bodies were the major consumed protein sources of people in this country, as they were easily accessible, less expensive, culturally acceptable and can be purchased in small quantities [4]. But nowadays, discharge of industrial wastewater/ effluents into river, canal or other surface water bodies without any treatment is the common scenario in Bangladesh [5-9]. These wastewater/ effluents contained metals and other toxic substances [10-14], which often lead to decrease in number of fish species in rivers and canals of Bangladesh.

However, mineral elements are essential for the normal maintenance of the human body, and these elements are mainly provided to the body through dietary intake. The human being may suffer from diseases like anaemia, osteoporosis, goitre, stunted growth and genetic disorders due to insufficient dietary intake of minerals [15-17]. The World Health Organisation reported that about 2 billion of the world's population is suffering from mineral and vitamin deficiencies and the majority of these are in the third world countries [18].

Surface water fishes are a great resource which is easily accessible to low income people and vulnerable communities prone to nutrient deficiency diseases. Nowadays mineral supplementation and food fortification are the strategies used for mitigating nutritional

36 deficiencies but unsustainable especially for developing countries [15]. On the other hand,
37 food based strategies are considered very much effective as well as sustainable for
38 mitigating mineral and nutritional deficiencies. Fish can play a big role for this strategy
39 because it can provide a variety of nutrients, including essential elements to the body [3].
40 Minerals commonly found in fish flesh are sodium, potassium, calcium, magnesium,
41 phosphorus, sulphur, iron, zinc and iodine [19-20]. These mineral elements participate in
42 several biochemical reactions, viz. calcium, magnesium and phosphorus are crucial in the
43 formation of bones and teeth; sodium and potassium work together in the transmission of
44 nerve impulses and keeping electrolyte balance; zinc is mostly found as a cofactor in
45 enzyme reactions, iron forms part of the haemoglobin molecule which transport oxygen
46 around the body [21]. But there are published reports that the nutritional quality of fish
47 depends on their species, age, size, feed and water quality [22-25].
48

49 The Brahmaputra River is one of the major rivers in Bangladesh, which provide a variety of
50 different fish species to the local people of the country. In the meantime, a total of 67 finfish
51 species including 63 indigenous and 4 exotic/alien species have been recorded from the
52 Brahmaputra River belonging to 46 genera, 24 families and 8 orders [26]. Considering the
53 fact stated above, this study was undertaken to quantify the protein and mineral contents in
54 common fish species found in the Brahmaputra River and identify the species with high
55 protein and mineral contents which can be recommended for consumption in combating
56 mineral and protein deficiencies of the people in Bangladesh.
57

58 2. MATERIAL AND METHODS

59 2.1 Study Area

60 The Brahmaputra River is an antecedent, snow fed, and large trans-Himalayan river
61 originating in southern Tibet (China). Geologically, and it is the youngest of the major rivers
62 of the world and unique in many respects. It runs for a length of 2880 km and its river system
63 drains parts of China, Bhutan, India and Bangladesh [27]. However, total of 32 fish samples
64 from 15 fish species were collected from three locations (viz. Bhabakhali bazar, BAU
65 campus and Mymensingh town) of the Brahmaputra River. Details about the fish species
66 along with their sampling locations are presented in Table 1 and Fig. 1, respectively.
67
68

69 2.2 Collection of Fish Samples

70 A reasonable amount (500 g to 1.0 kg) of fish samples were purchased directly from the
71 aforementioned locations of the river system, and requisite amount of samples were brought
72 to the laboratory of the Department of Agricultural Chemistry, BAU, Mymensingh and
73 processed for subsequent experiment.
74

75 2.3 Processing of Fish Samples

76 After collection, scales of fish samples were removed and cleaned first. Then the fish
77 samples were separated into edible part and non-edible part. After separation, edible part of
78 fish samples were sun dried for 2 days, and then the samples were oven dried at 50-60°C
79 for another 2-3 days until a constant weight was obtained. After drying, the samples were
80 ground well with the help of mortar and pestle, and then the powdered samples were
81 preserved in polythene bags with appropriate marking for further chemical analyses.

Deleted: r

Deleted: r

Deleted: r

Deleted: the Brahmaputra

Deleted: of

Deleted: r



Fig. 1. Map shows the sampling sites of collected fish samples from the Brahmaputra river of Bangladesh

2.4 Extraction of Fish Samples

Powdered fish samples were used to prepare extract for the determination of different mineral nutrients. Extract was prepared by wet oxidation method using di-acid mixture [28]. In this method, approximately 0.5 g of finely ground samples were taken into a 250 mL conical flask and 5 mL of di-acid mixture ($\text{HNO}_3:\text{HClO}_4 = 2:1$) was added to it. Then the flask was placed on an electric hot plate for heating at 180-200°C temperature until the solid particles disappeared and white fumes were evolved from the flask. Then, it was cooled at room temperature, washed with distilled water and filtered into 100 mL volumetric flask through filter paper (Whatman No. 1). Finally, the volume was made up to the mark with distilled water and preserved for the determination of major mineral nutrients in the fish samples.

2.5 Determination of Nitrogen and Protein

Total nitrogen content of the fish samples was determined by Kjeldahl method by digestion with concentrated H_2SO_4 and digestion mixture ($\text{K}_2\text{SO}_4:\text{CuSO}_4:\text{Selenium powder} = 100:10:1$) and then distilling with 40% NaOH. The ammonia distilled over was absorbed in boric acid indicator and titrated against 0.05N H_2SO_4 [29]. The % N in edible parts of fish sample was calculated by using the following formula-

Deleted: Preparation of

$$\% N = \frac{(T - B) \times N \times 1.4}{W}$$

111 Where, T = Actual titration reading (mL); B = Blank titration reading (mL); N = Normality of
 112 H₂SO₄ and W = Sample weight in g. The crude protein % was calculated by multiplying the
 113 total nitrogen % with a factor of 6.25.

114
 115
 116
 117

Table 1. Details of fish samples collected from three locations of the Brahmaputra river, Bangladesh

SL. No.	Bengali name	English name	Scientific name	Sampling location*			Total
				Bhabakhali bazar	BAU campus	Mymensingh town	
1.	<i>Punti</i>	Ticto barb	<i>Puntius ticto</i>	1	1	1	3
2.	<i>Tengra</i>	Striped dwarf catfish	<i>Mystus vittatus</i>	-	1	1	2
3.	<i>Bele</i>	Tank goby	<i>Glossogobius giuris</i>	1	1	1	3
4.	<i>Bangna</i>	Reba	<i>Labeo ariza</i>	1	1	-	2
5.	<i>Baim</i>	Zig-zag eel	<i>Macrogathus aculeatus</i>	-	1	1	2
6.	<i>Shingi</i>	Stinging catfish	<i>Heteropneustes fossilis</i>	-	1	1	2
7.	<i>Kalibaush</i>	Orange-fin labeo	<i>Labeo calbasu</i>	1	-	1	2
8.	<i>Mola</i>	Mola carplet	<i>Amblypharyngodon mola</i>	1	1	-	2
9.	<i>Chela</i>	Silver razor-belly minnow	<i>Salmophasia bacaila</i>	1	-	1	2
10	<i>Kaikka</i>	Garfish	<i>Xenentodon cancila</i>	1	-	1	2
11	<i>Meni</i>	Gangetic leaffish	<i>Nandus nandus</i>	1	-	1	2
12	<i>Chanda</i>	Elongate glassy perchlet	<i>Chanda nama</i>	1	-	1	2
13	<i>Chingri</i>	Prawns or shrimps	<i>Macrobrachium sp.</i>	1	-	1	2
14	<i>Guizza</i> or <i>Guizza ayre</i>	Giant-river catfish	<i>Sperata seenghala</i>	1	-	1	2
15	<i>Hiralo</i> or <i>Murari</i>	Carplet	<i>Aspidoparia murar</i>	-	1	1	2
Total				11	8	13	32

* Number in the column indicates type of fish species present in that particular location

118
 119
 120
 121
 122
 123
 124
 125
 126
 127

2.6 Determination of Major Mineral Elements

Among the major mineral nutrient elements, Ca and Mg were determined by titrimetrically, P and S were measured spectrophotometrically (660 and 425 nm absorbance wavelength, respectively; T60 UV-Visible Spectrophotometer, PG Instrument, UK) and Na and K were estimated by flame photometrically (589 and 766 nm emission wavelength, respectively; 0.2 ppm limit of detection; Jenway PFP7, Flame Photometer, UK) [28]. Determination of Fe in the fish extracts was done by atomic absorption spectrophotometer (AAS) using a Fe hollow cathode lamp operating at 248.3 nm as the radiation source and the lamp current was set at

128 15 mA. The instrumental parameters were adjusted according to the manufacturer's
129 recommendations.

130

131 **2.7 Data Recorded and Statistical Analysis**

132 Data on protein and different mineral element content of fish samples were measured thrice
133 and the mean value was recorded for presentation. Finally, obtained data were analysed
134 statistically with the help of computer package M-STAT. Single factor ANOVA at $P \leq 0.05$
135 significant level was applied to compare concentrations of minerals and proteins among the
136 fish species.

137

138 **3. RESULTS AND DISCUSSION**

139

140 **3.1 Mineral Nutrients Status in Different Fish Species**

141

142 **3.1.1 Calcium (Ca)**

143 Calcium is an essential macronutrient element, which is important for bone formation.
144 Fishes, particularly small fishes are known to be a good source of this mineral [3-4, 30]. The
145 minimum and maximum concentration of Ca in fish samples were 0.16% in *shingi* and 2.00%
146 in *chela*, respectively where the mean of Ca concentration was 0.80% (Fig.1). The Ca
147 concentration in fishes was in the sequence of *chela* > *chingri* > *bele* > *meni* > *kaika* =
148 *bangna* > *punti* > *chanda* = *baim* = *guizza ayre* > *hiralo* > *kali baush* > *tengra* > *mola* =
149 *shingi*. The recommended daily intake (RDI) of calcium for adults is 1000-1300 mg [18]. For
150 the poor households in Bangladesh, cereals contributed 27.3% of the total Ca intake
151 followed by fish (21.8%), vegetables (14.0%), and milk and dairy products (10.6%). In case
152 of non-poor households, the contribution of cereals, fish, vegetables, and milk and dairy
153 products was 27.3%, 21.7%, 14.9%, and 10.6% of the total Ca intake, respectively [31].
154 Present study results revealed that in most cases higher amount of Ca was present in
155 different available fish species collected from the Brahmaputra River of Bangladesh
156 compared to past studies in different countries of the world and FAO measured
157 concentrations (Table 2). The higher calcium concentrations obtained in this study may be
158 attributed from higher levels of calcium in their feeds, and therefore more trophic transfer
159 and accumulation of this mineral in fish flesh. However, it is also suggested that small fish
160 with bones may be an important source of Ca in human diets [30].

161


162 **3.1.2 Magnesium (Mg)**

163 Magnesium content in different available fish species collected from the Brahmaputra River of
164 Bangladesh ranged from trace to 4.17% (Fig. 2). The highest amount of Mg was obtained
165 from *chanda* (4.17%) followed by *mola* (2.54%), *chela* (2.52%) and *chingri* (1.59%), which
166 may be a very good source of Mg for human nutrition. On the other hand, trace amount of
167 Mg was obtained from *baim* and *kaikka* fishes. The recommended daily intake of magnesium
168 for adults is 220-260 mg [18]. So, it can be inferred from this result that 11 fish species
169 (except *tengra*, *guizza ayre*, *baim* and *kaikka*) can contribute 100% of this requirement by
170 taking 100g fish flesh portion. Comparison to other past studies, the present work obtained
171 much higher magnesium content in most of fish species (Table 2). The concentration of Mg
172 in the muscles of perch from a Siberian pond (Russia) was 136 mg in 100g⁻¹ dry weight [32].
173 The mean value of Mg in fish samples ranged 29-41 mg 100g⁻¹ [33] and 36.4 mg 100g⁻¹
174 [22]. Similarly, present study results also revealed that out of 15 fish species, 12 samples
175 contained higher amount of Mg compared to past studies conducted in different countries of
176 the world (Fig. 2 and Table 2). This could be due to the difference of species, seasons, area
177 of catch, feed and many other physical and environmental conditions of the area.

178


Deleted: r

Deleted: r



181
182
183
184
185
186

Fig. 2. Calcium content (%) in different fish species collected from the Brahmaputra river of Bangladesh. Each value is the mean for three replicates, and vertical bars indicate the standard errors.



187
188
189
190

Fig. 3. Magnesium content (%) in different fish species collected from the Brahmaputra river of Bangladesh. Each value is the mean for three replicates, and vertical bars indicate the standard errors.

191
192
193

Table 2. Comparison of mineral concentrations range (mg 100g⁻¹) in fresh water fishes of this work with past studies in abroad.

Mineral element	Lagoon in USA ^a	River in Pakistan ^b	Lake in Ethiopia ^c	Lagoon in Botswana ^d	Lake in Poland ^e	River in Sudan ^f	Rivers in India ^g	Present study	FAO ^h conc. in fish muscles	RDA (mg/day) for Indians ⁱ
Ca	760-2200	33-1080	124-981	413-1290	53-103	107-588	220-2023	160-2000	19-881	600-1200
Mg	27-140	81-156	60-81	34-48	84-143	68-75	74-120	5-4170	4.5-452	540-1002
Na	36-400	85-163	163-210	86-145	148-328	180-280	87-107	170-410	30-134	1100-3300
K	nd	282-371	1121-1728	245-443	1429-2387	954-1210	24-132	320-3240	19-502	1875-5625
P	nd	969-1730	25-56	435-1375	1047-1261	727-935	nd	60-170	68-550	600-1200
S	nd	nd	nd	nd	nd	nd	nd	95-129	130-257	nd
N	nd	11987-12736	2397-2760	nd	nd	9568-12656	958-4198	2810-3880	2400-3200	nd
Fe	2.3-9.0	2.5-7.4	1.6-2.6	1.7-6.4	0.8-1.1	1.7-6.1	7.9-24.7	0.04-22.7	1.0-5.6	17-35


194 ^a(USA)= Moeller et al. [36]; ^b(Pakistan)= Jabeen et al. [37]; ^c(Ethiopia)= Teame et al. [38]; ^d(Botswana)= Mogobe et al. [24]; ^e(Poland)=
195 Luczyiska et al. [39]; ^f(Sudan)= Mohamed et al. [40]; ^g(India)= Romharsha and Sarojinalini [41]; ^h= FAO [34]; ⁱ= RDA [42]; nd= not
196 determined.

197 **3.1.3 Sodium (Na)**

198 Sodium is good for muscle functions and electrolyte balancing but it is not usually a problem
199 in mineral deficiencies as it is frequently used to salt food [21]. Sodium content in different
200 available fish species collected from the Brahmaputra River of Bangladesh varied from 0.17-
201 0.41% with a mean value of 0.26% (Fig. 3), which was more than twice as recommended by
202 FAO (30-134 mg 100g⁻¹) [34]. The highest amount of Na was obtained from *chingri* (0.41%)
203 followed by *bele* (0.33%), and *chanda* (0.32%), which are a good source of Na for human
204 nutrition. On the other hand, the lowest amount of Na was obtained from *baim* and *bangna*
205 fishes. The muscle tissues of three species of fishes contained 381 mg Na 100g⁻¹ collected
206 from the fresh water of Dhanmondi Lake in Bangladesh [35], which is comparable to the
207 present study results. However, the results were at the higher end compared to some
208 studies carried out past in freshwater fishes of other countries like USA (36-400 mg 100g⁻¹),
209 Pakistan (85-163 mg 100g⁻¹), Ethiopia (163-210 mg 100g⁻¹), Botswana (86-145 mg 100g⁻¹),
210 Sudan (180-280 mg 100g⁻¹), India (87-107 mg 100g⁻¹) and Poland (148-328 mg 100g⁻¹).
211 Furthermore, the Na contents obtained by this study were also higher compared to FAO
212 reported concentrations in fish muscles (Table 2).
213

Deleted:

Deleted: r



214


215 **Fig. 4. Sodium content (%) in different fish species collected from the Brahmaputra**
216 **river of Bangladesh. Each value is the mean for three replicates, and vertical**
217 **bars indicate the standard errors.**

218
219 **3.1.4 Potassium (K)**

220 Potassium plays also an important role like Na for muscle functions, transmission of
221 impulses in the nerves and sugar metabolism. The highest concentration of potassium
222 among different available fish species collected from the Brahmaputra River of Bangladesh
223 was obtained from *shingi* fish (3.24%) and the lowest was in *chanda* fish (0.32%) (Fig. 4).
224 The K concentration in fishes was in the sequence of *shingi* > *tengra* > *bangna* > *bele* > *kali*
225 *baush* > *hiralo* > *guizza ayre* > *meni* > *punti* > *kaika* > *chingri* > *baim* > *mola* > *chela* >
226 *chanda*. However, some studies carried out past in freshwater fishes in Turkey (321-441 mg
227 100g⁻¹) [21], China (301-402 mg 100g⁻¹) [43], Pakistan (282-371 mg 100g⁻¹), Botswana

Deleted: r

231 (245-443 mg 100g⁻¹) and India (24-132 mg 100g⁻¹) were at the lower end compared to this
 232 study results (Table 2). But the studies from Ethiopia, Sudan and Poland obtained much
 233 higher K concentration levels (1121-1728, 954-1210 and 1429-2387 mg 100g⁻¹,
 234 respectively) which is at par with the present study (Table 2). The recommended daily
 235 allowance (RDA) of K for males aged between 25-50 years is 800 mg [18]. So, the
 236 consumption of 100g of this river fish flesh will provide 40-100% of the potassium daily
 237 requirement, assuming cooking will not affect the quantity of the mineral.
 238




239
 240 **Fig. 5. Potassium content (%) in different fish species collected from the Brahmaputra**
 241 **river of Bangladesh. Each value is the mean for three replicates, and vertical**
 242 **bars indicate the standard errors.**

243 3.1.5 Phosphorus (P)

244 Phosphorus as phosphate is an essential nutrient involved in many physiological processes,
 245 such as the cell's energy cycle, regulation of the whole body acid-base balance, as a
 246 component of the cell structure (as phospholipids), in cell regulation and signaling, and as a
 247 major constituents of bones and teeth [44]. The maximum concentration of P in available fish
 248 species collected from the Brahmaputra River of Bangladesh was 0.17% in *bele* fish
 249 followed by *mola* (0.16%), *chela* (0.15%), *hiralo* (0.15%) and *chanda* (0.15%). On the other
 250 hand, the minimum P concentration was 0.06% in *baim* fish (Fig. 5). The P concentration
 251 range obtained in this work was within the FAO range of 68-550 mg 100g⁻¹ sample, but the
 252 range was lower compared to previous studies conducted in Pakistan (969-1730 mg
 253 100g⁻¹), Botswana (435-1375 mg 100g⁻¹), Poland (1047-1261mg 100g⁻¹) and Sudan (727-
 254 935 mg 100g⁻¹) for freshwater fishes. On the contrary, the range was higher compared to
 255 the past study conducted in Ethiopia (25-56 mg 100g⁻¹) for freshwater fishes (Table 2).
 256 However, this variation of P content in different fish samples could be due to the difference
 257 of species, seasons, area of catch and many other physical and environmental conditions of
 258 the area.
 259
 260

Deleted: r



262
263 **Fig. 6. Phosphorus content (%) in different fish species collected from the**
264 **Brahmaputra river of Bangladesh. Each value is the mean for three**
265 **replicates, and vertical bars indicate the standard errors.**
266

267 **3.1.6 Sulphur (S)**


268 The minimum S content (0.095%) in available fish species collected from the Brahmaputra
269 River of Bangladesh was obtained from *kali baush* fish, while the maximum (0.129%) was
270 found in *baim* fish (Fig. 6). Sulphur concentration range obtained in this study was at the
271 lower end of FAO reported range of 130-257 mg 100g⁻¹ sample (Table 2). Similarly, S
272 content in some important fish species of Bangladesh ranged from 160 to 300 mg 100g⁻¹
273 sample [45]. However, such variations in mineral concentrations in different fish samples
274 could be due to the difference of species, seasons, area of catch and many other physical
275 and environmental conditions of the area.
276

277 **3.1.7 Iron (Fe)**


278 Iron (Fe) is the most abundant metal in the human body. Body Fe content is approximately
279 3-4 g, which almost corresponds to a concentration of 40-50 mg of Fe per kilogram of body
280 weight [46]. Iron deficiency causes anaemia, which is one of the commonest mineral
281 deficiency diseases in Africa with 206 million people at risk [47]. The rich sources of dietary
282 Fe include red meat, liver, lentils, beans, peas, nuts, seeds, poultry, fish, seafood, leafy
283 vegetables, watercress, tofu, chickpeas, black-eyed peas, blackstrap molasses, fortified
284 bread, and fortified breakfast cereals [48]. The average concentration of Fe in different fish
285 samples collected from the Brahmaputra river was 44.79 mg kg⁻¹. The minimum and
286 maximum concentrations of Fe in fish samples were 0.40 (*baim*) and 226.92 (*mola*) mg kg⁻¹,
287 respectively (Fig. 7). The highest amount of Fe was obtained from *mola* followed by *chingri*
288 and *chela* that might be due to contribution of fish bone. It is worth mentioning here that
289 these three fish samples were processed for digestion along with exoskeleton and bone.
290 Furthermore, in most cases these fish species also consumed by the people along with
291 exoskeleton and bone. The mean Fe contents in muscle of red mullet ranged from 8.93-
292 49.30 mg kg⁻¹ (wet wt.), whereas levels in the liver of the same species ranged from 83.90-
293 889.00 mg kg⁻¹ (wet wt.). The differences between two tissues were not that significant in

Deleted: r

295 whiting and found 21.90-160.00 mg kg⁻¹ (wet wt.) in the muscle and 49.90-328.00 mg kg⁻¹
 296 (wet wt.) in the liver tissues [49].
 297



298
 299 **Fig. 7. Sulphur content (%) in different fish species collected from the Brahmaputra**
 300 **river of Bangladesh. Each value is the mean for three replicates, and vertical**
 301 **bars indicate the standard errors.**
 302
 303



304
 305 **Fig. 8. Iron content (%) in different fish species collected from the Brahmaputra**
 306 **river of Bangladesh. Each value is the mean for three replicates, and vertical**
 307 **bars indicate the standard errors.**

308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332

3.2 Nitrogen and Protein Status in Different Fish Species


The highest concentration of N among different available fish species collected from the Brahmaputra River of Bangladesh was obtained from *shingi* fish (3.88%) and the lowest was in *mola* fish (2.81%). The mean N concentration among the fish samples was 3.39%. The N content in fishes was in the sequence of *Shingi* > *kaikka* > *punti* > *baim* > *kali baush* > *bele* > *bangna* > *hiralo* > *tengra* > *meni* > *guizza ayre* > *chanda* > *chela* > *chingri* > *mola*. However, the dry matter content of different fish species collected from fresh water examined averaged $2.17 \pm 2.93\%$ nitrogen [50]. Nitrogen content in different available fish species collected from the Brahmaputra River was comparatively higher, which contributed to higher protein content.

Deleted: r

Deleted: r

Fish protein provides a good combination of amino acids which is highly suited to man's nutritional requirements and compares favourably with that provided by meat, milk and eggs. Fish protein showed a high biological value, comparable with that of milk, as shown by the similar values for apparent N absorption and N retention at each protein level [30]. In addition aquatic protein is highly digestible and rich in several peptides and essential amino acids that are limited in terrestrial meat proteins, as for example methionine and lysine [51]. The maximum amount of protein in available fish species collected from the Brahmaputra River of Bangladesh was 24.27% in *shingi* fish followed by *kaikka* (23.59%), *punti* (23.52%), *baim* (23.31%) and *kali baush* (23.19%). On the other hand, the minimum amount of protein was 17.53% in *mola* fish. The amount of protein in fish muscle is usually somewhere between 15-20%, but values lower than 15% or as high as 28% are occasionally met with in some species. However, the amount of protein obtained by this study was within the range.

Deleted: r



333
334
335
336
337

Fig. 9. Protein content (%) in different fish species collected from the Brahmaputra river of Bangladesh. Each value is the mean for three replicates, and vertical bars indicate the standard errors.

341 4. CONCLUSION

342

343 Present study provides data on major mineral nutrients viz. Ca, Mg, Na, K, P, S and Fe in
344 different fish samples collected from the Brahmaputra River of Bangladesh. Study results
345 revealed comparatively higher amount of Ca, Mg, Na and K in different available fish species
346 collected from the river compared to some other studies carried out in different countries of
347 the world for freshwater fishes. The study results found that 3, 11, 12 fish species can
348 contribute 100% of Ca, Mg and K requirement by taking 100g fish flesh, respectively. On the
349 other hand, average consumption of 100g of this river fish flesh will provide 80-100% of the
350 daily requirement of Ca, Mg and K, assuming cooking will not affect the quantity of the
351 minerals. Thus the study inferred that fishes of the Brahmaputra River are a good source of
352 protein and minerals, which contributes in nutrition to the local people of the country. The
353 information generated from this study could be used as a baseline data for developing food
354 composition database of the country. Finally, the study concluded that the variation of major
355 mineral nutrient contents in different fish samples collected from the Brahmaputra River due
356 to the difference of species, age, area of catch, feed, quality of water and many other
357 physical and environmental conditions of the area.

Deleted: r

Deleted: river

Deleted: river

358

359 COMPETING INTERESTS

360

361 Authors have declared that no competing interests exist.

362

363 AUTHORS' CONTRIBUTIONS

364

365 'Author MSAE' collected samples, performed analysis and wrote the first draft of the
366 manuscript. 'Authors HMZ and QFQ' designed the study, supervise the work, performed the
367 statistical analysis and corrected the final draft of the manuscript, 'Author MSR' helped in
368 manuscript preparation. All authors read and approved the final manuscript."

369

370 REFERENCES

371

- 372 1. Pirestani S, Sahari A, Barzegar M, Seyfabadi SJ. Chemical compositions and minerals
373 of some commercially important fish species from the South Caspian Sea. *Int Food Res*
374 *J.* 2009;16:39-44.
- 375 2. Kawarazuka N, Bene C. The potential role of small fish species in improving
376 micronutrient deficiencies in developing countries: Building evidence. *Public Health Nutr.*
377 2011;14:1927-1938. DOI: 10.1017/S1368980011000814
- 378 3. Fawole OO, Ogundiran MA, Ayandiran TA, Lagunju OF. Proximate and mineral
379 composition of some selected fresh water fishes in Nigeria. *Internet J Food Safety,*
380 2007;9:52-55.
- 381 4. Roos N, Wahab AM, Chamnan C, Thilsted SH. The role of fish in Food-based strategies
382 to combat vitamin A and Mineral Deficiencies in developing countries. *J Nutr.*
383 2007;137:1106-1109. DOI: 10.1093/jn/137.4.1106
- 384 5. Rana M, Bakali B, Mia MY, Zakir HM. Physico-chemical properties of effluents
385 discharged from different industries of Gazipur, Bangladesh and its suitability for
386 agricultural land. *J Environ Sci Natural Resources,* 2014;7(1):157-167.
- 387 6. Zakir HM, Islam MM, Hossain MS. Impact of urbanization and industrialization on
388 irrigation water quality of a canal- a case study of Tongi canal, Bangladesh. *Adv Environ*
389 *Res.* 2016;5(2):109-123. DOI: 10.12989/aer.2016.5.2.109.

- 393 7. Hossain MB, Islam MN, Alam MS, Zakir HM. Industrialisation scenario at Sreepur of
394 Gazipur, Bangladesh and physic-chemical properties of wastewater discharged from
395 industries. *Asian J Environ Ecol*. 2019;9(4):1-14. DOI: 10.9734/AJEE/2019/v9i430103
396 8. Hossain MA, Zakir HM, Kumar D, Alam MS. Quality and metallic pollution level in
397 surface waters of an urban industrialized city: a case study of Chittagong city,
398 Bangladesh. *J Ind Safety Engg*. 2017;4(2):9-18.
399 9. Arifa Akter, Zakir HM, Atiqur Rahman, Rumana Yesmeen, Rahman MS. Appraisal of
400 surface water quality for irrigation collected from Sadar upazila of Jamalpur district,
401 Bangladesh. *Arc Agric Environ Sci*. 2018;3(3):216-225. DOI:
402 10.26832/24566632.2018.030302
403 10. Zakir HM, Rahman MM, Rahman A, Ahmed I, Hossain MA. Heavy metals and major
404 ionic pollution assessment in waters of midstream of the river Karatoa in Bangladesh. *J*
405 *Environ Sci and Natural Resources*. 2012;5(2):149-160.
406 11. Zakir HM, Sattar MA, Quadir QF. Cadmium pollution and irrigation water quality
407 assessment of an urban river: a case study of the Mayur river, Khulna, Bangladesh. *J*
408 *Chem Bio Phy Sci*. 2015;5(2):2133-2149.
409 12. Hossain MS, Zakir HM, Rahman MS, Islam MM. Toxic metallic contamination in
410 wastewater of some industrial areas of Mymensingh town, Bangladesh. *Adv Archit City*
411 *Environ*. 2015;1(3):7-13.
412 13. Zakir HM, Islam MM, Hossain MS. Heavy metal contents in sediments of an urban
413 industrialized area- a case study of Tongi canal, Bangladesh. *Asian J Water Environ*
414 *Pollut*. 2017;14(1):59-68. DOI 10.3233/AJW-170007
415 14. Yesmeen R, Zakir HM, Alam MS, Mallick S. Heavy metal and major ionic contamination
416 level in effluents, surface and groundwater of an urban industrialised city: A case study
417 of Rangpur city, Bangladesh. *Asian J Chem Sci*. 2018;5(1):1-16. DOI:
418 10.9734/AJOCS/2018/45061
419 15. Bhandari S, Banjara MR. Micronutrients deficiency, a hidden hunger in Nepal:
420 prevalence, causes, consequences and solutions. *Int Scholarly Res Notices*,
421 2015;2015:1-9. DOI: 10.1155/2015/276469
422 16. Komatsu F, Kagawa Y, Kawabata T, Kaneko Y, Kudoh H, Purvee B, Otgon J,
423 Chimedregzen U. Influence of essential trace minerals and micronutrient insufficiencies
424 on harmful metal overload in a Mongolian patient with multiple sclerosis. *Curr Aging Sci*.
425 2012;5:115-125. DOI: 10.2174/1874609811205020112
426 17. Hsieh BT, Chang CY, Chang YC, Cheng KY. Relationship between the level of essential
427 metal elements and in human hair and coronary heart disease. *J Radioanal Nucl Chem*.
428 2011;290:165-169. DOI: 10.1007/s10967-011-1174-z
429 18. FAO/WHO. Human Vitamin and Mineral Requirements. Report of a joint FAO/WHO
430 expert consultation on human vitamin and mineral requirements. Food and Nutrition
431 Division, FAO, Rome; 2001.
432 19. Dana JD, Hurlbut CS, Klein C. *Manual of Mineralogy*. 2nd ed., John Wiley and Sons Inc.,
433 New York, USA. 1985.
434 20. Waterman JJ. *The Composition of Fish*. Ministry of Agriculture, Fisheries and Food.
435 Torry Advisory Note No. 67. Torry Research Station Edingburgh. 1980.
436 21. Alas A, Ozcan MM, Harmankaya M. Mineral contents of head, caudal, central fleshy
437 part, and spinal columns of some fishes. *Environ Monit Assess*. 2014;186:889-894. DOI:
438 10.1007/s10661-013-3429-3
439 22. Martinez-Valverde I, Periago MJ, Santaella M, Ros G. The content and nutritional
440 significance of minerals on fish flesh in the presence and absence of bone. *Food Chem*.
441 2000;71:503-509. DOI: 10.1016/S0308-8146(00)00197-7
442 23. Daczowska-Kozon E, Sun-pan B (2011). *Environmental Effects on Seafood Availability,*
443 *Safety and Quality*. 1st ed., CRC Press, Taylor and Francis; 2011.

- 444 24. Mogobe O, Mosepele K, Masamba WRL. Essential mineral content of common fish
445 species in Chanoga, Okavango Delta, Botswana. *Afr J Food Sci.* 2015;9(9):480-486.
446 DOI: 10.5897/AJFS2015.1307
- 447 25. Rebole A, Velasco S, Rodriguez ML, Trevino J, Alzueta C, Tejedor JL, Ortiz LT. Nutrient
448 content in the muscle and skin of fillets from farmed rainbow trout (*Oncorhynchus*
449 *mykiss*). *Food Chem.* 2015;174:614-620. DOI: 10.1016/j.foodchem.2014.11.072
- 450 26. Galib, SM. Fish fauna of the Brahmaputra River, Bangladesh: richness, threats and
451 conservation needs. *J Fisheries* 2015;3(3):285-292. DOI: 10.17017/jfish.v3i3.2015.120
- 452 27. Sarma JN. An Overview of the Brahmaputra River System. *In: Singh VP, Sharma N,*
453 *Ojha CSP (eds). The Brahmaputra Basin Water Resources. Water Science and*
454 *Technology Library, vol. 47. Springer, Dordrecht; 2004. DOI: 10.1007/978-94-017-0540-*
455 *0_5*
- 456 28. Singh D, Chhonkar PK, Pandey RN. *Soil, Plant and Water Analysis: A Method Manual.*
457 IARI, New Delhi. India; 1999.
- 458 29. Piper CS. *Soil and Plant Analysis.* The University of Adelaide Press, Adelaide, Australia.
459 1950. 368 pp.
- 460 30. Larsen T, Thilsted HS, Kongsbak K, Hansen M. Whole fish as a rich calcium source. *Br J*
461 *Nutr.* 2000;83:191-196. DOI: 10.1017/S0007114500000246
- 462 31. Islam MR. Consumption of unsafe foods: Evidence from heavy metal, mineral and trace
463 element contamination. The National Food Policy Capacity Strengthening Programme
464 (NFPCSP) sponsored project, Dept. of Soil Science, BAU, Mymensingh. 2013.
- 465 32. Gladyshev MI, Gribovskaya IV, Moskvicheva AV, Muchkina EY, Chuprov SM, Ivanova
466 EA. Content of metals in compartments of ecosystem of a Siberian pond. *Arch Environ*
467 *Contam Toxicol.* 2001;41:157-162. DOI: 10.1007/s002440010233
- 468 33. Adeniyi SA, Orjiekwe CL, Ehiagbonare JE, Josiah SJ. Nutritional composition of three
469 different fishes (*Claria gariepinus*, *Malapterurus electricus* and *Tilapia guineensis*). *Pak J*
470 *Nutr.* 2012;11:793-797. DOI: 10.3923/pjn.2012.891.895
- 471 34. FAO. FAO in partnership with Support unit for International Fisheries and Aquatic
472 Research (SIFAR). FAO, Rome; 2001. <http://www.fao.org/3/x5916e00.htm>
- 473 35. Begum A, Amin MN, Kaneco S, Ohta K. Selected elemental composition of the muscle
474 tissue of three species of fish, *Tilapia nilotica*, *Cirrhina mrigala* and *Clarius batrachus*
475 from the fresh water Dhanmondi lake in Bangladesh. *Food Chem.* 2005;93:439-443.
476 DOI: 10.1016/j.foodchem.2004.10.021
- 477 36. Moeller A, MacNeil SD, Ambrose RF, Hee SSQ. Elements in fish of Malibu Creek and
478 Malibu Lagoon near Los Angeles, California. *Mar Pollut Bull.* 2003;46:424-429. DOI:
479 10.1016/S0025-326X(02)00466-6
- 480 37. Jabeen F, Noureen A, Hussain SM, Chaudhry AS, Irfan M, Shakeel M, Shabbir S,
481 Yaqub S, Ahmad S, Shaheen T. Chemical and mineral composition of *Cyprinus carpio*,
482 *Labeo rohita* and *Wallago attu* inhabiting river Indus in Mianwali district. *Int J Biosci.*
483 2015;6(5):333-342. DOI: 10.12692/ijb/6.5.333-342
- 484 38. Teame T, Natarajan P, Tesfay Z. Proximate and mineral composition of some
485 commercially important fish species of tekeze reservoir and lake Hashenge, Ethiopia. *Int*
486 *J Fish Aquat Stud.* 2016;4(1):160-164.
- 487 39. Luczynska J, Tonska E, Luczynski J. Essential mineral components in muscles of six
488 freshwater fish from the Mazurian Great Lakes (northeastern Poland). *Arch Pol Fish.*
489 2009;17:171-178. DOI:10.2478/v10086-009-0015-y
- 490 40. Mohamed EAH, Al-Maqbaly R, Mansour MH. Proximate Composition, amino acid and
491 mineral contents of five commercial Nile fishes in Sudan. *Afr J Food Sci.* 2010;10:650-
492 654. <http://www.academicjournals.org/ajfs>

- 493 41. Romharsha H, Sarojnalini C. Proximate composition, total amino acids and essential
494 mineral elements of some cyprinid fishes of Manipur, India. *Curr Res Nutr Food Sci.*
495 2018;6(1):157-164. DOI: 10.12944/CRNFSJ.6.1.18
- 496 42. RDA. Nutrient Requirements and Recommended Dietary Allowances for Indians. A
497 Report of the Expert Group of the Indian Council of Medical Research 2009. National
498 Institute of Nutrition. Hyderabad-500 604; 2010.
- 499 43. Tao NP, Wang LY, Gong X, Liu Y. Comparison of nutritional composition of farmed
500 puffer fish muscles among *Fugu obsurus*, *Fugu flavidus* and *Fugu rubripes*. *J Food*
501 *Comp Anal.* 2012;28:40-45. DOI: 10.1016/j.jfca.2012.06.004
- 502 44. EFSA (European Food Safety Authority). Tolerable Upper Intake Levels for Vitamins and
503 Minerals. Scientific committee on food; scientific panel on dietetic products, nutrition and
504 allergies. 2006.
- 505 45. Bogard JR, Thilsted SH, Marks GC, Wahab MA, Hossain MAR, Jakobsen J, Stangoulis
506 J. Nutrient composition of important fish species in Bangladesh and potential
507 contribution to recommended nutrient intakes. *J Food Comp Anal.* 2015;42:120-133.
508 DOI: 10.1016/j.jfca.2015.03.002
- 509 46. Crichton, RD, Danielson BG, Geisser P. Iron Therapy with Special Emphasis on
510 Intravenous Administration. 3rd ed., UNI-MED Verlag AG; 2006.
- 511 47. Latham MC. Human Nutrition in Developing World. Rome: FAO Food and Nutrition
512 Series No. 29, FAO, Rome; 1997.
- 513 48. Roger M. The Minerals You Need. Safe Goods Publishing, 561 Shunpike Road,
514 Sheffield, MA 01257, USA; 2011.
- 515 49. Tepe Y, Türkmen M, Türkmen A. Assessment of heavy metals in two commercial fish
516 species of four Turkish seas. *Environ Monit Assess.* 2008;146:277-284. DOI:
517 10.1007/s10661-007-0079-3.
- 518 50. Schreckenbach K, Knosche R, Ebert K. Nutrient and energy content of freshwater
519 fishes. *J Appl Ichthyol.* 2001;17(3):142-144. DOI: 10.1111/j.1439-0426.2001.00295.x
- 520 51. Tacon AGJ, Metian M. Fish matters: importance of aquatic foods in human nutrition and
521 global food supply. *Rev Fisher Sci.* 2013;21:22-38. DOI:
522 10.1080/10641262.2012.753405
- 523
524
525