

1  
2 **Original Research Article**

3  
4 **Real time PCR based quantification of Banana**  
5 **Bunchy Top Virus (BBTV) titre in banana cv.**  
6 **Grand Naine (*Musa acuminata*)**  
7

8  
9  
10  
11  
12  
13 **ABSTRACT**  
14

Banana Bunchy Top Disease (BBTD) is one of the most severe viral diseases affecting major banana growing belts in India. Banana Bunchy Top Virus (BBTV) is transmitted by a black banana aphid (*Pentalonia nigronervosa* L Coquerel) in a persistent manner. BBTV virions are limited to the phloem tissue of banana resulting in low titre in banana. A reliable method to quantify the BBTV in the banana will be useful for monitoring the insect vector mediated BBTV transmission in banana, an essential requirement for characterizing the transgenic banana transformed for BBTV resistance. A protocol for real time PCR based absolute quantification of BBTV is reported in the present study. The partial BBTV Coat protein gene (459 bp) was isolated, cloned into a plasmid vector and used to construct a standard curve using an SYBR green-based assay with known copies of BBTV CP gene. Using the standard curve, BBTV viral loads was estimated in BBTV infected symptomatic and asymptomatic leaf samples of banana cultivar Grand Naine through SYBR green-based quantitative Polymerase Chain Reaction (qPCR). The study proves that a higher viral titre is associated with BBTV disease symptoms appearance, whereas the low titre results in asymptomatic plants.

15  
16 *Keywords:* [Grand Naine, BBTV, Coat protein, PCR, Real time quantification]  
17

18 **1. INTRODUCTION**  
19

20 Banana bunchy top disease caused by a banana bunchy top virus (BBTV, family  
21 *Nanoviridae*, genus *Babuvirus*) is one of the significant viruses affecting banana plants  
22 leading to a 100 % loss in yield and has been recorded in all countries cultivating banana  
23 across the globe. BBTD is transmitted by the aphid *Pentalonia nigronervosa* [11] in a  
24 persistent, circulative, non-propagative manner; the virus enters the haemocoel and the  
25 aphids retain the virus throughout their lives without passing it on to their offspring [7].

26 *Pentalonia nigronervosa* was first described by [3] from banana. The BBTV is an ssDNA  
27 virus and comprises of six genome components and are termed as DNA-R, DNAU3, DNA-S,  
28 DNA-M, DNA-C, and DNA-N [10]. Based on the DNA-R sequences, the BBTV isolates were  
29 grouped into two: Asian and South Pacific groups [9]. The Asian group comprises of the  
30 isolates from the Philippines, Taiwan, China, Japan, Indonesia, and Vietnam. The South  
31 Pacific group includes India, Pakistan, Egypt, Australia, Burundi, Tonga, Myanmar, Fiji, and  
32 USA-Hawaii and isolates from Africa [15].

33 Earlier Polymerase Chain Reaction (PCR)-based method was employed for the detection of  
34 BBTV infection in banana [6, 12]. Recent methods demonstrate the quantification of BBTV  
35 through real-time PCR using SYBR Green or TaqMan chemistry [2, 5, 8 and 15]. Aphid  
36 mediated transmission is a standard method for the BBTV virus infection in banana.  
37 However, the BBTV transmission experiments require a suitable tool to monitor the virus  
38 accumulation in banana plants. Real time PCR is a sensitive method to quantify the viral  
39 load during the insect mediated transmission study in banana. The real time PCR based  
40 developed will be useful for the BBTV transmission study and also to characterize the  
41 transgenic banana plants engineered for BBTV resistance.

42

## 43 **2. MATERIAL AND METHODS**

44

### 45 **2.1 Source of plant material and virus inoculum**

46 BBTV infected samples of banana cv. Grand Naine were collected from the orchard of Tamil  
47 Nadu Agricultural University, Coimbatore, Tamil Nadu and confirmed the presence of BBTV  
48 infection through PCR. The tissue culture plants of banana cultivar Grand Naine obtained  
49 from Sunglow Biotech, Coimbatore were used throughout the study. Healthy banana adult  
50 aphids of *P. nigronervosa* were obtained from Dr. R. Selvarajan, NRCB, Trichy and  
51 multiplied in the tissue culture banana plants of cv. Grand Naine. The new aphid population  
52 was continuously obtained by culturing in new tissue culture plants.

53 The transmission studies were conducted by releasing the virus-free, same-sized, 8-10 days  
54 old aphids (apterae) to BBTV- infected plant for a period of 24 h to acquire the virus in an  
55 insect-proof chamber. After the acquisition of the virus by the aphids, 20 viruliferous aphids  
56 were allowed to feed on the virus- free tissue culture banana plants of cv. Grand Naine for a  
57 period of 48 h to transmit the virus. The banana plants were sprayed with 0.02 %  
58 Imidacloprid to kill the aphids.

### 59 **2.2 Isolation of BBTV coat protein gene**

60 The total DNA was extracted from BBTV infected leaf tissues of Grand Naine using the  
61 CTAB method [4]. One hundred milligram of leaf tissues was homogenised with one ml of  
62 CTAB buffer ([10 mM Tris-HCl, pH 8.0, 1.4 M NaCl, 20 mM EDTA, 2 % CTAB, 0.1 % (v/v)  $\beta$ -  
63 Mercaptoethanol and 2 (v/w) %PVP] to isolate the total DNA. Then DNA was precipitated  
64 with ice cold iso-propanol and washed with 70 % ethanol to remove the salts. The pellet  
65 obtained was air-dried and dissolved with 100  $\mu$ l of sterile water and stored at -20  $^{\circ}$ C for  
66 further analysis. The DNA samples were quantified using Nanodrop spectrophotometer ND-  
67 1000. The DNA samples were resolved on a 1.0 % agarose gel.


68 BBTV coat protein specific forward primer (5' TCCGAAGAAATCCATCAAGA 3') and reverse  
69 primer (5'CCAGAACTA CAATAGAATGCCAAA 3') were used for amplifying the coat protein  
70 gene. PCR condition for amplification of Coat protein gene of BBTV was as follows: Initial  
71 denaturation at 94 °C for 5 min followed by 35 cycles of Denaturation at 94 °C for 1 min, 60  
72 °C annealing for 30 sec, extension at 72 °C for 1 min and final extension was given at 72 °C  
73 for 5 min. PCR reactions were performed in Thermal Cycler (Eppendorf, Germany) in a final  
74 volume of 20 µl. The PCR reaction mixture contained 1 µl of banana total DNA (50 ng/µl),  
75 2.0 µl of 10X PCR buffer (10 mM Tris-HCl pH 9.0, 50 mM KCl and 1.5 mM MgCl<sub>2</sub>), 0.5 µl of  
76 100 mM dNTPs, 1.0 µl each of 10 µM respective forward and reverse primers, 0.3 µl of 1  
77 unit Taq DNA polymerase (TaKaRa Bio USA, Inc.) and 14.2 µl of sterile distilled water. The  
78 PCR amplified product was resolved on 1% Agarose gel. The amplified products were  
79 visualized on a UV transilluminator and documented in a gel documentation system  
80 (Syngene, UK).

### 81 **2.3 Cloning of partial BBTV Coat protein gene**

82 A 459 bp PCR amplified CP gene of BBTV was purified using PCR clean up kit (Macherey-  
83 Nagel GmbH & Co. KG, Germany) according to the manufacturer's instruction. The purified  
84 product was ligated and cloned using pJET1.2/blunt PCR Cloning Kit (Thermo Scientific Inc.,  
85 USA) and transformed using *E. coli* DH5α competent cells plated on Luria and Bertini agar  
86 media containing ampicillin (100 mg/L). Plates were incubated in 37 °C overnight. The  
87 transformed clones were analysed for the presence of recombinant clones through PCR  
88 followed by DNA sequencing (Agrigenomes Pvt. Ltd., Cochin, India).

### 89 **2.4 Standard curve for absolute quantification BBTV DNA through qPCR assay**

90 The purified 2.9 kb plasmid DNA containing a partial coat protein gene was used to prepare  
91 a standard curve for absolute quantification of the BBTV DNA-S component as described by  
92 [16]. The plasmid was serially diluted to obtain a standard series up to 10<sup>7</sup> with each step  
93 differing by 10- fold. The copy number of the DNA in each dilution was calculated with the  
94 following formula  $Y \text{ molecules} = X \text{ g/}\mu\text{l DNA} \times 6.022 \times 10^{23} / (\text{Base pair of recombinant plasmid} \times$   
95  $660)$ . qPCR was performed using these diluted DNA with three replicates. After the assay,  
96 the values of threshold cycles (C<sub>q</sub>) were obtained. A standard curve was established by  
97 plotting the C<sub>q</sub> value on the Y-axis and natural log of concentration (copies/µl) on the X-axis,  
98 and a regression equation was obtained and coefficient of determination (R<sup>2</sup>) were achieved  
99 (Fig. 2).



100

101 **Fig. 2. The standard curve obtained for BBTV DNA-S coat protein quantification. The**  
 102 **X-axis represents the number of DNA copies, while the Y-axis represents the Cq**  
 103 **value.**

#### 104 **2.5 Quantification of viral load through qPCR assay in BBTV infected plants**

105 Banana plant DNA isolated from three symptomatic and non-symptomatic Grand Naine  
 106 infected plants (120 dpi) and each sample was tested in triplicate using qPCR. DNA isolated  
 107 from leaf tissue samples from wild-type virus-free Grand Naine plants were used as negative  
 108 control and the cloned CP plasmid was used as a positive control. Equal quantity of banana  
 109 genomic DNA (25 ng) was taken for the absolute quantification of BBTV by qPCR using  
 110 SYBR green chemistry as described by [1] and reactions were carried out in a Biorad® CFX  
 111 connect Real time PCR system (Biorad, USA). The designed Real-time PCR primers, BBTV-  
 112 RT-CP-F (5'TCAACCAGCCGACAACCTGT3') and RT-CP-R  
 113 (5'TGTCCCTGTTGCGACTCCTG3') were used to amplify 116 bp of coat protein gene of  
 114 BBTV. The standard amplification profile was 94 °C for 4 min and 40 cycles of 94 °C for 15 s  
 115 and 60 °C for 1 min and followed by a melt curve analysis was performed by allowing the  
 116 reactions at 94 °C for 15 s and 60 °C for 1 min. The qPCR cycle provided the melt curve for  
 117 each sample, for assessing the specificity of amplification. For each PCR reaction, the  
 118 samples were taken in three replicates.

### 119 **3. RESULTS AND DISCUSSION**

#### 120 **3.1 Isolation of BBTV coat protein gene from infected banana samples**

121 Genomic DNA samples were isolated from BBTV infected Grand Naine banana collected  
 122 from Coimbatore, Tamil Nadu. PCR amplification of partial length BBTV DNA-S was  
 123 observed in all five samples of Grand Naine (Fig. 1).

124

125


126

127

128

129

130



131

**Fig. 1. PCR detection of BBTV in Grand Naine banana leaf samples**

132

Lane L- 1 Kb ladder; 1 and 2- healthy Grand Naine samples; 3 to 7- BBTV infected Grand Naine samples

133

134

### 3.2 Cloning of BBTV coat protein gene

135

The BBTV coat protein gene was amplified by PCR using gene specific primer BBTV-CPF and BBTV-CPR. The PCR amplified BBTV coat protein gene (459 bp) was cloned into pJET1.2/ blunt vector. The transformed clones were confirmed by DNA sequencing (Agrigenome, Kerala, India)

136

137

138

139

### 3.3 Standard curve for quantification of BBTV DNA in the infected banana samples

140

141

142

143

144

145

146

147


148

149

150

Absolute quantification of viral DNA is done using the standard curve made with the cloned viral gene with known copy number. Melt curve analysis confirmed the presence of specific real time PCR amplified product (116 bp). All the PCR products melt between 75 °C and 83.5 °C which indicates the breakdown of only one PCR product (Fig. 3). The Cq values obtained for each dilution of the plasmid were used for regression analysis taking copy number and Cq value. The mean squares of the coefficient of determination ( $R^2$ ) value was above 0.995 which indicates good linearity between the initial copy numbers and the fluorescence values (Cq values) for obtaining a high-quality standard curve for absolute quantification [16]. BBTV coat protein gene is an ideal target gene for the establishment of standard curve. Earlier studies for BBTV DNA quantification also used the CP gene for quantification [8].

151



152 **Fig. 3. The melt curve of 116 bp BBTV CP gene product amplified in the real time PCR**  
 153 **assay. The X-axis represents the temperature value and the Y-axis represents the**  
 154 **value of the fluorescence.**

155 **3.4 Viral titre in asymptomatic and symptomatic BBTV infected plants of cv. Grand**  
 156 **Naine**

157 The results revealed that viral titre was higher in the symptomatic plant compared to non-  
 158 symptomatic plants (Table 1). Equal quantity (25 ng/μl) of the total DNA of the banana  
 159 samples were taken for real time PCR analysis to ensure that BBTV DNA. This was  
 160 achieved by the quantification of DNA concentration in Nanodrop and diluted to get a DNA  
 161 concentration of 25 ng per μl, followed by further confirmation in Nanodrop. Based on qPCR  
 162 results, the viral load in symptomatic plants was higher compared to non- symptomatic. In  
 163 the symptomatic plants, the viral load ranges from  $1.4 \times 10^8$  to  $7.7 \times 10^7$ , whereas in the  
 164 non-symptomatic plants it ranges from  $1.1 \times 10^5$  to  $2.3 \times 10^5$ . Even though good correlation  
 165 exists among the analyzed plants and disease symptoms, one plant in each show's  
 166 deviation, which may be associated with an experimental error. Previous reports in BBTV  
 167 DNA quantification also show real time PCR is an effective and simple method for  
 168 quantification of viral load [2, 15]. It clearly proves that the symptomatic plants harbour more  
 169 viral load which is the reason for symptom expression. The viral titre plays a major role in  
 170 symptom expression [8] and the delay in symptom expression in non-symptomatic plants  
 171 may be due to unequal transmission of aphids during the incidence [13]. The absolute  
 172 amount of each BBTV DNA component were measured by real- time PCR [1,17]. qPCR  
 173 results will invoke the importance of latency of BBTV in non symptomatic plants. For various  
 174 studies on viral accumulation in plants, real time PCR provides a confirmation protocol to  
 175 study the quantity of virus for transmission studies.

176 **Table 1. Quantification data of qPCR analysis between symptomatic and non-**  
 177 **symptomatic BBTV infected plants of banana cv. Grand Naine**

<b>Target gene</b>	<b>Sample type</b>	<b>Cq value</b>	<b>Cq SEM</b>	<b>Copy number of BBTV coat protein*</b>
CP	Positive control	18.97	0.01790	$2.7 \times 10^6$
CP	Negative control	25.81	0.06283	$4.9 \times 10^2$
CP	Non- symptomatic -H1	24.03	0.81199	$1.1 \times 10^5$
CP	Non- symptomatic -H2	23.66	0.21572	$1.3 \times 10^5$
CP	Non- symptomatic -H3	22.79	0.35354	$2.3 \times 10^5$
CP	Symptomatic - I1	13.79	0.08139	$7.7 \times 10^7$
CP	Symptomatic - I2	15.33	0.14517	$2.8 \times 10^7$

CP	Symptomatic - I3	12.78	0.07265	1.4 x10 <sup>8</sup>
CP	NTC	0.00	0.00	0.00

178 \*Copy numbers have been calculated using the formula.

179 Copy No, 1µl of the sample=  $\frac{X \text{ g/}\mu\text{l DNA} \times 6.022 \times 10^{23}}$

180 (base pair of recombinant plasmid x 660)

#### 181 4. CONCLUSION

182

183 Real time is a rapid and sensitive method for the quantification of BBTV DNA in banana.  
 184 During BBTV transmission through banana aphids, real time can be employed as a suitable  
 185 tool for the accurate quantification of the BBTV DNA. The study confirms that BBTV disease  
 186 symptoms occur in banana at higher viral load and absence of symptom at lower viral load.  
 187 There exists good correlation with the BBTV viral load and disease symptom in banana. The  
 188 protocol described in this study can also be employed for BBTV DNA quantification of  
 189 transgenic banana developed for BBTV resistance.

190

#### 191 REFERENCES

192

193

194 1. Bernard Ouma. Development of Banana bunchy top virus resistance in bananas: RNAi  
 195 approach. Ph. D. thesis. Queensland University of Technology. 2016.

196

197 2. Chen Y, Hu X. High-throughput detection of banana bunchy top virus in banana plants  
 198 and aphids using real-time TaqMan® PCR. Journal of virological methods. 2013;193  
 199 (1):177-183.

200 3. Coquerel C. Observations entomologiques sur divers insectes recueillis a Madagascar.  
 201 Annales de la Societe entomologiques de France. 1859; 3(7), 239-260.

202 4. Doyle JJ, Doyle JL. Isolation of plant DNA from fresh tissue. Focus. 1990; 12:13.

203 5. Fu HC, Hu JM, Hung TH, Su HJ, Yeh HH. Unusual events involved in Banana bunchy top  
 204 virus strain evolution. Phytopathology. 2009; 99 (7):812-822.

205

206 6. Furuya N, Somowiyarjo S, Natsuaki KT. Virus detection from local banana cultivars and  
 207 the first molecular characterization of Banana bunchy top virus in Indonesia. J Agri Sci  
 208 Tokyo. 2004; 49(3):75–81.

209 7. Hu JS, Wang M, Sether D, Xie W, Leonhardt KW. Use of polymerase chain reaction  
 210 (PCR) to study transmission of banana bunchy top virus by the banana aphid (*Pentalonia*  
 211 *nigronevosa*). Annals of Applied Biology. 1996; 128(1): 55-64.

212 8. Jebakumar RM, Balasubramanian V, Selvarajan R. Virus titre determines the efficiency of  
 213 *Pentalonia nigronevosa* (Aphididae: Hemiptera) to transmit banana bunchy top virus. Virus  
 214 Disease. 2018; 29 (4):499-505.

- 215 9. Karan M, Harding RM, Dale JL. Evidence for two groups of banana bunchy top virus  
216 isolates. *Journal of General Virology*. 1994; 75 (12):3541-3546.
- 217 10. King AMQ, Adams MJ, Lefkowitz EJ. *Virus taxonomy: classification and nomenclature of*  
218 *viruses: ninth report of the international committee on taxonomy of viruses*. Springer, Wein.  
219 2011.
- 220 11. Magee, C.J. Investigation on the bunchy top disease of bananas. *Bulletin of the Council*  
221 *for Scientific and Industrial Research in Australia*. 1927; 30.
- 222 12. Mansoor S, Qazi J, Amin I, Khatri A, Khan IA, Raza S, et al. A PCR-based method, with  
223 internal control, for the detection of Banana bunchy top virus in banana. *Molecular*  
224 *biotechnology*. 2005; 30 (2):167-169.
- 225 13. Selvarajan R, Balasubramanian V, Anuradha C. Population Structure and Diversity of  
226 Banana Bunchy Top Virus and Banana Bract Mosaic Virus. In *Plant Viruses*. 2018; 149-170.  
227 CRC Press.
- 228 14. Stainton D, Martin DP, Muhire BM, Lolohea S, Halafihi MI, Lepoint P, et al. The global  
229 distribution of Banana bunchy top virus reveals little evidence for frequent recent, human-  
230 mediated long-distance dispersal events. *Virus evolution*. 2015;1 (1).
- 231 15. Watanabe S, Greenwell A, Bressan A. Localization, concentration, and transmission  
232 efficiency of Banana bunchy top virus in four asexual lineages of *Pentalonia aphids*. *Viruses*  
233 2013; 5 (2):758-776.
- 234 16. Xue B, Guo J, Que Y, Fu Z, Wu L, Xu L. Selection of suitable endogenous reference  
235 genes for relative copy number detection in sugarcane. *International Journal of Molecular*  
236 *Sciences*. 2014;15(5): 8846-8862.
- 237 17. Yu NT, Xie HM, Zhang YL, Wang JH, Xiong Z, Liu ZX. Independent modulation of  
238 individual genomic component transcription and a cis- acting element related to high  
239 transcriptional activity in a multipartite DNA virus. *BMC genomics*. 2019; 20: 573.