

Original Research Article

Pattern and Socio-Demographic Determinants of Gestational Age at Antenatal Booking at the Rivers State University Teaching Hospital, Nigeria: a two-year review.

Abstract:

Background: Antenatal care is one of the pillars of the SAFE Motherhood Initiative aimed at preventing adverse pregnancy outcome. Early initiation provides an opportunity for optimum utilization of this care with improved maternal and fetal outcomes.

Objective: To determine the pattern and Socio-demographic determinants of gestational age at antenatal booking at the Rivers State University Teaching Hospital (RSUTH) over a two-year period.

Methodology: A retrospective review of hospital records of all pregnant women booked for antenatal care (ANC) at RSUTH in a two-year period, from 1ST May 2017 to 30TH April 2019, was carried out. Data on patients' age, parity and educational level, and gestational age at booking were retrieved using structured pro-forma and analyzed using Epi Info Version 7. Test for significance using Chi-square was set at significant level of $P < 0.05$.

Results: There were 3560 cases, with a mean age of 31.5 ± 4.7 years and a mean gestational age at booking of 22.1 ± 6.8 weeks. Majority of the women (53.9%) booked in the second trimester. Only about a quarter (26.5%) booked early, with 73.5% booking late. Majority of the women (62.2%) fall within the 30-39 age group, are Multiparous (65.3%) and had tertiary education (72.5%). There was no statistically significant relationship between their ages and gestational age at booking ($\chi^2 = 3.372$, **p-value=0.761**). However, parity ($\chi^2 = 50.015$, **p-value=0.000**) and educational qualification ($\chi^2 = 18.358$, **p-value=0.001**) were statistically significant.

Conclusion: The majority of ANC attendees booked late for antenatal care in this study, and only about a quarter booked early. There was a statistically significant difference in parity and educational status as determinants of gestational age at booking, with the nullipara and primary education group booking earlier.

Key Words: Antenatal booking, Pattern, Gestational age, Socio-demographic Determinants, Rivers State,

Introduction:

Maternal and child health (MCH) are central to achieving the Sustainable Development Goal 3, related to maternal and neonatal outcomes of pregnancy [1]. As part of the global strategy to reduce the risk of stillbirth and maternal complications, the World Health Organization (WHO) aims to improve the quality of antenatal care by increasing the number of recommended visits (from four to eight), and advocating for the first ANC visit (also known as the booking visit) before 12 weeks of gestation. Early provision of ANC has been found crucial for ensuring optimal care and good maternal and fetal outcome [2].

This form of care is supervised to enable women attain and maintain a state of good health throughout pregnancy and ensures good outcomes for mother and child [3][4][5]. It exposes pregnant women to health education and counselling regarding the physiological changes that occur and risk factors to adverse pregnancy outcome. It affords opportunity for dietary counselling, immunization against tetanus and malaria chemoprophylaxis, and preexisting medical conditions are also detected early, with plans of management determined early. It is therefore important for women to book early to fully benefit from these interventions [6][7].

The booking visit, as recommended by WHO, should be initiated at ≤ 12 weeks in the focused antenatal care and < 14 weeks in the traditional antenatal care [8]. While this recommendation has been adopted by the national guidelines in developed countries like the United Kingdom and the USA, no such national guidelines on antenatal care exist in most developing countries including Nigeria. However, based on the WHO recommendation, commencement of ANC within the first 14 weeks of gestation is widely accepted as early booking, and late if commenced after this period [9][10][11].

In spite of the documented benefits of early ANC booking and subsequent care, low antenatal coverage, few visits and late booking are still common problems in Sub-Saharan Africa (SSA), making it difficult to achieve the WHO recommended schedule [10][11][12]. The minimum ANC visits recommended by WHO was possible only for less than about a third of the pregnant women in some SSA countries like Niger (15%), Ethiopia (19%), Chad (23%), Burundi (33%) and Rwanda (35%) [13]. Previous studies have reported the prevalence of late booking in Nigeria to be between 70.9% to 86% [3][14][15], with regional variation in the mean gestational age at booking ranging from 24.3 weeks in Abakaliki [5], 21.8 weeks in Ibadan [9], 23.6 weeks in Niger-Delta [10] and 23.6 weeks in Sokoto [16].

Various reasons for this late booking, as deduced by previous studies, ranges from a belief that there are no advantages for booking in the first trimester, financial constraint and distance from health facility; to busy work schedule, apparently problem-free pregnancy and perceived enemies who may harm the pregnancy [3][5][10][17]. Only a handful of studies have examined the biological or demographic determinants of the gestational age at booking. Onoh RC et al [5] in their Abakaliki study found that Socio-Biological factors did not significantly contribute to the Gestational age at booking. Aduloju OP et al [18] found that the age, parity and occupation (educational status) of women were significantly associated with early booking, although on further testing with logistic regression only occupation remained significant. Nwagha UI et al [19] specifically looked at the influence of Parity on gestational age at booking and found that the average gestational age at booking was significantly lower in Nulliparous women compared to multiparous and grand-multiparous women.

This study therefore, seeks to determine the pattern of gestational age at antenatal booking and possible socio-demographic (Age, Parity, Education) determinants, so as to determine the group of women that will need more targeted health education programmes to change the trend of high late booking in SSA.

Methodology:

A retrospective, quantitative study of hospital records of all pregnant women booked at the RSUTH for antenatal care in a two years period (1ST May 2017 to 30TH April 2019) was carried out. The study period of two years was chosen as it marked the period of conversion of the former Braithwaite Memorial Specialist Hospital to the RSUTH and is the expected period of child spacing, to limit duplication of cases.

Data on patients' age, parity and educational status and gestational age (GA) at booking were retrieved using structured pro-forma. The GA at booking were categorized into First Trimester (1-13 weeks), Second Trimester (14-26 weeks) and Third Trimester (27-40 weeks). The Age was categorized into ≤ 19 years, 20-29 years, 30-39 years and ≥ 40 years. The Parity was categorized into Nullipara (para 0), Multipara (para 1-4) and Grand Multipara (para ≥ 5).

All pregnant women who registered for antenatal care during the study period, with their gestational age recorded in the records books as confirmed by sure date of last menses or early ultrasound scan, were included and a formal sample size was not calculated. There were 3560 cases that met the stated criteria and formed the study population.

Data were analyzed using United States CDC Epi Info Version 7. Data were summarized using frequencies and proportions for categorical variables; and means, standard deviation, medians and range employed for quantitative variables. The test of significance for the categorical and discrete variables was done using Chi-square at statistically significant level of $P < 0.05$.

As the study involved review of existing MCH records, ethical approval, and a waiver for informed consent, was obtained from the Ethics and Research committee of the RSUTH.

The RSUTH is one of two tertiary hospital for referral from all private clinics, maternity homes, primary health centers and secondary health facilities from all the 23 Local government areas of Rivers State, Nigeria. The hospital is funded by the Government and patients are expected to pay directly for services (except few that participate in the National Health Insurance Scheme). It provides emergency obstetric services to women referred from other centers, as well as providing antenatal care and delivery services for low and high-risk pregnant women booked with the hospital. The hospital is well equipped and has round the clock availability of qualified team comprising of Obstetricians, Pediatricians and Anaesthetist. There is availability of laboratory and blood bank services in the hospital. Referrals are made directly from primary, as well as secondary, health facilities not following any pattern. The Department of Obstetrics and Gynaecology runs antenatal clinics Mondays through Fridays and about 2000 women book annually for ANC, with over 1500 deliveries per annum.

Results:

A total of 3560 women registered for antenatal care during the study period with a mean age of 31.5 ± 4.7 years and a mean gestational age at booking of 22.1 ± 6.8 weeks. The age range was 15-48 years and the modal parity was 3. Majority of the women (53.9%) booked in the second trimester and only about a quarter (26.5%) booked early in the first trimester, with 73.5% booking late. Majority of the women (62.2%) fall within the 30-39 age group, are Multiparous (65.3%) and had tertiary education (72.5%). See table 1 and 2.

Table 1: Pattern of Gestational age at booking over two years (n = 3560):

Gestational age	Frequency (N)	Percentage (%)
1 st Trimester	943	26.5

2 nd Trimester	1918	53.9
3 rd Trimester	699	19.6
Total	3560	100.0

Table 2: Socio-demographic pattern of the Antenatal Clinic Attendees (n = 3560).

Characteristics	Frequency	Percentage
Age group		
≤19 years	42	1.2
20-29 years	1137	31.9
30-39 years	2216	62.2
≥40 years	165	4.6
Total	3560	100.0
Parity		
Nullipara	1125	32.4
Multipara	2326	65.3
Grand-multipara	79	2.2
Total	3560	100.0
Educational qualification		
Primary	58	1.6
Secondary	922	25.9
Tertiary	2580	72.5
Total	3560	100.0

Figure 1, 2 and 3 show the distribution of the trimester at booking against the age, parity and educational status categories respectively. Majority of the women across the various age categories booked in the second trimester, and so also it was for the various categories of parity and educational status.




Fig 1: Age distribution of gestational age at booking of the clinic Attendees.




Fig 2: Parity distribution of gestational age at booking of the clinic Attendees.




Fig 3: Educational status distribution of gestational age at booking of the clinic Attendees.

Table 3 shows the statistical analysis of the Socio-demographic characteristics and gestational age at booking of antenatal clinic attendees at the RSUTH. There was no statistically significant relationship between their ages and gestational age at booking ($\chi^2 = 3.372$, **p-value = 0.761**). However, parity ($\chi^2 = 50.015$, **p-value = 0.000**) and educational qualification ($\chi^2 = 18.358$, **p-value = 0.001**) were statistically significant. Despite majority of the women across the various parity categories booking in the second trimester, a significant number of the nullipara booked more in the first trimester. So also, a significant number of those with primary education booked early than those with higher education.

Table 3: Socio-Demographic determinants of gestational age at booking of the Attendees (n = 3560):

Characteristics	Gestational age			Total	Chi-square	df	p-value
	1 st Trimester (n/%)	2 nd Trimester (n/%)	3 rd Trimester (n/%)				
Age group					3.372	6	0.761
≤19 years	12 (28.6)	19 (45.2)	11 (26.2)	42			
20-29 years	301 (26.5)	601 (52.9)	235 (20.7)	1137			
30-39 years	586 (26.4)	1211 (54.6)	419 (18.9)	2216			
≥40 years	44 (26.7)	87 (52.7)	34 (20.6)	165			
Parity					50.015	4	0.000*
Nullipara	390 (33.8)	555 (48.1)	210 (18.2)	1155			
Multipara ^R	531 (22.8)	1326 (57.0)	469 (20.2)	2326			
Grand-multipara	22 (27.8)	37 (46.8)	20 (25.3)	79			
Educational qualification					18.358	4	0.001*

Primary	18 (31.0)	26 (44.8)	14 (24.1)	58
Secondary ^R	201 (21.8)	513 (55.6)	208 (22.6)	922
Tertiary	724 (28.1)	1379 (53.4)	477 (18.5)	2580

***p-value is significant**

Discussion:

This study showed that about a quarter of the study population (26.5%) booked within the WHO recommended time while the rest booked late. The late booking of 73.5% found is however lower than the 77.3% in Ekiti [18], 82.0% in Lagos [15] and 85.9% in Ibadan [9] but higher than the 71.6% in Makurdi [17]. The mean gestational age at booking for ANC in this study was 22.1 weeks which was comparable to 21.1 weeks in Ekiti [18] and 21.8 weeks in Ibadan [9]; slightly higher than 19.1 weeks in Makurdi and 20.3 weeks in Lagos [15]; but lower than 23.5 weeks in Sokoto [16], 23.7 weeks in Benin [10], and 24.3 weeks in Abakaliki [5]. It is interesting to note that despite the documented benefits of early ANC booking and what is obtained in developed and some developing countries, late booking is still predominant in our environment, as shown by this study.

Our study showed that primigravidae (Nulliparous) women were more likely to book early for ANC than the multipara and this was similar to the findings of previous studies by Okunlola et al [9], Gharoro and Igbafe [12], Aduloju et al [18], and Nwagha et al [19]. The higher the parity, the more likely pregnant women may feel experienced enough as not to book early, which is actually contrary to the fact that they are more likely to be at risk of complications associated with high parity and advanced maternal age.

The educational level of women has been shown by some previous studies to significantly affect the timing of ANC booking. Higher education of women was found to affect early booking in the studies by Adekanle and Isawunmi [15], Ekele and Audu [16] and Aduloju et al [18]. The argument has been that higher educational level and occupation of the women encouraged early booking because they are well informed of the benefits of ANC and are financially empowered, since majority of them have good source of income. Contrary to the above reasoning, our study found a statistically significant association of lower educational status and early booking. This may be explainable from the fact that It is common practice in our environment for well-educated and financially empowered women to have multiple booking as they can afford it, booking early in private clinics which provide better comfort and privacy than Government owned hospital for antenatal care; and booking later in the Government hospitals close to their delivery for specialist obstetrician care. This multiple booking also stem from the fact that services in the government hospitals are frequently interrupted by frequent industrial strike actions. The poorly educated and supposedly less empowered cannot afford multiple booking and hence come early and straight to the government hospital for booking.

Contrary to the findings in our study, and others sited above, of parity and education significantly affecting early booking for ANC, it is at variance with the findings of Onoh et al [5] who reported that all the sociodemographic characteristics did not have any significant influence on the timing of antenatal care initiation.

This study may be limited by the fact that it is a retrospective study and there might be inaccuracies in the documented data such as gestational age. The study is also hospital based and may not reflect what happens in the general population in the state, as some pregnant women do not access the tertiary institution.

Conclusion:

The majority of ANC attendees booked late for antenatal care in this study, and only about a quarter booked early. There was a statistically significant difference in parity and educational status as

determinants of gestational age at booking, with the nullipara and primary education group booking earlier. A community-based study is advocated as this will help determine the actual determinants of antenatal booking peculiar to the pregnant women within and around the state.

Women should be adequately educated and informed about the concept of early antenatal registration and its benefits, since no pregnancy is the same as the other and unexpected adverse outcome may occur despite previous uncomplicated pregnancies. This should be done not only at the antenatal clinics but following discharge after delivery particularly for those who booked late or unbooked in that pregnancy.

References:

1. United Nations. Transforming our world: The 2030 agenda for sustainable development. 2015.
2. Moller A-B, Petzold M, Chou D, Say L. Early antenatal visit: a systematic analysis of regional and global levels and trends of coverage from 1990 to 2013. *Lancet Glob Heal*. 2017; 5: e977-83.
3. Oladokun A, Oladokun RE, Morhason-Bello I, Bello AF, Adedokun B. Proximate predictors of early antenatal registration among Nigerian pregnant women. *Ann Afr Med* 2010; 9: 222-5.
4. Adeyemi AB, Makinde ON, Ajenifuja KO, Soyinka AS, Ayinde AK, Ola BA, et al. Determinants of antenatal booking time in South-Western Nigeria setting. *West Afr J Med* 2007; 26: 293-7.
5. Onoh R., Umerora O, Agwu U, Ezegwui H, Ezeonu P, Onyebuchi A. Pattern and determinants of antenatal booking at Abakaliki South-East Nigeria. *Ann Med Health Sci Res* 2012; 2: 169-75.
6. Mpembeni RN, Killewo JZ, Leshabari MT, Massawe SN, Jahn A, Mushi D, et al. Use pattern of maternal health services and determinants of skilled care during delivery in Southern Tanzania: implications for achievement of MDG-5 targets. *BMC Pregnancy Childbirth* 2007; 7: 29.
7. Rockers PC, Wilson ML, Mbaruku G, Kruk ME. Source of antenatal care influences facility delivery in rural Tanzania: A population-based study. *Matern Child Health J* 2009; 13: 879-85.
8. UNDP/UNFPA/WHO/World Bank Special programme of Research development and Research training in Human Reproduction. WHO Antenatal Care Randomized trial: Manual for implementation of the new model. Geneva: Department of Reproductive Health and Research, Family and Community Health, World Health Organization; 2002.
9. Okunlola MA, Ayinde OA, Owonikoko KM, Omigbodun AO. Factors influencing gestational age of antenatal booking at the University College Hospital, Ibadan, Nigeria. *J Obstet Gynaecol* 2006; 26: 195-7.
10. Ebeigbe PN, Igberase GO. Reasons given by pregnant women for late initiation of antenatal care in Niger-Delta Nigeria. *Ghana Med J* 2010; 44: 47-51.
11. Zageye AM, Bitew BD, Koye DN. Prevalence and determinants of early antenatal care visit among pregnant women attending antenatal care in Debre Berhan Health Institutions, Central Ethiopia. *Afr J Reprod Health* 2013; 17: 130-6.
12. Gharoro EP, Igbafe AA. Antenatal care: some characteristics of the booking visit in a major teaching hospital in the developing world. *Med Sci Monit* 2000; 6: 519-22.
13. Berhan Y, Berhan A. Antenatal care as a means of increasing birth in the health facility and reducing maternal mortality: a systematic review. *Ethiop J Health Sci* 2014; 24: 93-104.
14. Okunlola MA, Owonikoko KM, Fawole AO, Adekunle AO. Gestational age at antenatal booking and delivery outcome. *Afr J Med Med Sci* 2008; 37: 165-9.
15. Adekanle OA, Isawunmi AI. Late antenatal care booking and its predictors among pregnant women in South-Western Nigeria. *J Allied Health Sci* 2008; 7: 4-7.

16. Ekele BA, Audu LR. Gestational age at antenatal clinic booking in Sokoto, Northern Nigeria. *Afr J Med Med Sci* 1998; 27: 161-3.
17. Ifenne DI, Utoo BT. Gestational age at booking for antenatal care in a tertiary health facility in North-Central Nigeria. *Niger Med J* 2012; 53: 236-9.
18. Aduloju OP, Akintayo AA, Ade-Ojo IP, Awoleke JO, Aduloju T, Ogundare OR. Gestational age at initiation of antenatal care in a tertiary hospital, Southwestern Nigeria. *Niger J Clin Pract* 2016; 19: 772-7.
19. Nwagha UI, Ugwu OV, Nwagha TU, Anyaehie USB. The influence of parity on the gestational age at booking among pregnant women in Enugu, South-East Nigeria. *Nigerian Journal of Physiological Sciences* 2008; 23(1-2): 67-70. Available at <http://www.bioline.org.br/np>.
<http://www.ajol.info/journal/njps>.

UNDER PEER REVIEW