

## **Original Research Article**

### **Effect of Aqueous Extract of *Khaya grandifoliola* C. DC. Stem Bark on some Disaccharidases Activity in Iron-Deficient Weanling Rats**

#### **ABSTRACT**

The effect of the aqueous extracts of *Khaya grandifoliola* stem bark on some disaccharidases (lactase and sucrase) in diet-induced anemia in weanling rats was investigated in this study. Weanling rats of 21 days old were maintained on iron-deficient diets for four weeks to induce anemia before treatment. A total of 35 weanling rats were used, grouped into five rats per group of iron-deficient diet/distilled water, iron-sufficient diet/distilled water, change of iron-deficient diet after four weeks to iron-sufficient diet, iron-deficient diet/ Standard drug, iron-deficient diet/25mg/kg body weight of aqueous plant extract, iron-deficient diet/50mg/kg body weight of aqueous plant extract, and iron-deficient diet/100mg/kg body weight of aqueous plant extract. Phytochemical screening of the extract revealed the presence of alkaloids, saponins, cardiac glycosides, tannins, anthraquinones and flavonoids. The extract administered orally produced significant increase in the activity of intestinal mucosa sucrase and lactase ( $P < 0.05$ ). The change of diet from iron deficient diet to iron sufficient diet increased disaccharidases activities in the intestinal mucosa. However, the aqueous extract of *Khaya grandifoliola* showed a higher disaccharidases activity when compared to the group of rats that were fed iron-sufficient diet. This study revealed that plant extract administered increased disaccharidases activity in the intestinal mucosa in diet-induced anemic group of weanling rats and thus lends credence to *Khaya grandifoliola* use in folklore medicine in the management of anemia.

Keywords: *Khaya grandifoliola*, iron-deficient, Lactase, Sucrase

## INTRODUCTION

Iron deficiency anemia (IDA) is the most common type of anemia that results from an inadequate iron supply to aid the production of healthy red blood cells (Adamson, 2005). Iron deficiency is usually a result of the depletion of the body's iron stores; hence, a restricted supply of iron to various tissues becomes apparent. This may result in a reduction in the synthesis of red blood cells and iron-dependent intra-cellular enzymes participating in many metabolic pathways (Earley et al., 2000). Iron deficiency has been reported to reduce the activity of some disaccharidases (Fernandes et al., 1997; Sriratanaban and Thayer 1971).

Although there are various drugs used for the treatment of anemia (Macdougall, 2005; Smith et al., 2008; Locatelli et al., 2001), they are not affordable to many poor people in the developing countries such as Nigeria. Furthermore, the rural populations in various parts of the world do not have appreciable access to high-quality drugs for the treatment of anemia, hence, the reliance on herbal products for the treatment of diseases and anemia. Some of the plants with ethnobotanical and scientific claims being used for the treatment of anemia include *Waltheria indica*, *Sorghum bicolor*, *Khaya grandifoliola*, and *Mangifera indica* (Peter et al., 2014; Oladiji et al., 2007; Nvvinuka et al., 2008; Olowokudejo et al., 2008; Modupe and Oladiji, 2016).

*Khaya grandifoliola* is a medicinal plant commonly found in Nigeria and many parts of the plant are valuable to traditional medicine. Some sicknesses like gastric pain, lumbago, stomach ache, worm infestation, cough, fever, and rheumatism are treated with concoctions made from the bark of *Khaya grandifoliola* (Stephen et al., 2009). The stem bark of this plant has also been previously reported to have anti-malarial, anti-ulcer, anti-hypoglycaemic, anti-hypoproteinaemia, and anti-hypocholesterolaemia activity (Makinde et al., 1988, Bumah et al., 2005, Njikam and Njikam, 2006). Several research studies have lent scientific credence to the use of various parts of *Khaya grandifoliola* in the treatment of several diseases, however, there is still no adequate information on the effect of the plant extract on lactase and sucrase activity as an indicator for anti-anemic potentials of *Khaya grandifoliola*. This study investigated the effect of aqueous

extract of *Khaya grandifoliola* stem bark on lactase and sucrase activity in the intestinal mucosa of iron-deficient weanling rats.

## **Materials and Methods**

### **Experimental Animals**

Forty-five weanling albino rats of both sexes (*Rattus norvegicus*) with a mean weight of 40.0 g  $\pm$  3.0 g were obtained from the Animal Holding Unit of the Department of Biochemistry, University of Ilorin, Nigeria. This study was carried out as approved by the Department of Biochemistry Ethical Committee on the use of laboratory animals at the University of Ilorin.

### **Feed Components**

Yellow maize (*Zea mays*) and locust bean (*Parkia biglobosa* A. Jacq) seeds were obtained locally from Baboko Market, Ilorin, Nigeria, while the soybean oil used was a product of Grand Cereals and Oil Mills Limited, Bukuru, Jos, Nigeria. The vitamin mix was a product of BASF Aktiengesellschaft, Germany Pantex, Netherland. Component chemicals of the mineral mix used were products of Sigma-Aldrich Chemical Limited, London, UK.

### **Reagents**

The reagents used were of analytical grade and were prepared in all glass-distilled water. The reagents were stored in reagent bottles (Plummer, 1987).

### **Plant Identification and Preparation of Extract**

The stem barks of the plant were obtained from Ikaro in the Ose Local Government Area of Ondo State in the South West region of Nigeria. It was authenticated in the Department of Plant Biology, University of Ilorin, Ilorin, Nigeria, where a voucher specimen was deposited in the herbarium.

The method described by Oladiji et al. (2007) was used in the preparation of the plant extract. The stem barks of the plant were air-dried until a constant weight was obtained. The dried pieces were then pulverized using an electric blender, and the powder obtained was stocked in a plastic container. A known weight (15 g) of the powder was poured into 100 ml of distilled water and

immediately boiled for 25 min. The resulting solution was left to cool and filtered with Whatman filter paper. The filtrate was concentrated on a water bath (kept at 45 °C) until a constant weight was obtained to give 2.32 g ± 0.03 of the residue (brownish-black slurry), which corresponded to percentage yield of 15.47%. The residues were pooled together and reconstituted in distilled water to give the doses of 25, 50, and 100mg/kg body weight.

The reconstituted aqueous extract was administered orally using an oropharyngeal cannula to all the animals in different groups.

### Diet Formulation

The composition of iron-deficient and iron-sufficient diets per kg diet is described in Table 1. The components of the diets were thoroughly mixed and made into pellets to ensure homogeneity and proper handling by the animals. It was produced weekly and packed into air-tight polythene bags to prevent rancidity, auto-oxidation of the oil, and microbial contamination. The proximate analysis of the compounded feeds was also carried out.

Table 1: Feed components of iron sufficient and iron-deficient diets

Feed components	Iron-sufficient (g/kg)	Iron-deficient (g/kg)
Locust bean seed	500	500
Maize flour	315	315
Soybean oil	40	40
Sucrose	100	100
Methionine	5	5
Mineral mix <sup>b</sup>	30	30
Vitamin mix <sup>a</sup>	10	10
FeSO <sub>4</sub> ·7H <sub>2</sub> O	157.36mg/Kg	38.216 mg/Kg

Soybean oil: polyunsaturated fatty acids (58%), monounsaturated fatty acids (29%), saturated fatty acids (13%).

<sup>a</sup>Vitamin mix (per kg of diet): vitamin A, 100,000 IU; vitamin D3, 10,000 IU; vitamin E, 100 mg; vitamin B1, 20 mg; vitamin B2, 40 mg; Lysine, 10g; d-calcium pantothenate, 100 mg; vitamin B6, 15 mg; vitamin C, 250 mg; vitamin K3, 15 mg; folic acid, 5000 mcg; nicotinic acid, 200 mg; biotin, 150 mcg; inositol, 80 mg.

<sup>b</sup>Mineral mix (g/kg): CoCl<sub>2</sub>·6H<sub>2</sub>O (0.001), CuSO<sub>4</sub>·5H<sub>2</sub>O (0.079), MnSO<sub>4</sub>·7H<sub>2</sub>O (0.178), KI (0.032), NaCl (3.573), ZnCO<sub>3</sub> (1.60), CaSO<sub>4</sub> (11.610), MgSO<sub>4</sub>·7H<sub>2</sub>O (2.292), K<sub>2</sub>HPO<sub>4</sub> (10.559). Control diet contained 1.078 g FeSO<sub>4</sub>·7H<sub>2</sub>O.

### **Animal Grouping and Extract Administration**

The animals were kept in well-ventilated house conditions (temperature:  $22\pm 3^{\circ}\text{C}$ ; photoperiod: 12h/12h light/dark cycle; relative humidity: 45-50 %). They were allowed free access to normal rat chow and distilled. The acclimatization was done for seven days, and they were then fasted for 24 hours (without food but water) before the commencement of the experiment. The animal grouping consisted of an initial two groups:

A – 10 weanling rats maintained on iron-sufficient diet designated as **ISF**.

B – 35 weanling rats maintained on an iron-deficient diet designated as **IDF**.

Animals in groups A and B were maintained on their respective diets for four weeks. At the end of the four weeks feeding period, five rats each from ISF and IDF groups were sacrificed, and th were determined. The remaining weanling rats in groups B were further grouped into six with five rats in each group as follows:

B1- Iron deficient weanling rats fed with iron-deficient feed for two weeks (iron deficient feed all through) designated as **IDF**

B2- Iron deficient weanling rats fed with iron sufficient feed for two weeks (change of feed) designated as **COF**

B3- Iron deficient weanling rats orally administered daily for two weeks with iron supplement syrup (a standard iron supplement) designated as **SD**

B4- Iron deficient weanling rats orally administered daily for two weeks with 25 mg/kg body weight of aqueous extract of *Khaya grandifoliola* stem bark designated as **KG 25mg**.

B5- Iron deficient weanling rats orally administered daily for two weeks with 50 mg/kg body weight of aqueous extract of *Khaya grandifoliola* stem bark designated as **KG 50mg**.

B6- Iron deficient rats orally administered daily for two weeks with 100 mg/kg body weight of aqueous extract of *Khaya grandifoliola* stem bark designated as **KG 100mg**.

The rest of the animals in group A were still fed with iron sufficient feed for two weeks (iron sufficient all through) designated as **ISF**.

The extracts and the distilled water were administered to the various groups using the oropharyngeal cannula.

### **Determination of lactase and sucrase activity**

Method described by Dahlqvist (1968) was used and the result of the intestinal mucosa disaccharidases was reported as units (U)

### **Statistical Analysis**

Results were expressed as the mean  $\pm$  SEM of five determinations. The data were analyzed using Duncan Multiple Range Test and complemented with Student's t-test. The differences were considered statistically significant at  $p < 0.05$ . All the analyses were done using SPSS version 16.0 software (SPSS Inc., Chicago, IL, USA).

## Results

Phytochemical screening of the aqueous stem barks of *K. grandifoliola* revealed the presence of alkaloids, flavonoids, tannins and saponins, cardiac glycosides, and anthraquinones in its aqueous extracts.

The proximate composition of both iron-deficient (ID) and iron sufficient (IS) formulated feeds are shown in Table 2. Proximate analysis of the diets showed that the components in the iron-deficient formulated feed (ID) were substantially similar to those in the iron-sufficient formulated feed.

Table 2: Proximate composition of iron-deficient and iron-sufficient diets

Components	ID (%)	IS (%)
Crude protein	20.78	21.44
Carbohydrate	60.00	59.65
Lipid	5.03	5.15
Crude fiber	2.97	3.02
Total ash	4.20	4.35
Moisture	7.02	6.39

ID, iron-deficient; IS, iron-sufficient; crude protein was obtained using the expression  $\text{nitrogen} \times 6.25$

### **Effect of *Khaya grandifoliola* on intestinal mucosa sucrase activity in iron-deficient weanling rats**

From Table 3, there was significant decrease in the intestinal mucosa sucrase activity in IDF group when compared to that of ISF group. The activity of sucrase in the SD group and ISF group showed a significant increase when compared IDF group. Also, there was also a significant increase in sucrase activity in COF and SD groups when compared to IDF group. The

result also showed a significant increase in the activity of sucrase in COF group when compared with both SD and ISF groups. Administration of the aqueous extracts of *K. grandifoliola* stem bark led to significant ( $p>0.05$ ) increase in the activity of the intestinal mucosa Sucrase (Table 3) at all dose levels investigated compared to the other groups. However, the dosage at 50mg/kg body weight of aqueous extract of *K. grandifoliola* stems bark has the highest significant increase ( $p<0.05$ ) in the activity of intestinal mucosa Sucrase.

**Table 3: Effect of *Khaya grandifoliola* on intestinal mucosa sucrase in iron deficient weanling Rats**

Group	Sucrase Specific Activity(U/g )
ISF	9.00±1.7 <sup>a,b</sup>
IDF	5.27±0.2 <sup>a</sup>
COF	9.13±1.4 <sup>a,b</sup>
SD	6.93±0.3 <sup>a,b</sup>
25mg/kg b. wt.	8.93±0.2 <sup>a,b</sup>
50mg/kg b. wt.	16.20±1.6 <sup>c</sup>
100mg/kg b. wt.	9.67±1.7 <sup>b</sup>

Values are expressed as mean of five replicates ± S.E.M. and those with different superscripts along a column are statistically different ( $p<0.05$ ). The extract was administered for 14days; IDF: Iron-deficient diet fed rats; ISF: Iron-sufficient diet fed rats; COF:Change of feed to iron-sufficient diet; SD: Group treated with Standard Drug; ; Kg b. wt.: Kilogramme Body weight

#### **Effect of *Khaya grandifoliola* on intestinal mucosa lactase in iron deficient weanling rats**

From Table 4, lactase activity showed a significant decrease in IDF group when compared with ISF group. SD group has significant increase in the intestinal mucosa lactase activity when compared with both ISF and COF groups. In this study, the SD group showed a significant



increase in lactase activity when compared with ISF group. This result also showed a significant increase in lactase activity in COF group when compared IDF group. Administration of the aqueous extract of *K. grandifoliola* stem bark led to significant increase in the activity of intestinal mucosa lactase (Table 4) at all doses investigated compared to other groups ( $p < 0.05$ ). However, the dosage at 50mg/kg body weight of aqueous extract of *K. grandifoliola* stems bark has the highest significant increase in the activity of intestinal mucosa lactase ( $p < 0.05$ ).

**Table 4: Effect of *Khaya grandifoliola* on intestinal mucosa lactase in Iron-deficient weanling rats**

Group	Lactase Specific Activity(U/g)
ISF	4.13±1.0 <sup>a,b</sup>
IDF	1.73±0.9 <sup>a</sup>
COF	3.27±2.0 <sup>a,b</sup>
SD	5.73±1.8 <sup>a,b</sup>
25mg/kg b. wt.	4.93±0.7 <sup>a,b</sup>
50mg/kg b. wt.	7.13±1.3 <sup>a,b</sup>
100mg/kg b. wt.	5.87±0.8 <sup>a,b</sup>

Values are expressed as mean of four replicates ± S.E.M. and those with different superscripts along a column are statistically different ( $p < 0.05$ ). The extract was administered for 14days; IDF: Iron-deficient diet fed rats ;ISF: Iron-sufficient diet fed rats ; COF:Change of feed to iron-sufficient diet; SD: Group treated with Standard Drug; ; Kg b.wt.: Kilogramme Body weight

## Discussion

Secondary metabolites screening carried out on the aqueous stem barks of *K. grandifoliola* indicated that, *K. grandifoliola* stem barks contain alkaloids, flavonoids, tannins and saponins, and anthraquinones in its aqueous extracts (Steven et al., 2009). These phytochemicals are known to perform several general and specific roles in plants, and may exhibit different biochemical and pharmacological actions in different species of animals when consumed.

In this study, the effect of aqueous extract of *Khaya grandifoliola* on some disaccharidases activity in iron-deficient weanling rats was investigated. It has been reported previously that disaccharidases have reduced activity in iron-deficient anemia (Fernandes et al., 1997; Sriratanaban and Thayer, 1971).

From tables (3&4), the results showed that, there was significant decrease in the activities of the disaccharidases in weanling rats fed iron- deficient diet in the study. Similar reports have shown that dietary iron deficiency can cause a significant decrease in disaccharidase activity (Hoffbrand and Broitman, 1969., Bolin et al., 1971., Sriratanaban and Thayer, 1971., Lanzkowsky et al., 1982., Buts et al., 1984., Buts et al., 1986., Fernandes et al., 1997., Vieira et al., 2000).

Iron-deficient diet fed rats treated with the reference drug and the group of iron deficient weanling rats administered aqueous extract of *K. grandifoliola* stem bark showed significant increase in the activity of disaccharidases in the intestinal mucosa. This result is in agreement with claim that, decrease in disaccharidases activity is reversible by iron supplementation (Fernandes et al., 1997., Lanzkowsky et al., 1982., Vieira et al., 2000).

The change of diet from iron deficient diet to iron sufficient was also able to increase the activities of intestinal mucosa sucrase and lactase, however, the plant aqueous extract of *K. grandifoliola* showed higher increase in the activity of the disaccharidases.

The significant increase in the activities of the disaccharidases in the iron-deficient weanling rats from administered aqueous extract of *K. grandifoliola* stem bark in this study may be as a result of the chemical constituent(s) of the of the aqueous extract of *K. grandifoliola* stem bark. It is possible that the extracts mechanism(s) of action for the significant increase in the activities of the disaccharidases may be due to an increase in the gene expression of both sucrase and lactase. Since the extract contains saponins and they are membrane active agents. It is possible that saponins stimulate the release of glucocorticoids that further initiate various biochemical reactions, hence, resulting in the induction of the disaccharidases in the intestinal cells.

The administered doses of aqueous extract of *Khaya grandifoliola* in this study were within the tolerable limit and produced no toxic effect. Tolerable limit of aqueous extract of *Khaya grandifoliola*, as investigated and reported by Njikam and Njikam (2006), showed that the LD<sub>50</sub> of the plant extract was 5.5g/kg bodyweight. The LD<sub>50</sub> is over a thousand times more than the 50mg/kg bodyweight shown to be the most efficacious of the three doses administered.

## **Conclusion**

In this study, the administration of the aqueous extract of *Khaya grandifoliola* stem bark increased the activity of both sucrase and lactase in the intestinal mucosa of iron-deficient weanling rats and 50mg/kg.bw dose showed to be the most efficacious of all the dose levels. This investigation further strengthens the claim of the usage of the aqueous extracts of *Khaya grandifoliola* stem bark in the management of iron-deficient anemia.

## **COMPETING INTERESTS DISCLAIMER:**

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly used products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of

knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

## References

- Adamson, J. W. 2005. Iron deficiency and other hypoproliferative anemias. *HARRISONS PRINCIPLES OF INTERNAL MEDICINE*, 16, 586.
- Bolin, T. D., McKern, A., Davis, A. E., 1971. The effects of iron deficiency, protein deficiency and worm infestation upon disaccharidase activity in the rat. *Aust NZ J Med* 1: 218–223.
- Bumah, V. V., Essien, E. U., Agbedahunsi, J. M. & Ekah, O. U. 2005. Effects of *Khaya grandifoliola* (Meliaceae) on some biochemical parameters in rats. *Journal of ethnopharmacology*, 102, 446-449.
- Buts, J. P., Delacroix, D. L., Dekeyser, N., Paquet, S., Horsmans, Y., Boelens, M., Van Craynest, M. P., De Meyer, R., 1984. Role of dietary iron in maturation of rat small intestine at weaning. *Am J Physiol Gastrointest Liver Physiol* 246: 725–731.
- Buts, J. P., Vamecq, J., van Hoof, F., 1986. Alteration of intracellular synthesis of surface membrane glycoproteins in small intestine of iron-deficient rats. *Amr J Physiol Gastrointest Liver Physiol*; 251: 737–743.
- Dahlqvist, A., (1968). Assay of intestinal disaccharidases. *Analytical Biochemistry*; 22: 99–107.
- Earley, C. J., Connor, J. R., Beard, J. L., Malecki, E. A., Epstein, D. K. & Allen, R. P. 2000. Abnormalities in CSF concentrations of ferritin and transferrin in restless legs syndrome. *Neurology*, 54, 1698-1700.
- Fermandes, M. I. M., Galvao, L. C., Bortolozzi, M. F., Oliveira, W. P., Zucoloto, S. & Bianchi, M. L. P. 1997. Disaccharidase levels in normal epithelium of the small intestine of rats with iron-deficient anaemia. *Brazilian Journal of Medical Biological Research*, 30, 849-854.
- Hoffbrand, A. V. and Broitman, S. A., 1969. Effect of chronic nutritional iron deficiency on the small intestinal disaccharidase activities of growing dogs. *Proc Soc Exp Biol Med* 130: 595–598.
- Lanzkowsky, P., Karayalcin, G., Miller, F., 1982. Disaccharidase levels in iron deficient rats at birth and during the nursing and postweaning periods: response to iron treatment. *Pediatr Res* 16: 318–323.

- Locatelli, F., Olivares, J., Walker, R., Wilkie, M., Jenkins, B., Dewey, C. & Gray, S. J. 2001. Novel erythropoiesis stimulating protein for treatment of anemia in chronic renal insufficiency. *Kidney international*, 60, 741-747.
- Macdougall, I. C. 2005. CERA (Continuous Erythropoietin Receptor Activator): a new erythropoiesis-stimulating agent for the treatment of anemia. *Current hematology reports*, 4, 436-440.
- Makinde, J. M., Awe, S. O. & Agbedahunsi, J. M. 1988. Effect of Khaya grandifoliola extract on Plasmodium berghei berghei in mice. *Phytotherapy Research*, 2, 30-32.
- Modupe, O. and Oladiji, T.A., 2016. Optimizing dose of aqueous extract of Mangifera indica L stem bark for treating anaemia and its effect on some disaccharidases activity in iron deficient weanling rats. *Journal of Nutrition & Intermediary Metabolism*, 3, pp.18-22.
- Njikam, R. N. & Njikam, N. 2006. Curative Dose of Khaya grandifoliola. Stem Bark for the Treatment of Gastric Ulcers Using Wistar Rats. *Pharmaceutical biology*, 44, 152-155.
- Nvvinuka, N. M., Monanu, M. O. & Nwiloh, B. I. 2008. Effects of aqueous extract of Mangifera indica L.(Mango) stem bark on haematological parameters of normal albino rats. *Pakistan Journal of Nutrition*, 7, 663-666.
- Oladiji, A. T., Jacob, T. O. & Yakubu, M. T. 2007. Anti-anaemic potentials of aqueous extract of Sorghum bicolor (L.) moench stem bark in rats. *Journal of Ethnopharmacology*, 111, 651-656.
- Olowokudejo, J. D., Kadiri, A. B. & Travih, V. A. 2008. An ethnobotanical survey of herbal markets and medicinal plants in Lagos State of Nigeria. *Ethnobotanical leaflets*, 2008, 116.
- Peter, E. L., Rumisha, S. F., Mashoto, K. O. & Malebo, H. M. 2014. Ethno-medicinal knowledge and plants traditionally used to treat Anemia in Tanzania: A cross sectional survey. *Journal of ethnopharmacology*.
- Plummer, D. T. 1987. *An introduction to practical biochemistry*, McGraw-Hill Book Company London.
- Smith, R. E., Aapro, M. S., Ludwig, H., Pintér, T., Šmakal, M., Ciuleanu, T. E., Chen, L., Lillie, T. & Glaspy, J. A. 2008. Darbepoetin alfa for the treatment of anemia in patients with active cancer not receiving chemotherapy or radiotherapy: results of a phase III, multicenter, randomized, double-blind, placebo-controlled study. *Journal of Clinical Oncology*, 26, 1040-1050.
- Sriratanaban, A. and Thayer, W. R. Jr., 1971. Small intestinal disaccharidase activities in experimental iron and protein deficiency. *Am J Clin Nutr* 24: 411–415.
- Stephen, U. A., Abiodun, F., Osahon, O. & Ewaen, E. 2009. Phytochemical analysis and antibacterial activity of Khaya grandifoliola stem bark. *Journal of Biological Sciences*, 9, 63-67.
- Vieira, M. R., Galvao, L. C., Fernandes, M. I., 2000. Relation of the disaccharidases in the small intestine of the rat to the degree of experimentally induced iron-deficiency anemia. *Braz J Med Biol Res*, 33: 539–544.

Yakubu, M. T., Akanji, M. A. & Oladiji, A. T. 2005. Aphrodisiac potentials of the aqueous extract of *Fadogia agrestis* (Schweinf. Ex Hiern) stem in male albino rats. *Asian Journal of Andrology*, 7, 399-404.

UNDER PEER REVIEW