
A sequence of models of generalized second-order Dedekind theory of real numbers with increasing powers

*Original
Research Article*

Abstract

The paper is devoted to construction of some closed inductive sequence of models of the generalized second-order Dedekind theory of real numbers with exponentially increasing powers. These models are not isomorphic whereas all models of the standard second-order Dedekind theory are. The main idea in passing to generalized models is to consider instead of superstructures with the single common set-theoretical equality and the single common set-theoretical belonging superstructures with several generalized equalities and several generalized belongings for first and second orders. The basic tools for the presented construction are the infraproduct of collection of mathematical systems different from the factorized Loś ultraproduct and the corresponding generalized infrafiltration theorem. As its auxiliary corollary we obtain the generalized compactness theorem for the generalized second-order language.

Keywords: second-order language; generalized models; infraproduct; ultraproduct; non-standard analysis

2010 Mathematics Subject Classification: 03C85; 03C20; 26E35; 03C98; 11U07; 11U09; 03H05

1 Introduction

It is well known that all *standard* models of the standard second-order Dedekind theory of real numbers are isomorphic (see, for example, [1, 7.2]). The paper is devoted to the exposition of some *generalized* second-order Dedekind theory of real numbers with non-isomorphic generalized models.

More precisely, the paper is devoted to construction of some closed inductive sequence R_i ($1 \leq i \leq \omega_0$) of models of the generalized second-order Dedekind theory of real numbers with

exponentially increasing powers. The models R_i ($0 \leq i < \omega_0$) are embedded in each other as submodels and at the same time they all are embedded in the *limit-closer model* R_{ω_0} as extending submodels. These generalized models are completely different from mathematical systems presented in [2, 2.14] under the name of *non-standard analysis*.

The main idea in passing to generalized models is to consider the *generalized second-order language* $L(\Sigma_2^g)$ of some *generalized signature* Σ_2^g containing, in addition to individual and predicative constants and variables, some symbols δ_τ of *generalized equalities* and some symbols ε_τ of *generalized belongings* for first-order types τ and second-order types $\tau \equiv [\tau_0, \dots, \tau_k]$.

Correspondingly, in the capacity of initial formulas of the language $L(\Sigma_2^g)$ the formulas of the following two forms are taken: the formula $y^\sigma \delta_\sigma z^\sigma$ and the formula $(x_0^{\tau_0}, \dots, x_k^{\tau_k}) \varepsilon_\tau u^\tau$, where y^σ and z^σ are the variables of the first- or the second-order type σ and $x_i^{\tau_i}$ and u^τ are the variables of the first-order types τ_i and the second-order type $\tau \equiv [\tau_0, \dots, \tau_k]$, respectively.

These atomic formulas are interpreted on an evaluated system $((A, S_2^g), \gamma)$ (with a superstructure S_2^g of the signature Σ_2^g over a support A and an evaluation γ on the system $U \equiv (A, S_2^g)$) in the following generalized way: $\gamma(y^\sigma) \approx_\sigma \gamma(z^\sigma)$ and $(\gamma(x_0^{\tau_0}), \dots, \gamma(x_k^{\tau_k})) \varepsilon_\tau \gamma(u^\tau)$, where \approx_σ is a *generalized ratio of equality* and ε_τ is a *generalized ratio of belonging*. Generalized equalities and generalized belongings are connected with each other by the *initial principle of change of equals* (see axiom E4 from 2.3).

With respect to the signature Σ_2^g formulas φ in the language $L(\Sigma_2^g)$ are defined by common induction, when we start from the above-mentioned atomic formulas.

To give a semantics of the language $L(\Sigma_2^g)$ a *satisfaction of a formula φ on the system U with respect to the evaluation of variables γ* is defined according to the above-mentioned generalized interpretation of the atomic formulas.

The semantics for the language $L(\Sigma_2^g)$ differs both from the standard semantics (see [2, Appendix], [3, §16]) and from the Henkin semantics (see [2, Appendix], [3, §21], [4, 4], and [5, 6, 7]), which restricts the range of values of the evaluation $\gamma(x^\tau)$ for a variable x^τ of a second-order type τ by some subset of the set $\mathcal{P}(\tau(A))$ of the *terminal* $\tau(A)$.

The general material about second-order notions mentioned above is presented in Sections 2 and 3 of the paper. More specific material about the *generalized second-order Dedekind theory of real numbers* $Th_{R_2}^g$ and about the canonical model $R_0 \equiv R_2^g \equiv (\mathbb{R}, S_{R_2})$ is presented in Section 4.

In Section 5 we construct some inductive sequence of non-canonical models $R_i \equiv (\widehat{\mathbb{R}}_i, S_i)$, $1 \leq i \leq \omega_0$, with exponentially increasing powers. The basic tool for construction of these systems is the *infraproduct of collection of systems of the signature Σ_2^g* , different from the factorized ultraproduct à la Łoś. To prove that the systems R_i are models for $Th_{R_2}^g$ we use the simplified variant of the generalized infrafiltration theorem for the generalized second-order language $L(\Sigma_2^g)$ presented in [8, 9]. Note that the corresponding proof of the infrafiltration property for the standard second-order language $L(\Sigma_2^{st})$ do not “pass”.

The main results of this paper were announced in [10].

Further, to shorten the writings we use for the designation of a symbol-string ρ by a symbol-string σ the symbol-strings $\sigma \equiv \rho$ or $\rho \equiv \sigma$ (σ is a *designation for ρ*).

2 The type theory in the language of the signature with generalized equalities and belongings

2.1 Types

Fix the canonical set ω_0 of all natural numbers and its subset $\mathbb{N} \equiv \omega_0 \setminus \{0\}$ constructed in the Neumann–Bernays–Gödel (NBG) or Zermelo–Fraenkel (ZF) set theories or in the local theory of

sets (LTS) (see [11] and [12, 1.1, A.2, B.1]). Hereinafter ST denotes any of these set theories.

Define by induction the *semitypes* and the *types*:

1. 0 is the *semitype* and the *type*;
2. if τ is a type, then τ is the *semitype*;
3. if τ is a semitype, then $[\tau]$ is the *type*;
4. if τ_0, \dots, τ_k are semitypes and $k \geq 1$, then (τ_0, \dots, τ_k) is the *semitype*.

This definition is a slight modification of the corresponding definition from [3, § 20].

Further, instead of $[(\tau_0, \dots, \tau_k)]$ we shall write simply $[\tau_0, \dots, \tau_k]$; then the notation $[\tau_0, \dots, \tau_k]$ may be used for $k \geq 0$.

Semantics of semitypes and types will be explained in the next subsection.

Types 0 will be called the *first-order type*. If τ_0, \dots, τ_k are first-order types and $k \geq 0$ then $[\tau_0, \dots, \tau_k]$ will be called the *second-order type*.

For a type $\tau \equiv [\tau_0, \dots, \tau_k]$ with $k \geq 0$ the types τ_0, \dots, τ_k will be called the *parents of the type* τ and will be denoted by $p_0\tau, \dots, p_k\tau$, respectively. Consider the set $P(\tau) \equiv \{p_0\tau, \dots, p_k\tau\}$ of all *parents of the type* τ .

For the first-order type τ put formally $p\tau \equiv \tau$ and $P(\tau) \equiv \{p\tau\} = \{\tau\}$.

With any type τ we associate the *semitype* $\check{\tau}$ of the type τ as follows:

1. if τ is the first-order type, then $\check{\tau} \equiv \tau$;
2. if $\tau = [\tau_1]$ and τ_1 is a semitype, then $\check{\tau} \equiv \tau_1$.

In other words, the semitype of a type is obtained by omitting the square brackets.

2.2 Terminals over set and mappings

Define the *terminals* $\tau(A)$ of the *semitypes* τ over a set A by induction:

1. $0(A) \equiv A$;
2. if τ is a semitype, then $[\tau](A) \equiv \mathcal{P}(\tau(A))$, where \mathcal{P} denotes the operation of taking power-set of the intended set;
3. if τ_0, \dots, τ_k are semitypes, $k \geq 1$, then $(\tau_0, \dots, \tau_k)(A) \equiv \tau_0(A) \times \dots \times \tau_k(A)$.

Thus, for semitypes τ_0, \dots, τ_k with $k \geq 1$, for the type $\tau \equiv [\tau_0, \dots, \tau_k]$, and for its semitype $\check{\tau} = (\tau_0, \dots, \tau_k)$ the equalities $\tau(A) = \mathcal{P}(\tau_0(A) \times \dots \times \tau_k(A))$ and $\check{\tau}(A) = \tau_0(A) \times \dots \times \tau_k(A)$ are fulfilled.

Let $u : A \rightarrow B$ be a mapping from the set A to the set B . Define the *terminals* $\tau^m(u)$ of the *semitypes* τ over the mapping $u : A \rightarrow B$ by induction:

1. $0^m(u) \equiv u : A \rightarrow B$;
2. if τ is a semitype, then $[\tau]^m(u) : \mathcal{P}(\tau(A)) \rightarrow \mathcal{P}(\tau(B))$ is the mapping such that $[\tau]^m(u)(P) \equiv (\tau^m(u))[P] \equiv \{q \in \tau(B) \mid \exists p \in P (q = \tau^m(u)(p))\}$ for every $P \in \mathcal{P}(\tau(A))$;
3. if τ_0, \dots, τ_k are semitypes and $k \geq 1$, then

$$(\tau_0, \dots, \tau_k)^m(u) : \tau_0(A) \times \dots \times \tau_k(A) \rightarrow \tau_0(B) \times \dots \times \tau_k(B)$$

is the mapping such that

$$((\tau_0, \dots, \tau_k)^m(u))(p_0, \dots, p_k) \equiv (\tau_0^m(p_0), \dots, \tau_k^m(p_k))$$

for every $(p_0, \dots, p_k) \in \tau_0(A) \times \dots \times \tau_k(A)$.

2.3 The signature with generalized equalities and belongings and its language

A non-empty set Θ of types τ will be called the *type domain* if $\tau \in \Theta$ implies $p\tau \in \Theta$ for every parent $p\tau$ of the type τ . In the type domain Θ select the *belonging type subdomain* $\Theta_b \equiv \{\tau \in \Theta \mid \exists k \in \omega_0 \exists \tau_0, \dots, \tau_k \in \Theta (\tau = [\tau_0, \dots, \tau_k])\}$.

A collection $\Sigma_c \equiv \{\Sigma_c^\tau \mid \tau \in \Theta\}$ of *collections* $\Sigma_c^\tau \equiv \{\sigma_\omega^\tau \mid \omega \in \Omega_\tau\}$ of *constants* σ_ω^τ of the types τ will be called the *signature of constants of the type domain* Θ . Sets Ω_τ may be empty, and then $\Sigma_c^\tau = \emptyset$.

The constants σ_ω^0 of the first-order type 0 are called *individual* or *objective*. The constants of other types are called *predicate*.

A collection $\Sigma_e \equiv \{\delta_\tau \mid \tau \in \Theta\}$ of binary *predicate symbols of (generalized) equalities* δ_τ of the types τ will be called the *signature of (generalized) equalities of the type domain* Θ . It follows from the definition of the type domain that for every equality symbol δ_τ the collection Σ_e contains necessarily the equality symbols $\delta_{p\tau}$ for every parent $p\tau$ of the type τ .

A collection $\Sigma_b \equiv \{\varepsilon_\tau \mid \tau \in \Theta_b\}$ of binary *predicate symbols of (generalized) belongings* ε_τ of the types τ will be called the *signature of (generalized) belongings of the type domain* Θ .

A collection $\Sigma_v \equiv \{\Sigma_v^\tau \mid \tau \in \Theta\}$ of denumerable *sets* Σ_v^τ of *variables* x^τ, y^τ, \dots of the types τ will be called the *signature of variables of the type domain* Θ . The sets Σ_v^τ may be empty. The variables x^0, y^0, \dots of the first-order type 0 are called *individual* or *objective*. The variables of other types are called *predicate*.

Further, we shall always assume that for every type $\tau \in \Theta$ there are either constants or variables of this type.

The quadruple $\Sigma^g \equiv \Sigma_c \mid \Sigma_e \mid \Sigma_b \mid \Sigma_v$ will be called the *generalized signature* or the *signature with generalized equalities and belongings*.

The *language* $L(\Sigma^g)$ of the *generalized signature* Σ^g consists of:

1. all types τ from the type domain Θ ;
2. all members of all signatures from Σ^g ;
3. the logical symbols $\neg, \vee, \wedge, \Rightarrow, \forall, \exists$, and \exists ;
4. parenthesis.

If the type domain Θ contains first- and second-order types only and at least one second-order type, then we shall say that the signature Σ^g and the language $L(\Sigma^g)$ have the *second order* (see [2, Appendix], [4, 4]). In this case the notations Σ_2^g and $L(\Sigma_2^g)$ will be used.

2.4 Terms, formulas, and the type theory for the language of the generalized signature

Constants and variables of a type τ are called *terms of the type* τ of the language $L(\Sigma^g)$.

The *atomic formulas of the language* $L(\Sigma^g)$ are defined in the following way:

1. if q and r are terms of a type $\tau \in \Theta$, then $q \delta_\tau r$ is an *atomic formula*;
2. if τ_0, \dots, τ_k are types from Θ for $k \geq 0$, $\tau \equiv [\tau_0, \dots, \tau_k] \in \Theta_b$, $q_0^{\tau_0}, \dots, q_k^{\tau_k}$ are terms of the types τ_0, \dots, τ_k , respectively, and r^τ is a term of the type τ , then $(q_0^{\tau_0}, \dots, q_k^{\tau_k}) \varepsilon_\tau r^\tau$ is the *atomic formula*; in particular, for $k = 0$ the symbol-string $q_0^{\tau_0} \varepsilon_{[\tau_0]} r^{[\tau_0]}$ is the *atomic formula*.

The *formulas of the language* $L(\Sigma^g)$ are constructed from atomic ones with the use of connectives $\vee, \wedge, \neg, \Rightarrow$, quantifiers $\exists x^\tau$ and $\forall x^\tau$ with respect to the variables x^τ , and parenthesis.

The *logical axiom schemes* of the type theory in the language $L(\Sigma^g)$ of the generalized signature Σ^g are the schemes of the predicate calculus, where variables and terms substituting each other must be of the same type $\tau \in \Theta$.

In addition to these axiom schemes, consider the following *equality axioms* for the types $\tau \in \Theta$.

- E1.** $\forall x^\tau (x \delta_\tau x)$.
- E2.** $\forall x^\tau, y^\tau (x \delta_\tau y \Rightarrow y \delta_\tau x)$.
- E3.** $\forall x^\tau, y^\tau, z^\tau (x \delta_\tau y \wedge y \delta_\tau z \Rightarrow x \delta_\tau z)$.
- E4.** (The *initial principle of change of equals*.)

$$\forall x_0^{\tau_0}, y_0^{\tau_0}, \dots, x_k^{\tau_k}, y_k^{\tau_k}, u^\tau, v^\tau (x_0 \delta_{\tau_0} y_0 \wedge \dots \wedge x_k \delta_{\tau_k} y_k \wedge u \delta_\tau v \Rightarrow ((x_0, \dots, x_k) \varepsilon_\tau u \Leftrightarrow (y_0, \dots, y_k) \varepsilon_\tau v)), \text{ where } \tau \equiv [\tau_0, \dots, \tau_k].$$

The *inference rules* in the depicted type theory are:

$$\frac{\varphi, \varphi \Rightarrow \psi}{\psi} (MP) \quad \text{and} \quad \frac{\varphi(x^\tau)}{\forall x^\tau \varphi(x^\tau)} (Gen).$$

If there are non-logical axioms or axiom schemes written by second-order formulas of the language $L(\Sigma_2^g)$, then we shall say that a (*mathematical*) *generalized second-order theory* is given.

3 Mathematical systems of the signature Σ^g with generalized equalities and belongings

3.1 The definition of mathematical systems and their homomorphisms of the generalized signature Σ^g

Generalized systems. Let Σ^g be a fixed signature defined in 2.3. Fix also a set A . For the set A and the signature Σ^g consider the following collections:

1. $S_c \equiv \{S_c^\tau \mid \tau \in \Theta\}$ of collections $S_c^\tau \equiv \{s_\omega^\tau \mid \omega \in \Omega_\tau\}$ of *constant structures* $s_\omega^\tau \in \tau(A)$ of the types τ ;
2. $S_e \equiv \{\approx_\tau \mid \tau \in \Theta\}$ of *generalized ratios of equality* $\approx_\tau \subset \tau(A) \times \tau(A)$ of the types τ on the sets $\tau(A)$, containing the usual set-theoretic ratios of equality $=$ on the sets $\tau(X)$, i. e., such ratios \approx_τ that for every elements $r, s \in \tau(A)$ the equality $r = s$ implies the generalized equality $r \approx_\tau s$;
3. $S_b \equiv \{\in_\tau \mid \tau \in \Theta_b\}$ of *generalized ratios of belonging* $\in_\tau \subset \tilde{\tau}(A) \times \tau(A)$ of the types τ , containing the usual set-theoretic ratios of belonging \in from the sets $\tilde{\tau}(X)$ into the sets $\tau(X)$, i. e., such ratios \in_τ that for every elements $p \in \tilde{\tau}(A)$ and $P \in \tau(A)$ the belonging $p \in P$ implies the generalized belonging $p \in_\tau P$;
4. $S_v \equiv \{\tau(A) \mid \tau \in \Theta\}$ of the terminals $\tau(A)$ of the types τ over the set A .

The quadruple $S \equiv (S_c, S_e, S_b, S_v)$ of the above-mentioned collections will be called a *superstructure* of the signature Σ^g over the set A .

The pair $U \equiv (A, S)$ will be called a *mathematical system of the generalized signature Σ^g with the support (carrier) A and the superstructure S* . This notion is a generalization of the notion of an *algebraic system of the signature Σ_1* (see [13, § 15]).

The mathematical system $U \equiv (A, S)$ will be called also an *interpretation of the signature Σ^g on the support A* .

Further, for a type $\tau = [\tau_0, \dots, \tau_k]$ and elements $p \equiv (p(0), \dots, p(k))$, $q \equiv (q(0), \dots, q(k)) \in \check{\tau}(A) = \tau_0(A) \times \dots \times \tau_k(A)$ along with

$$p(0) \approx_{\tau_0} q(0) \wedge \dots \wedge p(k) \approx_{\tau_k} q(k)$$

we shall also write $p \approx_{\check{\tau}} q$.

The generalized equalities \approx_{τ} and the generalized belongings \in_{τ} admit some additional conditions. A system U will be called *balanced* if

$$\forall P, Q \in \tau(A) (P \approx_{\tau} Q \Leftrightarrow \forall p \in P \exists q \in Q (q \approx_{\check{\tau}} p) \wedge \forall q \in Q \exists p \in P (p \approx_{\check{\tau}} q)),$$

where $\tau_0, \dots, \tau_k \in \Theta$, $k \geq 0$ and $\tau \equiv [\tau_0, \dots, \tau_k] \in \Theta$. A system U will be called *regular* if $\forall p \in \check{\tau}(A) \forall P \in \tau(A) (p \in_{\tau} P \Leftrightarrow \exists q \in P (p \approx_{\check{\tau}} q))$, where $\tau_0, \dots, \tau_k \in \Theta$, $k \geq 0$, and $\tau \equiv [\tau_0, \dots, \tau_k] \in \Theta$. A system U will be called *extensional* if

$$\forall P, Q \in \tau(A) (P \approx_{\tau} Q \Leftrightarrow \forall p (p \in_{\tau} P \Rightarrow p \in_{\tau} Q) \wedge \forall q (q \in_{\tau} Q \Rightarrow q \in_{\tau} P)),$$

where $\tau \in \Theta_b$.

Generalized homomorphisms. Let $U \equiv (A, S)$ and $V \equiv (B, T)$ be systems of the signature Σ^g from 3.1. A mapping $u : A \rightarrow B$ in the considered set theory ST from the set A to the set B is called a *homomorphism of the signature Σ^g from the system U into the system V* if for every type $\tau \in \Theta$, every index $\omega \in \Omega_{\tau}$, every corresponding constant structure $s_{\omega}^{\tau} \in \tau(A)$ of the collection S_c , and every corresponding constant structure $t_{\omega}^{\tau} \in \tau(B)$ of the collection T_c the following properties are fulfilled:

1. if $\tau = 0$, then $\tau^m(u)(s_{\omega}^{\tau}) = u(s_{\omega}^{\tau}) = t_{\omega}^{\tau}$;
2. if $\tau \in \Theta_b$, then every generalized belonging $p \in_{\tau, A} s_{\omega}^{\tau}$ implies the corresponding generalized belonging $\check{\tau}^m(u)(p) \in_{\tau, B} t_{\omega}^{\tau}$ for every $p \in \check{\tau}(A)$.

3.2 Evaluations and models

An *evaluation on a system $U \equiv (A, S)$ of the signature Σ^g* is a mapping γ defined on the set of all variables of the signature Σ^g and associating with the variable x^{τ} of the type $\tau \in \Theta$ the element $\gamma(x^{\tau})$ of the terminal $\tau(X)$ (see [13, § 16], [3, 16.17]). The pair (U, γ) consisting of the system U of the signature Σ^g and the evaluation γ on U will be called an *evaluated mathematical system of the signature Σ^g* .

Define the *value $q[\gamma]$ of a term q with respect to the evaluation γ on the system U* in the following way (see [13, § 16], [14, § 6], [2, 2.2], [15, 2.5]): for a constant σ_{ω}^{τ} of a type $\tau \in \Theta$ put $\sigma_{\omega}^{\tau}[\gamma] \equiv s_{\omega}^{\tau}$ and for a variable x^{τ} of a type $\tau \in \Theta$ put $x^{\tau}[\gamma] \equiv \gamma(x^{\tau})$.

Define the *satisfaction (translation) of a formula φ of the language $L(\Sigma_2^g)$ on a system U of the signature Σ_2^g with respect to an evaluation γ* (in notation, $U \models \varphi[\gamma]$) by induction in the following way (see [2, 2.2], [15, 2.5], [3, 16.17], [12, A.1.3]):

1. if q and r are terms of a type $\tau \in \Theta$ and $\varphi \equiv (q \delta_{\tau} r)$, then $U \models \varphi[\gamma]$ is equivalent to $q[\gamma] \approx_{\tau} r[\gamma]$;
2. if τ_0, \dots, τ_k are types from Θ for $k \geq 0$, $\tau \equiv [\tau_0, \dots, \tau_k] \in \Theta$, q_0, \dots, q_k are terms of the types τ_0, \dots, τ_k , respectively, r is a term of the type τ , and $\varphi \equiv (q_0, \dots, q_k) \varepsilon_{\tau} r$, then $U \models \varphi[\gamma]$ iff $(q_0[\gamma], \dots, q_k[\gamma]) \in_{\tau} r[\gamma]$;
3. if $\varphi \equiv \neg\psi$, then $U \models \varphi[\gamma]$ iff $U \models \psi[\gamma]$ is not true;
4. if $\varphi \equiv (\psi \vee \xi)$, then $U \models \varphi[\gamma]$ iff $U \models \psi[\gamma]$ or $U \models \xi[\gamma]$;
5. if $\varphi \equiv (\psi \wedge \xi)$, then $U \models \varphi[\gamma]$ iff $U \models \psi[\gamma]$ and $U \models \xi[\gamma]$;

6. if $\varphi \equiv (\psi \Rightarrow \xi)$, then $U \models \varphi[\gamma]$ iff that $U \models \psi[\gamma]$ implies $U \models \xi[\gamma]$;
7. if $\varphi \equiv \exists x^\tau \psi$, then $U \models \varphi[\gamma]$ is equivalent to $U \models \psi[\gamma']$ for some evaluation γ' such that $\gamma'(y^\sigma) = \gamma(y^\sigma)$ for every variable $y^\sigma \neq x^\tau$;
8. if $\varphi \equiv \forall x^\tau \psi$, then $U \models \varphi[\gamma]$ is equivalent to $U \models \psi[\gamma']$ for every evaluation γ' such that $\gamma'(y^\sigma) = \gamma(y^\sigma)$ for every variable $y^\sigma \neq x^\tau$.

Let Φ be a set of formulas of the language $L(\Sigma_2^g)$. An evaluated mathematical system (U, γ) of the signature Σ_2^g will be called an *(evaluated) model for the set Φ* if $U \models \varphi[\gamma]$ for every formula $\varphi \in \Phi$ (see [13, §17]). A mathematical system U of the signature Σ_2^g will be called a *model for the set Φ* if an evaluated mathematical system (U, γ) is a model for the set Φ for every evaluation γ on U .

A model (U, γ) will be called *balanced, regular, extensional*, etc. if the system U is the same.

A model (U, γ) for a set Φ will be called *second-order* if at least one formula from Φ contains at least one second-order variable.

Remark that if a system $U \equiv (A, S)$ is considered in an axiomatic set theory, then the satisfaction of a closed formula φ of the language $L(\Sigma_2^g)$ with respect to any evaluation γ is reduced to correctness of the relativization φ^r of φ on the corresponding terminals of the support A in this set theory. Here the correctness of φ^r means that φ^r is a deducible formula in this axiomatic set theory.

Thus, if Φ consists of closed formulas only, then U is a model for Φ iff (U, γ) is a model for Φ for some (and, consequently, for any) evaluation γ .

In particular, since equality axioms E1–E4 are closed formulas, their relativizations E1^r–E4^r take the following forms:

$$\begin{aligned}
 E1^r &\equiv \forall x \in \tau(A) (x \approx_\tau x); \\
 E2^r &\equiv \forall x, y \in \tau(A) (x \approx_\tau y \Rightarrow y \approx_\tau x); \\
 E3^r &\equiv \forall x, y, z \in \tau(A) (x \approx_\tau y \wedge y \approx_\tau z \Rightarrow x \approx_\tau z); \\
 E4^r &\equiv \forall x_0, y_0 \in \tau_0(A) \dots \forall x_k, y_k \in \tau_k(A) \forall u, v \in \tau(A) (x_0 \approx_{\tau_0} y_0 \wedge \dots \wedge \\
 &\quad \wedge x_k \approx_{\tau_k} y_k \wedge u \approx_\tau v \Rightarrow ((x_0, \dots, x_k) \in_\tau u \Leftrightarrow (y_0, \dots, y_k) \in_\tau v)), \\
 &\quad \text{where } \tau \equiv [\tau_0, \dots, \tau_k], k \geq 0, \text{ and all types are in } \Theta.
 \end{aligned}$$

The satisfaction of formulas E1^r–E3^r means that all generalized equalities \approx_τ are equivalence relations on corresponding sets $\tau(A)$, and the satisfaction of formula E4^r means the initial principle of change of equals in the atomic formula with the generalized belonging \in_τ .

Further on, we shall say that a system U of the signature Σ_2^g has *true generalized equalities and belongings* if axioms E1–E4 from 2.3 are satisfied on U with respect to some (and, consequently, to any) evaluation γ . This means that formulas E1^r–E4^r are correct for the system U in the used set theory.

3.3 The generalized equality of values of evaluations and satisfiability

For every formula φ of the language $L(\Sigma_2^g)$ we define the formula φ^* by induction:

1. $\varphi^* \equiv \varphi$ for every atomic formula φ ;
2. $(\psi \wedge \xi)^* \equiv \psi^* \wedge \xi^*$;
3. $(\neg\psi)^* \equiv \neg\psi^*$;
4. $(\exists x^\tau \psi)^* \equiv \exists x^\tau \psi^*$;
5. $(\psi \vee \xi)^* \equiv \neg(\neg\psi^* \wedge \neg\xi^*)$;
6. $(\psi \Rightarrow \xi)^* \equiv \neg(\psi^* \wedge \neg\xi^*)$;

7. $(\forall x^\tau \psi)^* \equiv \neg(\exists x^\tau (\neg \psi^*)).$

A formula φ is said to be *normalizable* if for every mathematical Σ_2^g -system U and every evaluation γ on U the following condition holds: $U \models \varphi[\gamma] \Leftrightarrow U \models \varphi^*[\gamma]$.

Lemma 1. *Let formulas ψ and ξ be normalizable. Then formulas $\psi \wedge \xi$, $\neg \psi$, $\psi \vee \xi$, $\psi \Rightarrow \xi$, $\forall x^\tau \psi$, and $\exists x^\tau \psi$ are normalizable as well.*

The proof of this lemma uses the definition of satisfiability and some well known tautologies only, so it is omitted.

Propositon 1. *Every formula of the language $L(\Sigma_2^g)$ of the generalized second-order signature Σ_2^g is normalizable.*

Proof. Denote by Φ the set of all formulas of the language $L(\Sigma_2^g)$. The subset of the set Φ consisting of formulas containing at most $n \in \omega_0$ logical symbols \neg , \wedge , \Rightarrow , \vee , \exists , \forall , denote by Φ_n . It is clear that $\Phi = \bigcup(\Phi_n \mid n \in \omega_0)$.

Prove by the complete induction principle the following assertion $A(n)$: *every formula $\varphi \in \Phi$ is normalizable.*

If $n = 0$, then the formula φ is atomic, and so by the definition of the operation $\varphi \mapsto \varphi^*$ we have $\varphi^* \equiv \varphi$. Consequently, the assertion $A(0)$ is true.

Suppose that for all $m < n$ the assertion $A(m)$ is true. Let $\varphi \in \Phi_n$. If $\varphi \equiv \psi \wedge \xi$, $\varphi \equiv \neg \psi$, $\varphi \equiv \exists x^\tau \psi$, $\varphi \equiv \psi \vee \xi$, $\varphi \equiv \psi \Rightarrow \xi$, or $\varphi \equiv \forall x^\tau \psi$, then $\psi, \xi \in \Phi_{n-1}$. Therefore by the induction hypothesis, the formulas ψ and ξ are normalizable. By Lemma 1 the formula φ is normalizable. Hence the assertion $A(n)$ is true. \square

Propositon 2. *Let U be a mathematical system of the second-order signature Σ_2^g with true generalized equalities and belongings. Then for every formula φ of the language $L(\Sigma_2^g)$ and every evaluations γ and δ on the system U such that $\gamma(x^\tau) \approx_\tau \delta(x^\tau)$ for every variable x^τ of every type $\tau \in \Theta$ the properties $U \models \varphi[\gamma]$ and $U \models \varphi[\delta]$ are equivalent.*

Proof. The set of all formulas φ of the language $L(\Sigma_2^g)$ constructed by induction from the atomic formulas with the use of connectives \neg and \wedge and quantifier \exists denote by Ψ . The subset of the set Ψ consisting of formulas containing at most $n \in \omega_0$ logical symbols \neg , \wedge , and \exists denote by Ψ_n . It is clear that $\Psi = \bigcup(\Psi_n \mid n \in \omega_0)$.

Prove by the complete induction principle the assertion $A(n)$: *for every formula $\varphi \in \Psi_n$ and every mentioned evaluations γ and δ the assertion of the Proposition holds.*

Let $n = 0$ and $\varphi \in \Psi_0$. Then φ is an atomic formula. At first consider the atomic formula φ of the form $q^\tau \delta_\tau r^\tau$. Suppose that $q^\tau = x^\tau$ and $r^\tau = s_\omega^\tau$. Then $U \models \varphi[\gamma]$ is equivalent to $\gamma(x) \approx_\tau s_\omega^\tau$ and $U \models \varphi[\delta]$ is equivalent to $\delta(x) \approx_\tau s_\omega^\tau$. Since, by our condition, $\gamma(x) \approx_\tau \delta(x)$, then assuming $U \models \varphi[\gamma]$ and using axioms $E2^\tau$ and $E3^\tau$ we infer $U \models \varphi[\delta]$. The inverse inference is checked in the same way. For the terms q^τ and r^τ of other forms the reasons are quite similar.

Now, consider the atomic formula φ of the form $(q_0^{\tau_0}, \dots, q_k^{\tau_k}) \varepsilon_\tau r^k$ for the type $\tau \equiv [\tau_0, \dots, \tau_k] \in \Theta_b$. Assume that $q_\lambda^{\tau_\lambda} = x_\lambda^{\tau_\lambda}$ and $r^\tau = u^\tau$ for some variables x_λ and u . Then $U \models \varphi[\gamma]$ is equivalent to $(\gamma(x_0), \dots, \gamma(x_k)) \varepsilon_\tau \gamma(u)$ and $U \models \varphi[\delta]$ is equivalent to $(\delta(x_0), \dots, \delta(x_k)) \varepsilon_\tau \delta(u)$.

Suppose $U \models \varphi[\gamma]$. Since, by our condition, $\gamma(x_\lambda^{\tau_\lambda}) \approx_{\tau_\lambda} \delta(x_\lambda^{\tau_\lambda})$, then using axiom $E4^\tau$, we infer $U \models \varphi[\delta]$. The inverse inference is checked in the same way. For the terms $q_\lambda^{\tau_\lambda}$ and r^τ of other kinds the reasons are quite similar.

Assume that assertion $A(m)$ is true for every $m < n$. Let $\varphi \equiv \exists x^\tau \psi$. Then $\psi \in \Psi_{n-1}$. Let be given some evaluations γ and δ such that $\gamma(x^\tau) \approx_\tau \delta(x^\tau)$.

Suppose $U \models \varphi[\gamma]$. It is equivalent to $U \models \psi[\gamma']$ for some evaluation γ' such that $\gamma'(y) = \gamma(y)$ for any $y^\sigma \neq x^\tau$.

Define an evaluation δ' on U setting $\delta'(y) \equiv \delta(y)$ for every $y^\sigma \neq x^\tau$ and $\delta'(x) \equiv \gamma'(x)$. Then $\delta'(y) = \delta(y) \approx_\sigma \gamma(y) = \gamma'(y)$ and $\delta'(x) = \gamma'(x)$, i. e., $\delta'(x) \approx_\tau \gamma'(x)$.

Since $\delta' \approx \gamma'$ in the above indicated sense, by our condition, we conclude that $U \models \psi[\gamma'] \Leftrightarrow U \models \psi[\delta']$. Consequently, we obtain the property $U \models \psi[\delta']$. By construction, $\delta'(y) = \delta(y)$ for every $y^\sigma \neq x^\tau$. By the definition of satisfiability, we conclude that $U \models \varphi[\delta]$. The inverse inference of $U \models \varphi[\gamma]$ from $U \models \varphi[\delta]$ is established quite analogously.

Now, let $\varphi \equiv \psi \wedge \xi$. Then $\psi, \xi \in \Psi_{n-1}$, whence $U \models \psi[\gamma] \Leftrightarrow U \models \psi[\delta]$ and $U \models \xi[\gamma] \Leftrightarrow U \models \xi[\delta]$. Hence $(U \models \psi[\gamma] \wedge U \models \xi[\gamma]) \Leftrightarrow (U \models \psi[\delta] \wedge U \models \xi[\delta])$. Thus, $U \models \varphi[\gamma] \Leftrightarrow U \models \varphi[\delta]$.

Finally, let $\varphi \equiv \neg\psi$. Then $\psi \in \Psi_{n-1}$. Consequently, $U \models \psi[\gamma] \Leftrightarrow U \models \psi[\delta]$. From here $U \models \varphi[\gamma] \Leftrightarrow \neg(U \models \psi[\gamma]) \Leftrightarrow \neg(U \models \psi[\delta]) \Leftrightarrow U \models \varphi[\delta]$.

This proves that the assertion $A(n)$ is true. By the complete induction principle, the assertion $A(n)$ is true for every natural number $n \in \omega_0$, i. e., the assertion of the Proposition holds for every formula $\varphi \in \Psi$.

Now let φ be an arbitrary formula of the language $L(\Sigma_2^g)$. By virtue of Proposition 1 we have $U \models \varphi[\gamma] \Leftrightarrow U \models \varphi^*[\gamma]$ and $U \models \varphi[\delta] \Leftrightarrow U \models \varphi^*[\delta]$. By the definition of the operation $\varphi \mapsto \varphi^*$, we have $\varphi^* \in \Psi$. As was shown above, $U \models \varphi^*[\gamma] \Leftrightarrow U \models \varphi^*[\delta]$. As a result, we obtain the equivalence $U \models \varphi[\gamma] \Leftrightarrow U \models \varphi[\delta]$. \square

3.4 Examples of good models for the second-order equality axioms

Construct for axioms E1–E4 two regular, balanced, extensional, second-order models.

Take $\rho \equiv 0$, $\sigma \equiv [\rho]$, $\Theta \equiv \{\rho, \sigma\}$, $\Omega_\rho = \emptyset$, $\Omega_\sigma = \emptyset$, $\Sigma_c^\rho = \emptyset$, and $\Sigma_c^\sigma = \emptyset$. Then $\Sigma_e \equiv (\delta_\rho, \delta_\sigma)$, $\Theta_b = \{\sigma\}$, $\Sigma_b \equiv (\varepsilon_\tau \mid \tau \in \Theta_b)$, i. e., Σ_b consists of the symbol $\varepsilon_\sigma = \varepsilon_{[\rho]}$ only, and the collection $\Sigma_v \equiv (\Sigma_v^\tau \mid \tau \in \Theta)$ consists of a denumerable set Σ_v^ρ of variables x^ρ, y^ρ, \dots of the first-order type ρ and a denumerable set Σ_v^σ of variables $u^\sigma, v^\sigma, \dots$ of the second-order type σ .

Consider the signature $\Sigma \equiv \Sigma_c \mid \Sigma_e \mid \Sigma_b \mid \Sigma_v$. This language contains the three atomic formulas: $x^\rho \delta_\rho y^\rho$, $u^\sigma \delta_\sigma v^\sigma$ and $x^\rho \varepsilon_\sigma u^\sigma$.

Example 1. Take the set $\mathbb{Q} \equiv \mathbb{Z} \times (\mathbb{Z} \setminus \{0\})$ of all rational fractions $p \equiv \frac{m}{s}$ as the set A_1 . Since $\Omega_\rho = \Omega_\sigma = \emptyset$, there are no constants.

For fractions $p \equiv \frac{m}{s}$ and $q \equiv \frac{n}{t}$ put $p \approx_\rho q$ if $mt = ns$ in \mathbb{Z} . For sets $P, Q \in \mathcal{P}(A_1)$ put $P \approx_\sigma Q$ if $(\forall p \in P \exists q \in Q (p \approx_\rho q)) \wedge (\forall q \in Q \exists p \in P (q \approx_\rho p))$. It is clear that the generalized ratio of equality \approx_σ is wider than the usual set-theoretical ratio of equality $=$ in ST. For example, for $P_0 \equiv \{\frac{3}{8}, \frac{2}{3}\}$ and $Q_0 \equiv \{\frac{6}{16}, \frac{2}{3}, \frac{4}{6}\}$ we have $P_0 \approx_\sigma Q_0$ but $P_0 \neq Q_0$.

For a fraction $p \in A_1$ and a set $P \in \mathcal{P}(A_1)$ put $p \in_\sigma P$ if $\exists q \in A_1 (q \approx_\rho p \wedge q \in P)$. It is clear that the generalized ratio of belonging \in_σ is wider than the usual set-theoretical ratio of belonging \in in ST. For example, $\frac{6}{16} \in_\sigma P_0$ and $\frac{6}{9} \in_\sigma P_0$ but $\frac{6}{16} \notin P_0$ and $\frac{6}{9} \notin P_0$.

The collection of terminals $S_v^1 \equiv (\tau(A_1) \mid \tau \in \Theta)$ consists of the terminal $\rho(A_1) = A_1$ and the terminal $\sigma(A_1) = \mathcal{P}(A_1)$.

The constructed collections form the superstructure S_1 over the set A_1 . Consider the mathematical system $U_1 \equiv (A_1, S_1)$ of the signature Σ .

Example 2. Take the set of all closed segments p of straight lines on the plane as the set A_2 . Since $\Omega_\rho = \Omega_\sigma = \emptyset$, there are no constants.

For segments $p, q \in A_2$ put $p \approx_\rho q$ if q is obtained from p by some parallel transfer. For sets $P, Q \in \mathcal{P}(A_2)$ of segments put $P \approx_\sigma Q$ if

$$(\forall p \in P \exists q \in Q (p \approx_\rho q)) \wedge (\forall q \in Q \exists p \in P (q \approx_\rho p)).$$

For a segment $p \in A_2$ and a set of segments $P \in \mathcal{P}(A_2)$ put $p \in_\sigma P$ if $\exists q \in A_2 (q \approx_\rho p \wedge q \in P)$, i. e., the segment p can be transferred into the set P by some parallel transfer.

The collection of terminals $S_v^2 \equiv (\tau(A_2) \mid \tau \in \Theta)$ consists of the terminal $\rho(A_2) = A_2$ and the terminal $\sigma(A_2) = \mathcal{P}(A_2)$.

The constructed collections form the superstructure S_2 over the set A_2 . Consider the mathematical system $U_2 \equiv (A_2, S_2)$ of the signature Σ .

Propositon 1. *The above-constructed mathematical systems U_1 and U_2 are the regular, balanced, extensional, second-order models for equality axioms E1–E4.*

Proof. The correctness of the equality axioms is evident. The regularity follows from the definition. The same is true for the balance property.

Check the extensionality property. Let $P, Q \in \sigma(A) = \mathcal{P}(A)$. Assume $p \in P$. Then $p \in_{\sigma} P$. Suppose the right side of the extensionality formula. By condition we conclude $p \in_{\sigma} Q$. By the regularity property there exists an element $q \in Q$ such that $q \approx_{\rho} p$. The inverse finding of an element $p \in P$ for a given element $q \in Q$ such that $p \approx_{\rho} q$ is established quite similarly. In accordance with the definition of the equality \approx_{σ} we conclude that $P \approx_{\sigma} Q$. Thus, we have inferred the left side of the extensionality formula. It follows from the correctness of axiom E4^r that the left side implies the right one. \square

4 The generalized second-order Dedekind theory of real numbers

4.1 The signature for the generalized and the standard second-order Dedekind theories of real numbers

Consider the first-order type $\pi \equiv 0$, the second-order types $\varkappa \equiv [\pi]$, $\rho \equiv [\pi, \pi]$, and $\lambda \equiv [\pi, \pi, \pi]$ and the type domain $\Theta \equiv \Theta_{R2}^g \equiv \{\pi, \varkappa, \rho, \lambda\}$ with the belonging type subdomain $\Theta_b \equiv \{\varkappa, \rho, \lambda\}$.

Put $\Omega_{\pi} \equiv 2$, $\Omega_{\varkappa} \equiv \emptyset$, $\Omega_{\rho} \equiv 3$, $\Omega_{\lambda} \equiv 2$, and consider the collections

$$\begin{aligned} \Sigma_c^{\pi} &\equiv (\sigma_{\omega}^{\pi} \mid \omega \in \Omega_{\pi}) = (\sigma_0^{\pi}, \sigma_1^{\pi}), & \Sigma_c^{\varkappa} &\equiv (\sigma_{\omega}^{\varkappa} \mid \omega \in \Omega_{\varkappa}) = \emptyset, \\ \Sigma_c^{\rho} &\equiv (\sigma_{\omega}^{\rho} \mid \omega \in \Omega_{\rho}) = (\sigma_0^{\rho}, \sigma_1^{\rho}, \sigma_2^{\rho}), & \text{and } \Sigma_c^{\lambda} &\equiv (\sigma_{\omega}^{\lambda} \mid \omega \in \Omega_{\lambda}) = (\sigma_0^{\lambda}, \sigma_1^{\lambda}). \end{aligned}$$

They compose the signature of constants of the type domain Θ of the form $\Sigma_c = (\Sigma_c^{\tau} \mid \tau \in \Theta) = ((\sigma_0^{\pi}, \sigma_1^{\pi}), \emptyset, (\sigma_0^{\rho}, \sigma_1^{\rho}, \sigma_2^{\rho}), (\sigma_0^{\lambda}, \sigma_1^{\lambda}))$ containing the objective first-order constants σ_0^{π} and σ_1^{π} for denoting the *real numbers* 0 (*null*) and 1 (*unit*), respectively, the predicate second-order constants σ_0^{ρ} , σ_1^{ρ} , and σ_2^{ρ} for denoting the *ratio of negation*, the *ratio of inversion*, and the *ratio of order*, respectively, and the predicate second-order constants σ_0^{λ} and σ_1^{λ} for denoting the *ratio of addition* and the *ratio of multiplication*, respectively.

Further, along with σ_0^{π} , σ_1^{π} , σ_0^{ρ} , σ_1^{ρ} , σ_2^{ρ} , σ_0^{λ} , and σ_1^{λ} we shall simply write 0, 1, $-$, $/$, \leq , $+$, and \cdot , respectively.

Take the signature of the generalized equalities of the type domain Θ of the form $\Sigma_e \equiv (\delta_{\tau} \mid \tau \in \Theta) = (\delta_{\pi}, \delta_{\varkappa}, \delta_{\rho}, \delta_{\lambda})$ containing the first-order equality δ_{π} , and the second-order equalities $\delta_{[\pi]}$, $\delta_{[\pi, \pi]}$, and $\delta_{[\pi, \pi, \pi]}$.

Take the signature of the generalized belongings of the type domain Θ of the form $\Sigma_b \equiv (\varepsilon_{\tau} \mid \tau \in \Theta_b) = (\varepsilon_{\varkappa}, \varepsilon_{\rho}, \varepsilon_{\lambda})$.

Finally, take a denumerable set Σ_v^{π} of objective variables x^{π}, y^{π}, \dots of the first-order type π and denumerable sets Σ_v^{\varkappa} , Σ_v^{ρ} , and Σ_v^{λ} of predicate variables $u^{\varkappa}, v^{\varkappa}, \dots$, $u^{\rho}, v^{\rho}, \dots$, and $u^{\lambda}, v^{\lambda}, \dots$ of the second-order types \varkappa , ρ , and λ , respectively.

They form the signature $\Sigma_v \equiv (\Sigma_v^{\tau} \mid \tau \in \Theta) = (\Sigma_v^{\pi}, \Sigma_v^{\varkappa}, \Sigma_v^{\rho}, \Sigma_v^{\lambda})$ of variables of the type domain Θ .

Consider the *generalized signature* $\Sigma_{R2}^g \equiv \Sigma_c \mid \Sigma_e \mid \Sigma_b \mid \Sigma_v$ and the corresponding language $L(\Sigma_{R2}^g)$. Terms p, q, r, s, \dots of this language are constants and variables only; the atomic equality formulas have the forms $q^{\pi} \delta_{\pi} r^{\pi}$, $q^{\varkappa} \delta_{\varkappa} r^{\varkappa}$, $q^{\rho} \delta_{\rho} r^{\rho}$, and $q^{\lambda} \delta_{\lambda} r^{\lambda}$. Respectively, the atomic belonging formulas have the forms $q^{\pi} \varepsilon_{\varkappa} r^{\varkappa}$, $(p^{\pi}, q^{\pi}) \varepsilon_{\rho} r^{\rho}$, and $(p^{\pi}, q^{\pi}, r^{\pi}) \varepsilon_{\lambda} s^{\lambda}$.

Further, along with x^{π} , y^{π} , and δ_{π} we shall simply write x , y , and δ , respectively.

Along with the generalized signature Σ_{R2}^g we consider the *standard signature* $\Sigma_{R2}^{st} \equiv \Sigma_c | \Sigma_e^{st} | \Sigma_b^{st} | \Sigma_v$, where in the signature of the standard equalities $\Sigma_e^{st} \equiv (\delta_\tau^{st} | \tau \in \Theta)$ the type equalities δ_τ^{st} are one and the same *standard equality* δ^{st} and in the signature of the standard belongings $\Sigma_b^{st} \equiv (\varepsilon_\tau^{st} | \tau \in \Theta_b)$ the type belongings ε_τ^{st} are one and the same *standard belonging* ε^{st} .

Respectively, this signature Σ_{R2}^{st} generates the standard language $L(\Sigma_{R2}^{st})$ with atomic equality formulas of the forms $q^\pi \delta^{st} r^\pi$, $q^\pi \delta^{st} r^\pi$, $q^\rho \delta^{st} r^\rho$, and $q^\lambda \delta^{st} r^\lambda$ and with atomic belonging formulas of the forms $q^\pi \varepsilon^{st} r^\pi$, $(p^\pi, q^\pi) \varepsilon^{st} r^\rho$, and $(p^\pi, q^\pi, r^\pi) \varepsilon^{st} s^\lambda$ for all terms p, q, r, s, \dots

4.2 The axiomatics for the generalized and the standard second-order Dedekind theories of real numbers

The signature Σ_{R2}^g gives the opportunity to define the language $L(\Sigma_{R2}^g)$ and to construct the desired models of the generalized second-order theory of real numbers, but the absence of functional variables in this signature makes the writing of generalized axioms for this theory very unusual. Only the names of these axioms placed in round brackets clarify their customary sense.

The *axioms of the generalized second-order Dedekind theory of real numbers* are the following ones.

A1 (the *existence and functionality of the negation*).

$$(\forall x \exists y ((x, y) \varepsilon_\rho -)) \wedge (\forall x, y, y' ((x, y) \varepsilon_\rho -) \wedge ((x, y') \varepsilon_\rho -) \Rightarrow y \delta y').$$

A2 (the *existence and functionality of the addition*).

$$(\forall x, y \exists z ((x, y, z) \varepsilon_\lambda +)) \wedge (\forall x, y, z, z' ((x, y, z) \varepsilon_\lambda +) \wedge ((x, y, z') \varepsilon_\lambda +) \Rightarrow z \delta z').$$

A3 (the *existence and functionality of the inversion*).

$$(\forall x (\neg(x \delta 0) \Rightarrow \exists y ((x, y) \varepsilon_\rho /))) \wedge (\forall x, y ((x, y) \varepsilon_\rho / \Rightarrow \neg(x \delta 0))) \wedge \\ \wedge (\forall x, y, y' ((x, y) \varepsilon_\rho /) \wedge ((x, y') \varepsilon_\rho /) \Rightarrow y \delta y').$$

A4 (the *existence and functionality of the multiplication*).

$$(\forall x, y \exists z ((x, y, z) \varepsilon_\lambda \cdot)) \wedge (\forall x, y, z, z' ((x, y, z) \varepsilon_\lambda \cdot) \wedge ((x, y, z') \varepsilon_\lambda \cdot) \Rightarrow z \delta z').$$

The appearance of axioms A1–A4 in this list is directly impelled by the absence of functional variables in the signature Σ_{R2}^g .

A5 (the *non-equality of the unit and the null*). $\neg(1 \delta 0)$.

A6 (the *associativity of the addition*).

$$\forall x, y, z \forall u_1, u_2, v_1, v_2 (((x, y, u_1) \varepsilon_\lambda +) \wedge ((u_1, z, u_2) \varepsilon_\lambda +) \wedge \\ \wedge ((y, z, v_1) \varepsilon_\lambda +) \wedge ((x, v_1, v_2) \varepsilon_\lambda +) \Rightarrow u_2 \delta v_2).$$

The writing of axiom A6 in the common way: $\forall x, y, z (((x + y) + z) \delta (x + (y + z)))$.

A7 (the *neutrality of the null*).

$$\forall x \forall u, v (((x, 0, u) \varepsilon_\lambda +) \Rightarrow u \delta x) \wedge (((0, x, v) \varepsilon_\lambda +) \Rightarrow v \delta x).$$

A8 (the *elimination of the negation*).

$$\forall x \forall u_1, u_2, v_1, v_2 (((x, u_1) \varepsilon_\rho -) \wedge ((x, u_1, u_2) \varepsilon_\lambda +) \Rightarrow u_2 \delta 0) \wedge \\ \wedge (((x, v_1) \varepsilon_\rho -) \wedge ((v_1, x, v_2) \varepsilon_\lambda +) \Rightarrow v_2 \delta 0).$$

A9 (the *commutativity of the addition*).

$$\forall x, y \forall u, v ((x, y, u) \varepsilon_\lambda +) \wedge ((y, x, v) \varepsilon_\lambda +) \Rightarrow u \delta v).$$

A10 (the *right distributivity of the multiplication with respect the addition*).

$$\begin{aligned} \forall x, y, z \forall u_1, u_2, v_1, v_2, v_3 & (((y, z, u_1) \varepsilon_\lambda +) \wedge ((x, u_1, u_2) \varepsilon_\lambda \cdot) \wedge \\ & \wedge ((x, y, v_1) \varepsilon_\lambda \cdot) \wedge ((x, z, v_2) \varepsilon_\lambda \cdot) \wedge ((v_1, v_2, v_3) \varepsilon_\lambda +) \Rightarrow u_2 \delta v_3). \end{aligned}$$

The writing of this axiom in the common way: $\forall x, y, z ((x \cdot (y + z)) \delta (x \cdot y + x \cdot z))$.

A11 (the *left distributivity of the multiplication with respect the addition*).

$$\begin{aligned} \forall x, y, z \forall u_1, u_2, v_1, v_2, v_3 & (((x, y, u_1) \varepsilon_\lambda +) \wedge ((u_1, z, u_2) \varepsilon_\lambda \cdot) \wedge \\ & \wedge ((x, z, v_1) \varepsilon_\lambda \cdot) \wedge ((y, z, v_2) \varepsilon_\lambda \cdot) \wedge ((v_1, v_2, v_3) \varepsilon_\lambda +) \Rightarrow u_2 \delta v_3). \end{aligned}$$

A12 (the *associativity of the multiplication*).

$$\begin{aligned} \forall x, y, z \forall u_1, u_2, v_1, v_2 & (((x, y, u_1) \varepsilon_\lambda \cdot) \wedge ((u_1, z, u_2) \varepsilon_\lambda \cdot) \wedge \\ & \wedge ((y, z, v_1) \varepsilon_\lambda \cdot) \wedge ((x, v_1, v_2) \varepsilon_\lambda \cdot) \Rightarrow u_2 \delta v_2). \end{aligned}$$

A13 (the *neutrality of the unit*).

$$\forall x \forall u, v (((x, 1, u) \varepsilon_\lambda \cdot) \Rightarrow u \delta x) \wedge (((1, x, v) \varepsilon_\lambda \cdot) \Rightarrow v \delta x).$$

A14 (the *elimination of the inversion*).

$$\begin{aligned} \forall x \forall u_1, u_2, v_1, v_2 & (-(x \delta 0) \Rightarrow (((x, u_1) \varepsilon_\rho /) \wedge \\ & \wedge ((x, u_1, u_2) \varepsilon_\lambda \cdot) \Rightarrow u_2 \delta 1) \wedge (((x, v_1) \varepsilon_\rho /) \wedge ((v_1, x, v_2) \varepsilon_\lambda \cdot) \Rightarrow v_2 \delta 1)). \end{aligned}$$

The writing of A14 in the common way is the following:

$$\forall x (-(x \delta 0) \Rightarrow (x \cdot (x^{-1}) \delta 1) \wedge ((x^{-1}) \cdot x \delta 1)).$$

A15 (the *commutativity of the multiplication*).

$$\forall x, y \forall u, v (((x, y, u) \varepsilon_\lambda \cdot) \wedge ((y, x, v) \varepsilon_\lambda \cdot) \Rightarrow u \delta v).$$

Further, along with $(x, y) \varepsilon_\rho \leq$ we shall write $x \leq y$ as well. It gives the opportunity to write the subsequent axioms in a more customary form.

A16 (the *reflexivity of the order*). $\forall x (x \leq x)$.

By E4 we get $x \delta y \Rightarrow (x \leq x \Leftrightarrow x \leq y)$. Applying A16, we conclude that $x \delta y \vdash x \leq y$.

A17 (the *antisymmetry of the order*). $\forall x, y ((x \leq y) \wedge (y \leq x)) \Rightarrow x \delta y$.

A18 (the *transitivity of the order*). $\forall x, y, z ((x \leq y) \wedge (y \leq z)) \Rightarrow x \leq z$.

A19 (the *linearity of the order*). $\forall x, y ((x \leq y) \vee (y \leq x))$.

A20 (the *compatibility of the addition and the order*).

$$\forall x, y, z \forall u, v (x \leq y \Rightarrow (((x, z, u) \varepsilon_\lambda +) \wedge ((y, z, v) \varepsilon_\lambda +) \Rightarrow u \leq v)).$$

A21 (the *compatibility of the multiplication and the order*).

$$\forall x, y \forall u ((x \geq 0) \wedge (y \geq 0) \Rightarrow (((x, y, u) \varepsilon_\lambda \cdot) \Rightarrow u \geq 0)).$$

A22 (the *existence of Dedekind cuts*).

$$\begin{aligned} \forall u^\times, v^\times & ((\exists x (x \varepsilon_\times u^\times)) \wedge (\exists y (y \varepsilon_\times v^\times)) \wedge \\ & \wedge (\forall z ((z \varepsilon_\times u^\times) \vee (z \varepsilon_\times v^\times))) \wedge (\forall x, y ((x \varepsilon_\times u^\times) \wedge (y \varepsilon_\times v^\times) \Rightarrow x \leq y)) \Rightarrow \\ & \Rightarrow (\exists z \forall x, y ((x \varepsilon_\times u^\times) \wedge (y \varepsilon_\times v^\times) \Rightarrow (x \leq z) \wedge (z \leq y))))). \end{aligned}$$

Consider the following generalized *extensionality properties*.

PE1. $\forall u^\varkappa, v^\varkappa (u^\varkappa \delta_\varkappa v^\varkappa \Leftrightarrow \forall x (x \varepsilon_\varkappa u^\varkappa \Leftrightarrow x \varepsilon_\varkappa v^\varkappa))$.

PE2. $\forall u^\rho, v^\rho (u^\rho \delta_\rho v^\rho \Leftrightarrow \forall x, y ((x, y) \varepsilon_\rho u^\rho \Leftrightarrow (x, y) \varepsilon_\rho v^\rho))$.

PE3. $\forall u^\lambda, v^\lambda (u^\lambda \delta_\lambda v^\lambda \Leftrightarrow \forall x, y, z ((x, y, z) \varepsilon_\lambda u^\lambda \Leftrightarrow (x, y, z) \varepsilon_\lambda v^\lambda))$.

The theory determined by the language $L(\Sigma_{R2}^g)$ and the set of axioms $\Psi_2^g \equiv \{\text{E1–E4, A1–A22, PE1–PE3}\}$ can be called the *generalized second-order Dedekind theory of real numbers*. It will be denoted by Th_{R2}^g .

Respectively, in the language $L(\Sigma_{R2}^{st})$ we can write formulas $\text{E1}^{st}\text{–E4}^{st}$, $\text{A1}^{st}\text{–A22}^{st}$, $\text{PE1}^{st}\text{–PE3}^{st}$, which are obtained from the corresponding formulas $\text{E1–E4, A1–A22, PE1–PE3}$ of the language $L(\Sigma_{R2}^g)$ by the substitution of the generalized type equalities and belongings δ_τ and ε_τ by the standard ones δ^{st} and ε^{st} , respectively.

The theory determined by the language $L(\Sigma_{R2}^{st})$ and axioms $\text{E1}^{st}\text{–E4}^{st}$, $\text{A1}^{st}\text{–A22}^{st}$, $\text{PE1}^{st}\text{–PE3}^{st}$ can be called the *standard second-order Dedekind theory of real numbers*. It will be denoted by Th_{R2}^{st} .

4.3 The canonical generalized and standard second-order Dedekind real axes

Consider the canonical set \mathbb{R} of all real numbers constructed in the considered set theory ST (see, e.g., [12, 1.4] for NBG set theory and [11] and [12, B.1] for the LTS).

For the set \mathbb{R} and the signature Σ_{R2}^g consider the collections

$$\begin{aligned} S_c^\pi &\equiv (s_\omega^\pi \mid \omega \in \Omega_\pi) = (s_0^\pi, s_1^\pi), & S_c^\varkappa &\equiv (s_\omega^\varkappa \mid \omega \in \Omega_\varkappa) = \emptyset, \\ S_c^\rho &\equiv (s_\omega^\rho \mid \omega \in \Omega_\rho) = (s_0^\rho, s_1^\rho, s_2^\rho), & \text{and } S_c^\lambda &\equiv (s_\omega^\lambda \mid \omega \in \Omega_\lambda) = (s_0^\lambda, s_1^\lambda). \end{aligned}$$

They compose the collection of constants structures

$$S_c = (S_c^\tau \mid \tau \in \Theta) = ((s_0^\pi, s_1^\pi), \emptyset, (s_0^\rho, s_1^\rho, s_2^\rho), (s_0^\lambda, s_1^\lambda))$$

containing the constant structures $s_0^\pi, s_1^\pi \in \pi(\mathbb{R}) = \mathbb{R}$ which are the *neutral real numbers*, the constant structures $s_0^\rho, s_1^\rho, s_2^\rho \in \rho(\mathbb{R}) = \mathcal{P}(\mathbb{R}^2)$, which are the *ratio of negation*, the *ratio of inversion*, and the *ratio of order on* \mathbb{R} , respectively, and the constant structures $s_0^\lambda, s_1^\lambda \in \lambda(\mathbb{R}) = \mathcal{P}(\mathbb{R}^3)$ which are the *ratio of addition* and the *ratio of multiplication on* \mathbb{R} , respectively.

Further, along with $s_0^\pi, s_1^\pi, s_0^\rho, s_1^\rho, s_2^\rho, s_0^\lambda$, and s_1^λ we shall simply write $0_{\mathbb{R}}, 1_{\mathbb{R}}, -_{\mathbb{R}}, /_{\mathbb{R}}, \leq_{\mathbb{R}}, +_{\mathbb{R}}$, and $\cdot_{\mathbb{R}}$, respectively.

Consider the collection of the equality ratios of the form

$$S_e \equiv (\approx_\tau \mid \tau \in \Theta) = (\approx_\pi, \approx_\varkappa, \approx_\rho, \approx_\lambda) \equiv (= \mid \mathbb{R}^2, = \mid \mathcal{P}(\mathbb{R})^2, = \mid \mathcal{P}(\mathbb{R}^2)^2, = \mid \mathcal{P}(\mathbb{R}^3)^2)$$

containing in the capacity of the first-order equality ratio \approx_π and of the second-order equality ratios $\approx_\varkappa, \approx_\rho$, and \approx_λ the restrictions on the indicated sets one and the same set-theoretical equality in ST.

Consider the collection of the belonging ratios of the form

$$S_b \equiv (\in_\tau \mid \tau \in \Theta) = (\in_\varkappa, \in_\rho, \in_\lambda) \equiv (\in \mid (\mathbb{R} \times \mathcal{P}(\mathbb{R})), \in \mid (\mathbb{R}^2 \times \mathcal{P}(\mathbb{R}^2)), \in \mid (\mathbb{R}^3 \times \mathcal{P}(\mathbb{R}^3)))$$

containing in the capacity of the belonging ratios \in_\varkappa, \in_ρ , and \in_λ the restrictions on the indicated sets one and the same set-theoretical belonging ratio \in in ST.

Finally, take the collection of the terminals over the set \mathbb{R} of the form

$$S_v \equiv (\tau(\mathbb{R}) \mid \tau \in \Theta) = (\pi(\mathbb{R}), \varkappa(\mathbb{R}), \rho(\mathbb{R}), \lambda(\mathbb{R})) = (\mathbb{R}, \mathcal{P}(\mathbb{R}), \mathcal{P}(\mathbb{R}^2), \mathcal{P}(\mathbb{R}^3)).$$

These collections compose the superstructure $S_{R2} \equiv \{S_c, S_e, S_b, S_v\}$ of the signature Σ_{R2}^g . The system (\mathbb{R}, S_{R2}) of the signature Σ_{R2}^g can be called the *canonical generalized second-order Dedekind real axis in ST*. It will be denoted by R_2^g .

Consider an evaluation ζ on the system R_2^g such that $\zeta(x) \in \pi(\mathbb{R}) = \mathbb{R}$, $\zeta(u^\varkappa) \in \varkappa(\mathbb{R}) = \mathcal{P}(\mathbb{R})$, $\zeta(u^\rho) \in \rho(\mathbb{R}) = \mathcal{P}(\mathbb{R}^2)$, and $\zeta(u^\lambda) \in \lambda(\mathbb{R}) = \mathcal{P}(\mathbb{R}^3)$. Thus, we get the evaluated system (R_2^g, ζ) .

The above constructed superstructure S_{R2} is also the superstructure of the signature Σ_{R2}^{st} . Therefore the system (\mathbb{R}, S_{R2}) is also the system of the signature Σ_{R2}^{st} . It can be called the *canonical standard second-order Dedekind real axis in ST*. It will be denoted by R_2^{st} .

The evaluation ζ on the system R_2^g considered above is also an evaluation on the system R_2^{st} . Therefore we may consider the evaluated system (R_2^{st}, ζ) .

Let B be a set and T_{R2}^{st} be a superstructure on B of the signature Σ_{R2}^{st} . Consider the system $V \equiv (B, T_{R2}^{st})$ and some evaluation η on V . For the evaluated system (V, η) we shall use the following designations: $0_B \equiv \sigma_0^\circ[\eta]$, $1_B \equiv \sigma_1^\tau[\eta]$, $-_B \equiv \sigma_0^\rho[\eta]$, $/_B \equiv \sigma_1^\rho[\eta]$, $\leqslant_B \equiv \sigma_2^\rho[\eta]$, $+_B \equiv \sigma_0^\lambda[\eta]$, and $\cdot_B \equiv \sigma_1^\lambda[\eta]$.

The (*standard*) *satisfaction* $U \models_{st} \varphi[\eta]$ of a formula φ of the language $L(\Sigma_{R2}^{st})$ on the system V of the signature Σ_{R2}^{st} with respect to the evaluation η differs from the (generalized) satisfaction from 3.2 only in the first two points:

- 1'. if q and r are terms of a type $\tau \in \Theta$ and $\varphi \equiv (q \delta^{st} r)$, then $V \models_{st} \varphi[\eta]$ is equivalent to $q[\eta] = r[\eta]$;
- 2'. if τ_0, \dots, τ_k are types from Θ for $k \geqslant 0$, $\tau \equiv [\tau_0, \dots, \tau_k] \in \Theta$, q_0, \dots, q_k are terms of the types τ_0, \dots, τ_k , respectively, r is a term of the type τ , and $\varphi \equiv (q_0, \dots, q_k) \varepsilon^{st} r$, then $U \models \varphi[\eta]$ iff $(q_0[\eta], \dots, q_k[\eta]) \in r[\eta]$.

Let Φ be a set of formulas of the language $L(\Sigma_{R2}^{st})$. As in 3.3 the evaluated system (V, η) of the signature Σ_{R2}^{st} is called a *standard model for the set Φ* if $V \models_{st} \varphi[\eta]$ for every $\varphi \in \Phi$.

Now we can formulate some initial theorem about the standard R_2^{st} and the generalized R_2^g Dedekind real axes.

Theorem 1.

1. *The mathematical system R_2^{st} is a standard model for the theory Th_{R2}^{st} in the language $L(\Sigma_{R2}^{st})$.*
2. *The mathematical system R_2^g is a (generalized) model for the theory Th_{R2}^g in the language $L(\Sigma_{R2}^g)$.*

Proof. 1. Note that all the axioms from the set Ψ_2^g are closed formulas. Therefore the satisfaction $R_2^{st} \models \alpha[\zeta]$ for $\alpha \in \Psi_2^g$ means the deducibility of the relativization α^r of α on \mathbb{R} in the considered axiomatic set theory ST. But the corresponding deducibility of every α^r is very well demonstrated in mathematical literature (see, for example, [16, 1, 17, 18, 19, 12]).

2. This assertion follows directly from assertion 1 by virtue of the inclusions $|\tau(\mathbb{R}) \times \tau(\mathbb{R}) \subset \approx_\tau$ and $\in |\tilde{\tau}(\mathbb{R}) \times \tau(\mathbb{R}) \subset \in_\tau$ from 3.2, where the left parts of the inclusions are the restrictions of the usual set-theoretical ratios $=$ and \in on the indicated sets. \square

The models from Theorem 1 are called *canonical*.

It is well known that the theory Th_{R2}^{st} is categorical. On the contrary, we shall prove that the theory Th_{R2}^g is non-categorical. More exactly, using the initial canonical model R_2^g with the support \mathbb{R} we shall prove the existence of some non-canonical models for the theory Th_{R2}^g having arbitrary large powers.

This statement can be proven with the help of the generalized infrafiltration theorem (see, e. g., [9] or [12, C.3.2]). But to make the paper self-contained we prefer to prove here some more simple variant of the generalized infrafiltration theorem than it is presented in the indicated works.

5 The infraproduct construction of evaluated systems of the signature Σ_2^g

5.1 Infraproducts of collections of evaluated systems of the signature Σ_2^g

Let F be a fixed set and $(U_f \mid f \in F)$ be a fixed collection of mathematical systems of the signature Σ_2^g with true generalized equalities and belongings.

By definition, $U_f \equiv (A_f, S_f)$. Consider the set $A \equiv \prod(A_f \mid f \in F)$.

Let $\tau \equiv [\tau_0, \dots, \tau_k]$ be a second-order type and $k \geq 0$. If $\mu \in k + 1$, then $\tau_\mu = 0$. Thus, we see that $\tau_\mu(A) = A = \prod(A_f \mid f \in F) = \prod(\tau_\mu(A_f) \mid f \in F)$. For elements $p \in \check{\tau}(A) \equiv \tau_0(A) \times \dots \times \tau_k(A) = A^{k+1}$ and $f \in F$ define the element $p(f) \in \check{\tau}(A_f) = \tau_0(A_f) \times \dots \times \tau_k(A_f) = A_f^{k+1}$ setting $p(f)(\mu) \equiv p(\mu)(f)$ for every $\mu \in k + 1$.

For elements $P \subset \check{\tau}(A)$ and $f \in F$ define the element $P\langle f \rangle \subset \check{\tau}(A_f)$ setting $P\langle f \rangle \equiv \{\xi \in \check{\tau}(A_f) \mid \exists p \in P (p(f) = \xi)\}$.

Let \mathcal{D} be a subset of the set $\mathcal{P}(F)$, i. e., an ensemble on F . Define some superstructure S of the signature Σ_2^g over the set A .

First, define constant structures $s_\omega^\tau \in \tau(A)$ for $\tau \in \Theta$ and $\omega \in \Omega_\tau$.

If τ is a first-order type, then $\tau(A) = \prod(\tau(A_f) \mid f \in F)$. Therefore define $s_\omega^\tau \in \tau(A)$ setting $s_\omega^\tau(f) \equiv s_{\omega_f}^\tau$ for every $f \in F$.

Put $s_\omega^\tau \equiv \{p \in \check{\tau}(A) \mid \forall f \in F (p(f) \in s_{\omega_f}^\tau)\}$ if $\tau \equiv [\tau_0, \dots, \tau_k]$ is a second-order type.

As a result, we obtain the collections $S_c^\tau \equiv \{s_\omega^\tau \mid \omega \in \Omega_\tau\}$ and the collection $S_c \equiv \{S_c^\tau \mid \tau \in \Theta\}$.

Now define generalized equality ratios $\approx_\tau \subset \tau(A) \times \tau(A)$. If τ is the first-order type, then for $p, q \in \tau(A)$ put $p \approx_\tau q$ if $\exists G \in \mathcal{D} \forall g \in G (p(g) \approx_{\tau, g} q(g))$.

If $\tau \equiv [\tau_0, \dots, \tau_k]$ is a second-order type, then for $P, Q \subset \check{\tau}(A)$ put $P \approx_\tau Q$ if $\exists G \in \mathcal{D} \forall g \in G (P\langle g \rangle \approx_{\tau, g} Q\langle g \rangle)$.

As a result, we obtain the collection $S_e \equiv \{\approx_\tau \mid \tau \in \Theta\}$.

Now define generalized belonging ratios $\in_\tau \subset \check{\tau}(A) \times \tau(A)$.

By definition, $\tau = [\tau_0, \dots, \tau_k]$ for some $\tau_0, \dots, \tau_k \in \Theta$. For $p \in \check{\tau}(A)$ and $P \subset \check{\tau}(A)$ put $p \in_\tau P$ if $\exists G \in \mathcal{D} \forall g \in G (p(g) \in_{\tau, g} P\langle g \rangle)$. Note that the usage of a generalized belonging ratio was explored in the forcing method in the form $x \in_p y$ (see, e. g., [15, 9.8]).

Thus, we obtain the collection $S_b \equiv \{\in_\tau \mid \tau \in \Theta_b\}$.

Consider also the collection $S_v \equiv \{\tau(A) \mid \tau \in \Theta\}$ consisting of the τ -terminals of the set A .

The constructed collections compose the superstructure $S \equiv \{S_c, S_e, S_b, S_v\}$ over the set A . Therefore we can consider the mathematical system $U \equiv (A, S)$ of the signature Σ_2^g . It will be called the *infra- \mathcal{D} -product of the collection of mathematical systems $(U_f \mid f \in F)$ of the generalized second-order signature Σ_2^g* and will be denoted by $\text{infra-}\mathcal{D}\text{-prod}(U_f \mid f \in F)$.

An ensemble \mathcal{D} on F is called a *filter on F* if it has the following properties:

1. $\forall G, H \in \mathcal{D} (G \cap H \in \mathcal{D})$;
2. $\forall G \in \mathcal{D} \forall H \in \mathcal{P}(F) (G \subset H \Rightarrow H \in \mathcal{D})$.

A filter \mathcal{D} is called *proper* if $\mathcal{D} \neq \mathcal{P}(F)$. A proper filter \mathcal{D} is called an *ultrafilter* if for any proper filter \mathcal{E} on F such that $\mathcal{D} \subset \mathcal{E}$ we have $\mathcal{D} = \mathcal{E}$, i. e., \mathcal{D} is a maximal element in the set of all proper filters on F .

A pair (G, H) of subsets of F is called a *binary partition of F* if $G \cap H = \emptyset$ and $G \cup H = F$. A filter \mathcal{D} is a ultrafilter iff it has the *binary partition property*, i. e., if for every binary partition (G, H) of F either $G \in \mathcal{D}$ or $H \in \mathcal{D}$ (see [2, Exercise 2.119]).

Further on, we assume that \mathcal{D} is a filter.

Now let $((U_f, \gamma_f) \mid f \in F)$ be a collection of evaluated mathematical systems of the second-order signature Σ_2^g with true generalized equalities and belongings.

Define an evaluation γ on the system $U \equiv \text{infra-}\mathcal{D}\text{-prod}(U_f \mid f \in F)$ in the following way.

Let x be a variable of a type τ . If τ is the first-order type, then define $\gamma(x) \in \tau(A)$ setting $\gamma(x)(f) \equiv \gamma_f(x)$ for every $f \in F$. If $\tau = [\tau_0, \dots, \tau_k]$ is a second-order type, then put $\gamma(x) \equiv \{p \in \check{\tau}(A) \mid \forall f \in F (p(f) \in \gamma_f(x))\}$.

The evaluation γ will be called the *crossing of the collection of evaluations* $\{\gamma_f \mid f \in F\}$ and will be denoted by $\bowtie (\gamma_f \mid f \in F)$.

Lemma 1. *Let $\{(U_f, \gamma_f) \mid f \in F\}$ be a collection of evaluated mathematical systems of the second-order signature Σ_2^g and let every system (U_f, γ_f) be a model for equality axioms E1–E4. Then the pair $(\text{infra-}\mathcal{D}\text{-prod}(U_f \mid f \in F), \bowtie (\gamma_f \mid f \in F))$ is also a model for axioms E1–E4.*

Proof. Let $t_0, t'_0 \in \tau_0(A), \dots, t_k, t'_k \in \tau_k(A), P, P' \subset \check{\tau}(A) = \tau_0(A) \times \dots \times \tau_k(A), p \equiv (t_0, \dots, t_k), p' \equiv (t'_0, \dots, t'_k), p \approx_{\check{\tau}} p'$, and $P \approx_{\tau} P'$.

Assume that $p \in_{\tau} P$. According to the definition of the belonging, we get $\exists G_1 \in \mathcal{D} \forall g \in G_1 (p(g) \in_{\tau, g} P\langle g \rangle)$. By the definition of the first-order equality, $\exists G_2 \in \mathcal{D} \forall g \in G_2 (p(g) \approx_{\tau, g} p'(g))$. Finally, by the definition of the second-order equalities $\exists G_3 \in \mathcal{D} \forall g \in G_3 (P\langle g \rangle \approx_{\tau, g} P'\langle g \rangle)$. Since every system (U_g, γ_g) satisfies E4, we see that $p'(g) \in_{\tau, g} P'\langle g \rangle$ for every $g \in G \equiv G_1 \cap G_2 \cap G_3$. Thus, $p' \in_{\tau} P'$. Hence, $p \in_{\tau} P \Rightarrow p' \in_{\tau} P'$. The inverse implication is checked quite similarly. This proves axiom E4. The validity of axioms E1, E2, E3 is obvious. \square

Further, for a formula $\varphi \in L(\Sigma)$ the set $\{f \in F \mid U_f \models \varphi[\gamma_f]\}$ will be denoted by G_{φ} .

Lemma 2. *Let $\tau = [\tau_0, \dots, \tau_k]$ be a second-order type. Let s_{ω}^{τ} be the constants constructed above for the support $A \equiv \prod (A_f \mid f \in F)$. Then $s_{\omega}^{\tau}\langle f \rangle = s_{\omega_f}^{\tau}$ for every $f \in F$.*

Proof. Let $\xi \in s_{\omega}^{\tau}\langle f \rangle$, i. e., $\xi = p(f)$ for some $p \in s_{\omega}^{\tau}$. By definition, $\xi = p(f) \in s_{\omega_f}^{\tau}$. Consequently, $s_{\omega}^{\tau}\langle f \rangle \subset s_{\omega_f}^{\tau}$.

Conversely, let $\xi_f \in s_{\omega_f}^{\tau}$. Using the axiom of choice we can find a collection $(\xi_g \mid g \in F \setminus \{f\})$ such that $\xi_g \in s_{\omega_g}^{\tau}$. Define the element $p \in \check{\tau}(A)$ setting $p(\mu)(g) \equiv \xi_g(\mu)$ for every $g \in F$ and every $\mu \in k+1$. Then $p(g) = \xi_g \in s_{\omega_g}^{\tau}$ for every $g \in F$ implies $p \in s_{\omega}^{\tau}$. Since $\xi_f = p(f)$, we have $\xi_f \in s_{\omega}^{\tau}\langle f \rangle$. Hence, $s_{\omega_f}^{\tau} \subset s_{\omega}^{\tau}\langle f \rangle$. \square

Lemma 3. *Let $\tau = [\tau_0, \dots, \tau_k]$ be a second-order type. Let x be a variable of the type τ and $\gamma(x)$ be the evaluation constructed above for the system $U \equiv (A, S)$. Then $\gamma(x)\langle f \rangle = \gamma_f(x)$ for every $f \in F$.*

The proof is completely similar to the proof of the previous lemma.

5.2 Infrafiltration of formulas of the second-order language $L(\Sigma_2^g)$

Consider a non-empty set F and a filter \mathcal{D} on F .

By analogy with the first order language (see [13, § 17], [14, 8.2]) a formula φ of the language $L(\Sigma_2^g)$ of the second-order signature Σ_2^g with generalized equalities and belongings will be called *infiltrated with respect to the filter \mathcal{D}* if for every collection $\{(U_f, \gamma_f) \mid f \in F\}$ of evaluated mathematical systems of the second-order signature Σ_2^g with true generalized equalities and belongings the property $\text{infra-}\mathcal{D}\text{-prod}(U_f \mid f \in F) \models \varphi[\bowtie (\gamma_f \mid f \in F)]$ is equivalent to the property $\{g \in F \mid U_g \models \varphi[\gamma_g]\} \in \mathcal{D}$.

Lemma 1. *Every atomic formula is infiltrated with respect to any filter \mathcal{D} on the set F .*

Proof. First, consider an atomic formula φ of the form $q^{\tau} \delta_{\tau} r^{\tau}$. Assume that $q^{\tau} = x^{\tau}$ and $r^{\tau} = \sigma_{\omega}^{\tau}$. Then $U \models \varphi[\gamma]$ is equivalent to $\gamma(x) \approx_{\tau} s_{\omega}^{\tau}$, and analogously for the pair (U_f, γ_f) .

Let τ be the first-order type. Let $G_\varphi \in \mathcal{D}$, i. e., $\gamma_g(x) \approx_{\tau,g} s_{\omega_g}^\tau$ for every $g \in G_\varphi \in \mathcal{D}$. Then $\gamma_g(x) = \gamma(x)$ and $s_{\omega_g}^\tau = s_\omega^\tau(g)$ implies $\gamma(x)(g) \approx_{\tau,g} s_\omega^\tau(g)$ for every $g \in G_\varphi \in \mathcal{D}$. Thus, $\gamma(x) \approx_\tau s_\omega^\tau$, i. e., $U \models \varphi[\gamma]$.

Conversely, let $U \models \varphi[\gamma]$, i. e., $\gamma(x) \approx_\tau s_\omega^\tau$. Then there exists $G \in \mathcal{D}$ such that $\gamma(x)(g) \approx_{\tau,g} s_\omega^\tau(g)$ for every $g \in G$. But it means that $\gamma_g(x) \approx_{\tau,g} s_{\omega_g}^\tau$, i. e., $U_g \models \varphi[\gamma_g]$ for every $g \in G \in \mathcal{D}$. Since $G \subset G_\varphi$, we have $G_\varphi \in \mathcal{D}$.

Now let $\tau \equiv [\tau_0, \dots, \tau_k]$ be a second-order type. Let $G_\varphi \in \mathcal{D}$, i. e., $\gamma_g(x) \approx_{\tau,g} s_{\omega_g}^\tau$ for every $g \in G_\varphi \in \mathcal{D}$. According to Lemmas 2 and 3, the equalities $s_{\omega_g}^\tau = s_\omega^\tau\langle g \rangle$ and $\gamma_g(x) = \gamma(x)\langle g \rangle$ are correct. Therefore $\gamma(x)\langle g \rangle \approx_{\tau,g} s_\omega^\tau\langle g \rangle$ for every $g \in G_\varphi \in \mathcal{D}$. Consequently, $\gamma(x) \approx_\tau s_\omega^\tau$, i. e., $U \models \varphi[\gamma]$.

Conversely, let $U \models \varphi[\gamma]$, i. e., $\gamma(x) \approx_\tau s_\omega^\tau$. By the definition of the second-order equality, $\gamma(x)\langle g \rangle \approx_{\tau,g} s_\omega^\tau\langle g \rangle$ for some $G \in \mathcal{D}$ and every $g \in G$. Using Lemmas 2 and 3 we obtain $\gamma_g(x) \approx_{\tau,g} s_{\omega_g}^\tau$, i. e., $U_g \models \varphi[\gamma_g]$ for every $g \in G \in \mathcal{D}$. Since $G \subset G_\varphi$, we infer that $G_\varphi \in \mathcal{D}$.

For the terms q^τ and r^τ of other forms the reasons are quite similar.

Now consider an atomic formula φ of the form $(q_0^{\tau_0}, \dots, q_k^{\tau_k}) \varepsilon_\tau r^\tau$ for $\tau \equiv [\tau_0, \dots, \tau_k] \in \Theta_b$. Assume that $q_\lambda^{\tau_\lambda} = x_\lambda^{\tau_\lambda}$ and $r^\tau = u^\tau$ for some variables x_λ and u . Then $U \models \varphi[\gamma]$ is equivalent to $(\gamma(x_0), \dots, \gamma(x_k)) \varepsilon_\tau \gamma(u)$ and analogously for the pair (U_f, γ_f) .

Let $G_\varphi \in \mathcal{D}$, i. e., $(\gamma_g(x_0), \dots, \gamma_g(x_k)) \varepsilon_{\tau,g} \gamma_g(u)$ for every $g \in G_\varphi$. Consider the elements $\xi_f \equiv (\gamma_f(x_0), \dots, \gamma_f(x_k))$ and $p \equiv (\gamma(x_0), \dots, \gamma(x_k)) \in \tilde{\tau}(A)$. Let $f \in F$. Then $p(f)(\mu) \equiv p(\mu)(f) = \gamma(x_\mu)(f) = \gamma_f(x_\mu) = \xi_f(\mu)$ for every $\mu \in k+1$. Consequently, $p(f) = \xi_f$. By Lemma 2 $\gamma_f(u) = \gamma(u)\langle f \rangle$. As a result, we obtain $p(g) \varepsilon_{\tau,g} \gamma(x)\langle g \rangle$ for every $g \in G_\varphi \in \mathcal{D}$. By definition, it means that $p \varepsilon_\tau \gamma(x)$, i. e., $U \models \varphi[\gamma]$.

Conversely, let $U \models \varphi[\gamma]$, i. e., $(\gamma(x_0), \dots, \gamma(x_k)) \varepsilon_\tau \gamma(u)$. By the definition of the second-order belonging, for $p \equiv (\gamma(x_0), \dots, \gamma(x_k))$ there exists $G \in \mathcal{D}$ such that $p(g) \varepsilon_{\tau,g} \gamma(u)\langle g \rangle$ for every $g \in G$. By Lemma 3 $\gamma(u)\langle g \rangle = \gamma_g(u)$. By the previous subsection, $\xi_g = p(g)$. Consequently, $\xi_g \varepsilon_{\tau,g} \gamma_g(u)$, i. e., $U_g \models \varphi[\gamma_g]$ for every $g \in G$. Since $G \subset G_\varphi$, we infer that $G_\varphi \in \mathcal{D}$.

For the terms $q_\lambda^{\tau_\lambda}$ and r^τ of other forms the reasons are quite similar. \square

A proof of the property of infrafiltration for the quantified formula $\exists x^\tau \varphi$ for the language $L(\Sigma_2^g)$ of the generalized second-order signature Σ_2^g is more delicate than for the first-order language. Therefore we begin it with a subsidiary proposition.

Let $\{(U_f, \gamma_f) \mid f \in F\}$ be a collection of evaluated mathematical systems of the second-order signature Σ_2^g with true generalized equalities and belongings. Let β be an evaluation on the system $U \equiv \text{infra-}\mathcal{D}\text{-prod}(U_f \mid f \in F)$.

For the evaluation β and for every $f \in F$ define the evaluation δ_f on the system U_f in the following way. Let x be a variable of a type τ . If τ is the first-order type, then put $\delta_f(x) \equiv \beta(x)(f)$. If τ is a second-order type, then put $\delta_f(x) \equiv \beta(x)\langle f \rangle$. Consider the evaluation $\delta \equiv \bowtie(\delta_f \mid f \in F)$.

Propositon 1. *The equalities $\delta(x^\tau) \approx_\tau \beta(x^\tau)$ hold for any variable x^τ .*

Proof. If τ is the first-order type, then by the definition of the evaluations δ and δ_f we obtain $\delta(x)(f) \equiv \delta_f(x) = \beta(x)(f)$ for any $f \in F$, i. e., $\delta(x) = \beta(x)$.

Let τ be a second-order type. Lemma 3 implies $\delta(x)\langle f \rangle = \delta_f(x) = \beta(x)\langle f \rangle$ for any $f \in F$. By the definition of the second-order equality, we conclude that $\delta(x) \approx_\tau \beta(x)$. \square

Propositon 2. *Let a formula ψ be infrafiltrated with respect to the filter \mathcal{D} . Then the formula $\exists x^\tau \psi$ is infrafiltrated with respect to \mathcal{D} as well.*

Proof. Denote the formula $\exists x^\tau \psi$ by φ . Let $G_\varphi \in \mathcal{D}$, i. e., $U_g \models \varphi[\gamma_g]$ for every $g \in G_\varphi \in \mathcal{D}$. Further, we shall write simply G instead of G_φ .

The presented satisfaction property means that $U_g \models \psi[\gamma'_g]$ for some evaluation γ'_g such that $\gamma'_g(y) = \gamma_g(y)$ for every $y^\sigma \neq x^\tau$. For every $f \in F$ define the evaluation δ_f setting $\delta_f \equiv \gamma_f$ if $f \in F \setminus G$ and $\delta_f \equiv \gamma'_f$ if $f \in G$.

Check that the evaluated systems (U_f, δ_f) and (U_g, δ_g) are H -concordant for every $f, g \in F$. If $f, g \in F \setminus G$, then $\delta_f = \gamma_f$ and $\delta_g = \gamma_g$. Since the evaluations γ_f and γ_g are H -concordant, our assertion is true. Let $f, g \in G$. Then $\delta_f = \gamma'_f$ and $\delta_g = \gamma'_g$. Let x be a variable of a type τ .

Consider the evaluation $\delta \equiv \bowtie \{\delta_f \mid f \in F\}$. Check that $\delta(y) = \gamma(y)$ for every $y^\sigma \neq x^\tau$.

Let σ be the first-order type. Then $\delta(y)(g) = \delta_g(y) = \gamma'_g(y) = \gamma_g(y) = \gamma(y)(g)$ for $g \in G$. If $f \in F \setminus G$, then $\delta(y)(f) = \delta_f(y) = \gamma_f(y) = \gamma(y)(f)$. Consequently, $\delta(y) = \gamma(y)$.

Let σ be a second-order type. If $f \in G$, then $\delta_f(y) = \gamma'_f(y) = \gamma_f(y)$. If $f \in F \setminus G$, then $\delta_f(y) = \gamma_f(y)$. Let $p \in \delta(y)$. By the definition of the crossing, $p(f) \in \delta_f(y)$ for every $f \in F$. By the above, $p(f) \in \gamma_f(y)$ for every $f \in F$. This means that $p \in \gamma(y)$, whence $\delta(y) \subset \gamma(y)$. The inverse inclusion is checked in the same way. Consequently, $\delta(y) = \gamma(y)$.

Thus, for every $y \neq x$ we have $\delta(y) = \gamma(y)$.

By condition and construction, $U_g \models \psi[\delta_g]$ for every $g \in G \in \mathcal{D}$. Since the formula ψ is infrafiltrated, the obtained property implies the property $U \models \psi[\delta]$. Since $\delta(y^\sigma) = \gamma(y^\sigma)$ for every $y^\sigma \neq x^\tau$, we obtain the property $U \models \varphi[\gamma]$.

Conversely, let $U \models \varphi[\gamma]$. It is equivalent to $U \models \psi[\beta]$ for some evaluation β , H -concordant with the evaluation γ and such that $\beta(y) = \gamma(y)$ for every $y^\sigma \neq x^\tau$.

Consider the evaluation $\delta \equiv \bowtie \{\delta_f \mid f \in F\}$ from Proposition 1, corresponding to the evaluation β . According to Proposition 1, $\delta(z^\rho) \approx_\rho \beta(z^\rho)$ for every variable z^ρ . It follows from Proposition 2 that the property $U \models \psi[\beta]$ is equivalent to the property $U \models \psi[\delta]$. Since the formula ψ is infrafiltrated, the property $U \models \psi[\delta]$ is equivalent to the property $G \equiv \{g \in F \mid U_g \models \psi[\delta_g]\} \in \mathcal{D}$.

Let $y^\sigma \neq x^\tau$. If σ is the first-order type, then $\delta_g(y) = \beta(y)(g) = \gamma(y)(g) = \gamma_g(y)$. If σ is a second-order type, then $\delta_g(y) = \beta(y)\langle g \rangle = \gamma(y)\langle g \rangle$. Since by Lemma 3 $\gamma(y)\langle g \rangle = \gamma_g(y)$, we have $\delta_g(y) = \gamma_g(y)$. Consequently, in all the cases $\delta_g(y) = \gamma_g(y)$ for every $y^\sigma \neq x^\tau$. Therefore the property $U_g \models \psi[\delta_g]$ is equivalent to the property $U_g \models \varphi[\gamma_g]$. Thus, $\{g \in F \mid U_g \models \varphi[\gamma_g]\} = G \in \mathcal{D}$. This implies $G_\varphi \in \mathcal{D}$. \square

The following two lemmas are the same as ones for the first-order language.

Lemma 2. *Let formulas ψ and ξ be infrafiltrated with respect to the filter \mathcal{D} . Then the formula $\psi \wedge \xi$ is infrafiltrated with respect to \mathcal{D} as well.*

Proof. Denote the formula $\psi \wedge \xi$ by φ . Let $G_\varphi \in \mathcal{D}$, i.e., $U_g \models \varphi[\gamma_g]$ for all $g \in G_\varphi \in \mathcal{D}$. This property is equivalent to the conjunction of the properties $U_g \models \psi[\gamma_g]$ and $U_g \models \xi[\gamma_g]$. Since these formulas are infrafiltrated, it is equivalent to the conjunction of the properties $U \models \psi[\gamma]$ and $U \models \xi[\gamma]$, but it is equivalent to the property $U \models \varphi[\gamma]$.

Conversely, let $U \models \varphi[\gamma]$. It is equivalent to the conjunction of the properties $U \models \psi[\gamma]$ and $U \models \xi[\gamma]$. Then $G_\psi \in \mathcal{D}$ and $G_\xi \in \mathcal{D}$. Consider $G \equiv G_\psi \cap G_\xi$. Then $U_g \models \psi[\gamma_g]$ and $U_g \models \xi[\gamma_g]$ implies $U_g \models \varphi[\gamma_g]$ for every $g \in G \in \mathcal{D}$. Hence, $G_\varphi \in \mathcal{D}$. \square

Lemma 3. *Let a formula ψ be infrafiltrated with respect to the ultrafilter \mathcal{D} . Then the formula $\neg\psi$ is infrafiltrated with respect to \mathcal{D} as well.*

Proof. Denote the formula $\neg\psi$ by φ . By assumption, the properties $G_\psi \in \mathcal{D}$ and $U \models \psi[\gamma]$ are equivalent.

By definition, $F \setminus G_\varphi = \{g \in F \mid \text{the property } U_g \models \varphi[\gamma_g] \text{ does not hold}\}$. But $U_g \models \varphi[\gamma_g]$ is equivalent to the assertion that the property $U_g \models \psi[\gamma_g]$ does not hold. Consequently the property $U_g \models \psi[\gamma_g]$ is equivalent to the assertion that the property $U_g \models \varphi[\gamma_g]$ does not hold. It implies $F \setminus G_\varphi = G_\psi$.

Let $G_\varphi \in \mathcal{D}$. Since \mathcal{D} is an ultrafilter, we have $G_\psi = F \setminus G_\varphi \notin \mathcal{D}$. So the property $U \models \psi[\gamma]$ does not hold. By the definition of the satisfiability, it means that $U \models \varphi[\gamma]$.

Conversely, let $U \models \varphi[\gamma]$. Then the property $U \models \psi[\gamma]$ does not hold. Therefore $G_\psi \notin \mathcal{D}$. Since \mathcal{D} is an ultrafilter, we have $G_\varphi = F \setminus G_\psi \in \mathcal{D}$. \square

Theorem 1 (the generalized infrafiltration theorem). *Every formula φ of the language $L(\Sigma_2^g)$ of the second-order signature Σ_2^g with generalized equalities and belongings is infrafiltrated with respect to any ultrafilter \mathcal{D} on the set F .*

Proof. The set of all formulas φ of the language $L(\Sigma_2^g)$, constructed by induction from atomic formulas by means of the connectives \neg and \wedge and the quantifier \exists , will be denoted by Ψ . The subset of the set Ψ , consisting of all formulas containing at most n logical symbols \neg , \wedge , and \exists , will be denoted by Ψ_n . Obviously, $\Psi = \bigcup\{\Psi_n \mid n \in \omega_0\}$.

Using the complete induction principle we shall prove the following assertion $A(n)$: *every formula $\varphi \in \Psi_n$ is infrafiltrated.*

If $n = 0$, then φ is an atomic formula. By Lemma 1, it is infrafiltrated. Consequently, $A(0)$ holds.

Assume that for every $m < n$ the assertion $A(m)$ holds. Let $\varphi \in \Psi_n$. If $\varphi = \neg\psi$, then $\psi \in \Psi_{n-1}$. Therefore, ψ is infrafiltrated. By Lemma 3, the formula φ is infrafiltrated as well. If $\varphi = \psi \wedge \xi$, then $\psi, \xi \in \Psi_{n-1}$. Therefore, by the inductive assumption, the formulas ψ and ξ are infrafiltrated. By Lemma 2, the formula φ is infrafiltrated as well. Finally, if $\varphi = \exists x^\tau \psi$, then $\psi \in \Psi_{n-1}$. Consequently, as above, the formula ψ is infrafiltrated. By Proposition 2 the formula φ is infrafiltrated as well. Thus, the assertion $A(n)$ holds.

By the complete induction principle the assertion $A(n)$ holds for every $n \in \omega_0$. This means that any formula $\varphi \in \Psi$ is infrafiltrated.

Let φ be an arbitrary formula of the language $L(\Sigma_2^g)$. Consider for φ the accompanying formula φ^* defined in 3.3. By the definition of the operation $\varphi \mapsto \varphi^*$, we have $\varphi^* \in \Psi$. By the proven above, the formula φ^* is infrafiltrated, i. e., $\{g \in F \mid U_g \models \varphi^*[\gamma_g]\} \in \mathcal{D} \Leftrightarrow U \models \varphi^*[\gamma]$. Proposition 1 implies the equivalences $U \models \varphi^*[\gamma] \Leftrightarrow U \models \varphi[\gamma]$ and $U_g \models \varphi^*[\gamma_g] \Leftrightarrow U_g \models \varphi[\gamma_g]$. As a result we get the following chain of equivalences:

$$\{g \in F \mid U_g \models \varphi[\gamma_g]\} \in \mathcal{D} \Leftrightarrow \{g \in F \mid U_g \models \varphi^*[\gamma_g]\} \in \mathcal{D} \Leftrightarrow U \models \varphi^*[\gamma] \Leftrightarrow U \models \varphi[\gamma].$$

It means that the formula φ is infrafiltrated. □

This theorem has one important corollary. Let Φ be some set of formulas of the language $L(\Sigma_2^g)$ of the generalized second-order signature Σ_2^g . Let the set Φ has a model (U_0, γ_0) of the signature Σ_2^g with true generalized equalities and belongings. Take an arbitrary set F and an arbitrary ultrafilter \mathcal{D} on F . Consider the collection of the models $\{(U_f, \gamma_f) \mid f \in F\}$ such that $(U_f, \gamma_f) \equiv (U_0, \gamma_0)$. The infra- \mathcal{D} -product $\text{infra-}\mathcal{D}\text{-prod}(U_f \mid f \in F)$ of the collection $(U_f \mid f \in F)$ will be called the *infra- \mathcal{D} -power of the system U_0 with the exponent F* and will be denoted by $\text{infra-}\mathcal{D}\text{-power}(U_0, F)$. The crossing $\bowtie \{\gamma_f \mid f \in F\}$ of the collection $(\gamma_f \mid f \in F)$ will be called the *crossing of the evaluation γ_0 in the quantity F* and will be denoted by $\bowtie \{\gamma_0, F\}$.

Corollary 1. *Let Φ be some set of formulas of the language $L(\Sigma_2^g)$. If the set Φ has a model (U_0, γ_0) of the signature Σ_2^g with true generalized equalities and belongings, then for every set F and every ultrafilter \mathcal{D} on F the set Φ has also the model $(\text{infra-}\mathcal{D}\text{-power}(U_0, F), \bowtie \{\gamma_0, F\})$ of the signature Σ_2^g with true generalized equalities and belongings.*

5.3 Compactness theorem for formulas of the language $L(\Sigma_2^g)$

In the capacity of some pleasant complementary corollary to the infrafiltration theorem we deduce the generalized compactness theorem for the language $L(\Sigma_2^g)$. It is well-known that it does not hold for the standard language $L(\Sigma_2^{st})$ [2, Appendix].

Theorem 1. *Let Φ and Ψ be some sets of formulas of the language $L(\Sigma_2^g)$ of the generalized second-order signature Σ_2^g . Let for every finite subset f of the set Φ the set of formulas $f + (E1-E4) + \Psi$ has a model (U_f, γ_f) of the signature Σ_2^g . Then the set of formulas $\Phi + (E1-E4) + \Psi$ has a model (U, γ) of the signature Σ_2^g .*

Proof. Consider the set $F \equiv \{f \subset \Phi \mid 0 < |f| < \omega\}$ of all finite non-empty subsets from Φ .

For an element $f \in F$ consider the set $F_f \equiv \{g \in F \mid f \subset g\}$. Since $f \in F_f$, we have $F_f \neq \emptyset$. The ensemble $\mathfrak{C} \equiv \{F_f \mid f \in F\}$ has the finite intersection property, i. e., it is multiplicative. Hence, there is some ultrafilter \mathcal{D} on the set F including the set \mathfrak{C} .

Consider the system $U \equiv \text{infra-}\mathcal{D}\text{-prod}(U_f \mid f \in F)$ and the evaluation $\gamma \equiv \bowtie(\gamma_f \mid f \in F)$ on the system U constructed in 5.1. By Lemma 1, U is a system with the true generalized equalities and belongings.

Prove that the evaluated system (U, γ) is a model for the set Φ .

Suppose $\varphi \in \Phi$. Consider the set $F_{\{\varphi\}}$. By condition, $U_{\{\varphi\}} \models \varphi[\gamma_{\{\varphi\}}]$. Consider the set $G_\varphi \equiv \{g \in F \mid U_g \models \varphi[\gamma_g]\}$. If $g \in F_{\{\varphi\}}$, then $\{\varphi\} \subset g$ implies $\varphi \in g$. Therefore $U_g \models \varphi[\gamma_g]$. Consequently, $F_{\{\varphi\}} \subset G_\varphi$. Since $F_{\{\varphi\}} \in \mathcal{D}$, we have $G_\varphi \in \mathcal{D}$.

By Theorem 1 we infer the property $U \models \varphi[\gamma]$. Thus, (U, γ) is a model for the set Φ . The fact that (U, γ) is a model for the set Ψ follows immediately from Theorem 1. \square

6 Inductive sequence of models of non-canonical generalized second-order Dedekind real axes with exponentially increasing powers

6.1 The formulation of Final theorem

Final theorem.

- (I) *Let F be a fixed set. Then there exist some sequence $(\widehat{\mathbb{R}}_i \mid i \in \omega_0)$ of sets $\widehat{\mathbb{R}}_i$, some sequence $(S_i \mid i \in \omega_0)$ of superstructures S_i of the signature Σ_{R2}^g over the sets $\widehat{\mathbb{R}}_i$, and some sequence $(u_i \mid i \in \omega_0)$ of mappings $u_i : \widehat{\mathbb{R}}_i \rightarrow \widehat{\mathbb{R}}_{i+1}$ such that:*
- (1) $R_0 \equiv (\widehat{\mathbb{R}}_0, S_0) \equiv (\mathbb{R}, S_{R2})$;
 - (2) *every system $R_i \equiv (\widehat{\mathbb{R}}_i, S_i)$ of the signature Σ_{R2}^g is a model for the theory Th_{R2}^g ;*
 - (3) *every mapping u_i is an $(\approx_{\pi, i}, \approx_{\pi, i+1})$ -injective homomorphism of the signature Σ_{R2}^g from the system R_i into the system R_{i+1} ;*
 - (4) *the image of the system R_i in the system R_{i+1} respectively to the homomorphism u_i is a submodel of the model R_{i+1} ;*
 - (5) *the support $\widehat{\mathbb{R}}_{i+1}$ of the system R_{i+1} is the set $\widehat{\mathbb{R}}_i^F$;*
 - (6) $(u_i p)(f) = p$ for every $p \in \widehat{\mathbb{R}}_i$ and every $f \in F$, i. e., $u_i p$ is the $\{p\}$ -valued function on F .
- (II) *There exists some superstructure S_{ω_0} of the signature Σ_{R2}^g over the set $\widehat{\mathbb{R}}_{\omega_0} \equiv \prod(\widehat{\mathbb{R}}_i \mid i \in \omega_0)$ and some sequence of mappings $w_i : \widehat{\mathbb{R}}_i \rightarrow \widehat{\mathbb{R}}_{\omega_0}$ such that:*
- (1) *the system $R_{\omega_0} \equiv (\widehat{\mathbb{R}}_{\omega_0}, S_{\omega_0})$ of the signature Σ_{R2}^g is a model for the theory Th_{R2}^g ;*
 - (2) *every mapping w_i is an $(\approx_{\pi, i}, \approx_{\pi, \omega_0})$ -injective homomorphism of the signature Σ_{R2}^g from the system R_i into the system R_{ω_0} ;*
 - (3) *the image of the system R_i in the system R_{ω_0} respectively to the homomorphism w_i is a submodel of the model R_{ω_0} ;*
 - (4) $w_i = w_{i+1} \circ u_i$ for every $i \in \omega_0$.

6.2 Detailed superstructures in Final theorem

Here we give the detailed description of the superstructures S_i from Final theorem in the same manner as it is given for the superstructure S_{R2} in 4.3.

The superstructure S_i is the quadruple $(S_{c,i}, S_{e,i}, S_{b,i}, S_{v,i})$, where:

- the *collection of constant structures* $S_{c,i}$ is the suit

$$((s_0^{\pi,i}, s_1^{\pi,i}), \emptyset, (s_0^{\rho,i}, s_1^{\rho,i}, s_2^{\rho,i}), (s_0^{\lambda,i}, s_1^{\lambda,i})) = (0_i, 1_i), \emptyset, (-_i, /_i, \leq_i), (+_i, \cdot_i));$$

- the *collection of the equality ratios* $S_{e,i}$ is the suit $(\approx_{\pi,i}, \approx_{\varkappa,i}, \approx_{\rho,i}, \approx_{\lambda,i})$;
- the *collection of the belonging ratios* $S_{b,i}$ is the suit $(\in_{\varkappa,i}, \in_{\rho,i}, \in_{\lambda,i})$;
- the *collection of the terminals* $S_{v,i}$ over the set $\widehat{\mathbb{R}}_i$ is the suit

$$(\pi(\widehat{\mathbb{R}}_i), \varkappa(\widehat{\mathbb{R}}_i), \rho(\widehat{\mathbb{R}}_i), \lambda(\widehat{\mathbb{R}}_i)) = (\widehat{\mathbb{R}}_i, \mathcal{P}(\widehat{\mathbb{R}}_i), \mathcal{P}(\widehat{\mathbb{R}}_i^2), \mathcal{P}(\widehat{\mathbb{R}}_i^3)).$$

6.3 The proof of Final theorem

(I) The construction of the infra- \mathcal{D} -power of the system U_0 with the exponent F from 5.2 gives the opportunity to prove part I of the Final theorem.

Fix some ultrafilter \mathcal{D} on F . We shall construct the necessary sequence of models by natural induction. Take for the initial model $R_0 \equiv (\mathbb{R}_0, S_0)$ the canonical model $R_2^g \equiv (\mathbb{R}, S_{R2})$ from 4.3. Assume that the model $R_i \equiv (\widehat{\mathbb{R}}_i, S_i)$ with some evaluation ζ_i is constructed.

Take the system $R_{i+1} \equiv (\widehat{\mathbb{R}}_{i+1}, S_{i+1}) \equiv \text{infra-}\mathcal{D}\text{-power}(R_i, F)$ and the evaluation $\zeta_{i+1} \equiv \bowtie(\zeta_i, F)$ defined in 5.1. According to Corollary to Theorem 1 the evaluated system (R_{i+1}, ζ_{i+1}) is a model for the theory Th_{R2}^g . And the support $\widehat{\mathbb{R}}_{i+1}$ of this model is the set $\widehat{\mathbb{R}}_{i+1}^F \equiv \prod(\widehat{\mathbb{R}}_{i_f} \mid f \in F)$, where $\widehat{\mathbb{R}}_{i_f} \equiv \widehat{\mathbb{R}}_i$ for every $f \in F$. Since the set Ψ_2^g of axioms of the theory Th_{R2}^g from 4.2 consists of closed formulas only, the system R_{i+1} is a model for this theory.

Define the mapping $u_i : \widehat{\mathbb{R}}_i \rightarrow \widehat{\mathbb{R}}_{i+1}$ setting $(u_i(p))(f) \equiv p$ for every $p \in \widehat{\mathbb{R}}_i$ and every $f \in F$. Check that u_i is $(\approx_{\pi,i}, \approx_{\pi,i+1})$ -injective. Take some $p, q \in \widehat{\mathbb{R}}_i$ and suppose that $u_i(p) \approx_{\pi,i+1} u_i(q)$. By the construction from 5.1 there exists $G \in \mathcal{D}$ such that $(u_i(p))(g) \approx_{\pi,g,i} (u_i(q))(g)$ for every $g \in G$. Since $G \neq \emptyset$ we can take $g_0 \in G$. Then $(u_i(p))(g_0) \equiv p$ and $(u_i(q))(g_0) \equiv q$ implies $p \approx_{\pi,i} q$.

The construction of constant structures presented in 5.1 implies immediately that u_i is a homomorphism of the signature Σ_{R2}^g from the system R_i into the system R_{i+1} .

(II) The construction of the infra- \mathcal{D} -product of the collection of mathematical systems from 5.1 gives the opportunity to prove part II of the Final theorem. Fix some ultrafilter \mathcal{E} on ω_0 . Take the system $R_{\omega_0} \equiv (\widehat{\mathbb{R}}_{\omega_0}, S_{\omega_0}) \equiv \text{infra-}\mathcal{E}\text{-prod}(R_i \mid i \in \omega_0)$ and the evaluation $\zeta_{\omega_0} \equiv \bowtie(\zeta_i \mid i \in \omega_0)$ defined in 5.1. According to part I and Theorem 1 the evaluated system $(R_{\omega_0}, \zeta_{\omega_0})$ is a model for the theory Th_{R2} . Since the set Ψ_2^g of axioms of this theory from 4.2 consists of closed formulas only, the system R_{ω_0} is a model for this theory.

Fix $i \in \omega_0$. Construct some mapping $w_i : \widehat{\mathbb{R}}_i \rightarrow \widehat{\mathbb{R}}_{\omega_0}$ by the inverse and direct natural inductions. For the base of direct induction put $(w_i p)(i) \equiv p$ and $(w_i p)(i+1) \equiv u_i p$. For the step of direct induction put $(w_i p)(j+1) \equiv u_j((w_i p)(j))$ for $j \geq i+1$. Fix some $f_0 \in F$. Put $(w_i p)(i-1) \equiv p(f_0)$ for the base of inverse induction. For the step of inverse induction put $(w_i p)(j-1) \equiv ((w_i p)(j))(f_0)$ for $1 \leq j \leq i-1$. These constructions can be described in a more rigorous form based on [12, Theorem 1(1.2.8)].

By the natural induction in ST it can be proved that w_i is a homomorphism of the signature Σ_{R2}^g from the system R_i into the system R_{ω_0} (see the example of scrupulous arguments below).

Check that w_i is $(\approx_{\pi,i}, \approx_{\pi,\omega_0})$ -injective. Take some $p, q \in \widehat{\mathbb{R}}_i$ and suppose that $w_i(p) \approx_{\pi,\omega_0} w_i(q)$. By the construction from 5.1 there exists $J \in \mathcal{E}$ such that $(w_i p)(j) \approx_{\pi,j} (w_i q)(j)$ for every

$j \in J$. Consider the binary partition $(i, \omega_0 \setminus i)$ of ω_0 . Since \mathcal{E} is a ultrafilter, we infer that either $i \in \mathcal{E}$ or $\omega_0 \setminus i \in \mathcal{E}$. If $i \in \mathcal{E}$, then $\omega_0 \in \mathcal{E}$ but it is not so. Hence, $\omega_0 \setminus i \in \mathcal{E}$. This implies $J \cap (\omega_0 \setminus i) \in \mathcal{E}$, and, therefore, there is $j \in J$ such that $j \geq i$. Take $k_0 \equiv j - i$.

If $j = i$, then by the definition of w_i we have $(w_i p)(i) = p$ and $(w_i q)(i) = q$. Hence, $p \approx_{\pi, i} q$. If $j = i + 1$, then $(w_i p)(i + 1) \equiv u_i p$ and $(w_i q)(i + 1) \equiv u_i q$ imply $(u_i p)(i + 1) \approx_{\pi, i+1} (u_i q)(i + 1)$. Since by assertion 3 of part I the mapping u_i is $(\approx_{\pi, i}, \approx_{\pi, i+1})$ -injective, we infer that $p \approx_{\pi, i} q$. Consider in ST the set $K \equiv \{k \in \mathbb{N} \mid ((w_i p)(i + k) \approx_{\pi, i+k} (w_i q)(i + k)) \Rightarrow p \approx_{\pi, i} q\}$. Let Φ_{ST} be a totality of axioms of the theory ST, i. e., Φ_{ST} consists of all explicit proper axioms of this theory, all implicit proper axioms of this theory, and all implicit logical axioms of the predicate calculus (see, e. g., [12, 1.1.3–1.1.11 and A.1.2]). Denote the first formula in the definition of K by $\varphi(i + k)$ and the second one by ψ . We have proved in ST the existence of deduction $\Phi_{ST}, \varphi(i + 1) \vdash \psi$. Since ST is the first-order theory, we conclude that $\Phi_{ST} \vdash \varphi(i + 1) \Rightarrow \psi$ by virtue of the deduction theorem (see, for example, [2, Proposition 2.5] or [12, 1.1.3]). Hence, $1 \in K$.

Suppose that $k \in K$ and $(w_i p)(i + k + 1) \approx_{\pi, i+k+1} (w_i q)(i + k + 1)$. By the definition of w_i we have $(w_i p)(i + k + 1) \equiv u_{i+k}((w_i p)(i + k))$ and the same for q . Since by assertion 3 of part I the mapping u_{i+k} is $(\approx_{\pi, i+k}, \approx_{\pi, i+k+1})$ -injective, we infer that $(w_i p)(i + k) \approx_{\pi, i+k} (w_i q)(i + k)$. Now from $k \in K$ we deduce that $p \approx_{\pi, i} q$. Thus, we have proved the existence of deduction $\Phi_{ST}, \varphi(i + k + 1) \vdash \psi$. As above this implies $\Phi_{ST} \vdash \varphi(i + k + 1) \Rightarrow \psi$, and, therefore, $k + 1 \in K$. By the principle of natural induction in ST (see [12, 1.2.6]) we get $K = \mathbb{N}$.

This means that for our $j = i + k_0$ we have $k_0 \in K$. Since $j \in J$, we conclude that $p \approx_{\pi, i} q$. This proves assertion 2.

Now we must only prove assertion 4. Fix $p \in \widehat{\mathbb{R}}_i$. Then $(u_i p)(f) \approx p$ for every $f \in F$. By the definition we have $(w_{i+1}(u_i p))(i + 1) \approx u_i p \approx (w_i p)(i + 1)$. For the base of direct induction we have

$$(w_{i+1}(u_i p))(i + 2) \approx u_{i+1}((w_{i+1}(u_i p))(i + 1)) \approx u_{i+1}((w_i p)(i + 1)) \approx (w_i p)(i + 2).$$

For the base of inverse induction we have

$$(w_{i+1}(u_i p))(i) \approx (u_i p)(f_0) \approx p \approx (w_i p)(i).$$

Then by the direct and inverse inductions we check that $(w_{i+1}(u_i p))(j) \approx_{\pi, j} (w_i p)(j)$ for every $j \in \omega_0$. Hence, $(w_{i+1} \circ u_i)(p) \approx_{\pi, \omega_0} w_i(p)$ for every $p \in \widehat{\mathbb{R}}_i$. \square

Remark 1. Since every set $\widehat{\mathbb{R}}_i \equiv \widehat{\mathbb{R}}_{i-1}^F$ for $i \geq 1$ consists of “real”-valued functions $p : F \rightarrow \widehat{\mathbb{R}}_{i-1}$, it is necessary to clarify directly the satisfaction of non-evident axioms A3 (the existence and functionality of the inversion) and A19 (the linearity of the order) on the systems R_i .

In case of A3 take any function $p \in \widehat{\mathbb{R}}_i$ such that $p \not\approx_{\pi, i} 0_i$, where 0_i denotes the null in $\widehat{\mathbb{R}}_i$. Consider the binary partition of F consisting of the sets $\text{zer}(p) \equiv \{f \in F \mid p(f) \approx_{\pi, f, i-1} 0_{i-1}\}$ and $\text{coz}(p) \equiv F \setminus \text{zer}(p)$. Since \mathcal{D} has the binary partition property, we have either $\text{zer}(p) \in \mathcal{D}$ or $\text{coz}(p) \in \mathcal{D}$. In the first case we conclude that $p \approx_{\pi, i} 0_i$ but it contradicts our assumption. Hence, $\text{coz}(p) \in \mathcal{D}$ and $p(g) \not\approx_{\pi, i-1} 0_{i-1}$ for every $g \in \text{coz}(p)$. By A3 for every $g \in \text{coz}(p)$ there exists $p(g)^{-1}$ such that $(p(g), p(g)^{-1}) \in_{\rho, i-1} /_{i-1}$. Define p^{-1} setting $p^{-1}(g) \equiv p(g)^{-1}$ for every $g \in \text{coz}(p)$ and $p^{-1}(f) \equiv p(f)$ for every $f \in \text{zer}(p)$. By the definition of ρ -belonging $\in_{\rho, i}$ from 5.1 $(p, p^{-1}) \in_{\rho, i} /_i$. Thus, we deduced the existence of the inversion in R_i from the existence of the inversion in R_{i-1} using the binary partition property of the ultrafilter \mathcal{D} .

In case of A19 take any functions $p, q \in \widehat{\mathbb{R}}_i$. Since $\widehat{\mathbb{R}}_{i-1}$ is linearly ordered with respect to the order \leq_{i-1} , we can take the binary partition of F consisting of the sets

$$\begin{aligned} G &\equiv \{g \in F \mid (p(g), q(g)) \in_{\rho, i-1} \leq_{i-1}\} \text{ and} \\ H' &\equiv \{h \in F \mid ((q(h), p(h)) \in_{\rho, i-1} \leq_{i-1}) \wedge (q(h) \not\approx_{\pi, i-1} p(h))\}. \end{aligned}$$

By binary partition property of \mathcal{D} we have either $G \in \mathcal{D}$ or $H' \in \mathcal{D}$. In the first case we conclude that $(p, q) \in_{\rho, i} \leq_i$. In the second case we can see that $H' \subset H \equiv \{h \in F \mid (q(h), p(h)) \in_{\rho, i-1} \leq_{i-1}\} \in \mathcal{D}$

implies $(q, p) \in_{\rho, i \leq i}$. Thus, we deduced the linearity of the order in R_i from the linearity of the order in R_{i-1} using again the binary partition property of the ultrafilter \mathcal{D} .

Open question 1. Part II of Final theorem shows that the model R_{ω_0} can be considered as some pretender for the *inductive limit of the inductive sequence* $s \equiv (R_i \mid i \in \omega_0)$ in the sense of [20, 11.8]. But this is an open question.

Open question 2 (about transfinite extension of the inductive sequence s). Let λ be an ordinal number such that $\lambda > \omega_0$ and \mathcal{E} be an ultrafilter on λ . Since \mathcal{E} has the binary partition property, we can consider the ultrafilters $\mathcal{E}_\alpha \equiv \{E \subset \alpha \mid \exists G \in \mathcal{E} (E = G \cap \alpha)\}$ for every ordinal number $\alpha \in \lambda \setminus \omega_0 = [\omega_0, \lambda[$. Starting from R_{ω_0} we can construct by the transfinite procedure some collection $t \equiv (R_\alpha \mid \alpha \in \lambda \setminus \omega_0)$ of models for the theory $Th_{R_2}^q$ such that: 1) $R_\alpha \equiv \text{infra-}\mathcal{E}_\alpha\text{-prod}(R_\gamma \mid \gamma \in \alpha \setminus \omega_0)$ for limit ordinal number α and 2) $R_{\alpha+1} \equiv \text{infra-}\mathcal{E}_\alpha\text{-power}(R_\alpha, F)$. *Is the collection t inductive with respect to some injective homomorphisms $u_{\alpha\beta} : \hat{\mathbb{R}}_\alpha \rightarrow \hat{\mathbb{R}}_\beta$ for every $\alpha < \beta$ and does t extend s?*

Supplement 1. In [12, C.3.4] the *generalized second-order Peano – Landau theory $Th_{N_2}^q$ of natural numbers* is considered. It is clear that some inductive sequence $(N_i \mid i \in \omega_0)$ of models of this theory can be constructed, which is similar to the inductive sequence $s \equiv (R_i \mid i \in \omega_0)$ constructed above. And also the inductive “quasilimit” N_{ω_0} of this sequence can be constructed similarly to the inductive “quasilimit” R_{ω_0} . Moreover, its own Final theorem can be proved for the generalized models N_i and N_{ω_0} of the theory $Th_{N_2}^q$. Besides, open questions 1 and 2 are valid for these hypothetical models.

References

- [1] Feferman S. The number systems. Foundations of algebra and analysis. Addison-Wesley Publishing, Reading, 1963.
- [2] Mendelson E. Introduction to Mathematical Logic, 4th ed. Chapman&Hall / CRC, London, 1997.
- [3] Takeuti G. Proof theory. Dover Publications, Mineola, NY, 2013.
- [4] Dalen D.v. Logic and Structure. Springer, Berlin, 1997.
- [5] Rossberg M. First-order logic, second-order logic, and completeness, in: First-order logic revisited. Proceedings of the conference FOL75 – 75 years of first-order logic. Logos Verlag, Berlin, 2004, pp. 303–321.
- [6] Shapiro S. Foundations without Foundationalism: A Case for Second-Order Logic. Oxford University Press, Oxford, 1991.
- [7] Väänänen J. Second-order logic and foundations of mathematics. Bull. Symb. Log. 2001. 7(4). 504–520. [doi:10.2307/2687796](https://doi.org/10.2307/2687796).
- [8] Zakharov V.K. Compactness theorem for generalized second-order language, in: Contemporary problems of fundamental and applied mathematics. MPhTI, Dolgoprudnii, 2008, pp. 11–31 (in Russian).
- [9] Zakharov V.K., Yashin A.D. Compactness theorem for some generalized second-order language. J. Math. Research 2014. 6(3). 21–38. [doi:10.5539/jmr.v6n3p21](https://doi.org/10.5539/jmr.v6n3p21).
- [10] Zakharov V.K., Rodionov T.V. Some inductive sequence of models of generalized second-order Dedekind theory of real numbers with exponentially increasing powers, in: Proc. of Lobachevsky Math. Center, 57. “Theory of functions, its appl., and adj. probl.”. Kazan Math. Soc., Kazan, 2019, pp.155–159.

-
- [11] Zakharov V.K. Local set theory. *Math. Notes*. 2005 77(2). 177–193. doi:[10.1007/s11006-005-0019-x](https://doi.org/10.1007/s11006-005-0019-x).
- [12] Zakharov V.K., Rodionov T.V. *Sets, Functions, Measures. Volume I: Fundamentals of Set and Number Theory*. De Gruyter Studies in Math. 68/1, de Gruyter, Berlin, 2018. doi:[10.1515/9783110550948](https://doi.org/10.1515/9783110550948).
- [13] Ershov Yu.L., Palyutin E.A. *Mathematical logic*. Mir, Moscow, 1984.
- [14] Maltsev A.I. *Algebraic systems*. Springer, Berlin, 1973.
- [15] J. R. Shoenfield, *Mathematical logic*, Assoc. of Symb. Logic, Natick, MS, 2001.
- [16] Bourbaki N. *Eléments de Mathématique. Livre III. Topologie générale. Chapitre 3. Groupes topologiques (théorie élémentaire). Chapitre 4. Nombres réels*, 3rd Edition. Actualities sci. Ind., 1143. Hermann, Paris, 1960.
- [17] Landau E. *Elementary number theory*. Chelsea, New York, 1967.
- [18] Hewitt E., Stromberg K. *Real and abstract analysis. A modern treatment of the theory of functions of a real variable*. Springer, New York, 1965.
- [19] Grauert H., Lieb I. *Differential- und Integralrechnung. I: Funktionen einer reellen Veränderlichen*. Springer, Berlin–Heidelberg, 1967.
- [20] Semadeni Z. *Banach spaces of continuous functions*. Monografie Matematyczne, 55. PWA, Warszawa, 1971.