

**PERFORMANCE OF PRE RELEASED RICE (*ORYZA SATIVA L.*) GENOTYPES UNDER DIFFERENT SOWING WINDOWS IN RABI SEASON.**

**ABSTRACT**

The aim of this study was to investigate the effect of different sowing dates on growth and yield potential of pre released rice genotypes under irrigated conditions of Northern Telangana zone. The field experiments were carried out during two consecutive rabi seasons of 2018-19 and 2019-20, on clay soils of agricultural research station, Kunaram, Telangana state, India. The experiment was laid out in strip plot design with three replications. The treatments comprised of three sowing dates i.e. 20<sup>th</sup> November, 5<sup>th</sup> December and 20<sup>th</sup> December in horizontal factor and four genotypes i.e. KNM 733, RNR 15048, KNM 1638 and KNM 118 in vertical factor. Pooled data analysis results revealed that the different sowing dates and varieties significant effect on all the studied growth and yield characters. The rice crop sown on 20<sup>th</sup> December recorded significantly higher grain yield ( 8138 kg ha<sup>-1</sup>) and Among the genotypes, the short slender, short duration genotype KNM 733 recorded the recorded the maximum grain yield ( 8024 kg ha<sup>-1</sup>), which was on par with the other genotypes . The treatment combinations data results concluded that the, among the genotypes the genotype KNM 118 was recorded highest grain yield (8438 kg ha<sup>-1</sup>) when sowing was taken up on 20<sup>th</sup> December and followed by the genotype KNM 733 with sown on 20<sup>th</sup> November. In respect of economics of treatment combinations, the highest net returns (Rs.91,165 ha<sup>-1</sup>) and B:C (2.47) ratio were obtained when rice crop was sown during 20<sup>th</sup> December with the genotype KNM 118 and followed by sown on 20<sup>th</sup> November with the genotype KNM 733.

*Key words:* Dates, Economics, Genotypes, Grain yield, KNM, Sown,

**1. INTRODUCTION**

Rice (*Oryza sativa* L.) is one of the most important staple food grain crops of the world, which constitute the principle food for 60 per cent of the world's population and 2/3rd of Indian population. Rice is intensively grown in 88 countries across the world on an area about 160.43million hectares with annual production of 495.73 million tones [1]. In India rice is grown in 43.79 million hectares, the production level is 116.40 million tones and the productivity is about 2659 kg ha<sup>-1</sup> (Agricultural statistics at a glance-2018) [2]. In Telangana rice is grown in 1.95 million hectares with annual production of 4.46 million tones and the productivity is about 3436 kg ha<sup>-1</sup> (Agricultural statistics at a glance-2018) More than 80 per cent of the world's rice is produced and consumed in Asia [3], where it is an integral part of culture and tradition. Rice, it is believed, is associated with wet, humid climate, though it is not a tropical plant. It is probably a descendent of wild grass that was most likely cultivated in the foothills of the far Eastern Himalayas. The optimum period of time for sowing and transplanting of rice is critical in achieving high grain yield. However, optimum rice planting dates vary with regional, location and genotypes [4].

Rice plants require a particular temperature for its phenological affairs such as panicle initiation, flowering, panicle exertions from flag leaf sheath and maturity. Performance of a genotype entirely depends upon the time of planting. Rice needed before or after the window of optimum dates usually has slow germination and emergence, poor crop stand establishment, increased soil borne, seedling diseases damage under cold conditions, and seeds lose by birds or mice. Seedling at the optimum time is an important factor of transplanting for uniform stand establishment of rice. On the other hand, seedling sown with the delay of sowing more than optimum produces fewer tillers due to the reduction of the vegetative period and hence results in poor yield. Among the crop production tools, optimum time and method of sowing are the important agronomic tools that allow the crop to complete its growth timely and successfully under specific agro-ecology zone [5].

The significantly highest effective tillers /hill, leaf area index (LAI) at flowering , filled grain / panicle, 1000 grain weight coupled with significantly lower sterility percentage were observed with 15 July planting during the rainy season [6].

Varieties play a unique role in maximizing yield by improving the input- use efficiency as the genetic potential of variety limits the expression of its yield and affects plant growth in response to environment condition. The genotype KNM 733 is a short slender, short duration, non-shattering and good quality traits

with high yield potential and the genotype KNM 1638 is an early duration, medium slender and non-lodging culture with high yield potential.

Pre released KNM rice cultures developed from breeding experiments are to be evaluated for their performance with best checks under different dates of sowings before recommending to the farmers. There is a need to find out the optimum dates of sowings during rabi for getting higher yield in Northern Telangana Zone.

## 2. MATERIALS AND METHODS

A field experiment was conducted for two consecutive rabi seasons of 2018-19 and 2019-20 at Agriculture Research Station, Kunaram situated at an altitude of 231 m above mean sea level at 18.5272° N latitude and 79.4943° E longitude. The soil of the experiment sites was clay in texture, saline in reaction. It was normal electrical conductivity (0.40 dS m<sup>-1</sup>) and just below neutral in reaction (pH 6.36). The organic carbon content was low (0.26-0.40%) while medium in available Nitrogen and phosphorus respectively but high in potash content. The experiment was laid out in strip plot design replicated thrice. The treatments combination comprised three sowing dates in the horizontal factor viz. 20<sup>th</sup> November, 5<sup>th</sup> December and 20<sup>th</sup> December, and four genotypes in the vertical factor viz. KNM 733, RNR 15048, KNM 1638 and KNM 118. The net size of each plot was 27 m<sup>2</sup> (6.0 × 4.5 m). Row to row and plant to plant distance was made at 15 cm apart and seedlings were transplanted according to different dates of sowings. Seeds were sown at the rate of 50 kg ha<sup>-1</sup> in each date of sowing in the nursery. The seeds were treated with bavistin @ 2g kg<sup>-1</sup> seed before sowing. One third of the recommended dose of nitrogen (150 kg ha<sup>-1</sup>), full dose of phosphorus (60 kg ha<sup>-1</sup>) and half dose of potash (40 kg ha<sup>-1</sup>) were applied at after main field preparation, and the remaining nitrogen was top-dressed in two equal splits dose, at active tillering (18 -20 DAT), and at panicle initiation stage. The remaining half of the potassium applied at panicle initiation stage in the D<sub>1</sub>, D<sub>2</sub> and D<sub>3</sub> transplanted plots. For effective weed management, oxadiazyl (70 grams a.i.ha<sup>-1</sup>) was used in moist condition at morning sunshine hours in all the treatments just after D<sub>1</sub>, D<sub>2</sub> and D<sub>3</sub> transplanted fields. Zinc sulphate (2 grams liter<sup>-1</sup>) was sprayed to foliage at 25 and 30 DAT to avoid zinc deficiency in the crop. Irrigation was applied @ 5 cm at 7 to 8 days interval to maintain soil moisture at field capacity from sowing to one week before harvest during dry spells in the season. The plant height, tillers production and dry matter accumulation were recorded at tiller initiation,

maximum tillering, panicle initiation, 50 % flowering and maturity stage of the crop growth. The yield attributes and grain yield was recorded at harvest and sun dried straw yield was recorded 15 days after harvest.

### **3. RESULTS AND DISCUSSION**

#### **Growth parameters:**

##### **3.1. Plant height (cm):**

The crop sown on 20<sup>th</sup> December recorded significantly maximum plant height (103.0 cm) and followed by 5<sup>th</sup> December and 20<sup>th</sup> November. Plant height is directly proportional to the length of the vegetative phase of the crop. Increased plant height in late planting dates was due to response of different varieties to different photoperiodic conditions. Among the genotypes, the genotype RNR 15048 registered the highest plant height (107.0 cm) and which was significantly difference with the other genotypes. The interaction effect of dates of sowing and varieties on plant height was significant. The difference in plant height among the genotypes due to variation in the duration, photo-sensitiveness, thermo-sensitiveness and vegetative lag period of the variety. These results are in line with Safdar et al. [7].

##### **3.2. Number of tillers per square meter:**

At harvest stage, crop sown on 20<sup>th</sup> December recorded significantly more number of effective tillers m<sup>-2</sup> (457.0) over the crop sown on 5<sup>th</sup> December; however, this was statistically at par to crop sown on 20<sup>th</sup> November. This may be due to availability of favorable soil and air temperature during growing cycle of the crop. Similarly these findings were supported by Sharma [8] and Osman et al. [9]. Among the short duration genotypes viz., KNM 118 and KNM 733 registered higher number of effective tillers m<sup>-2</sup> (447 and 438) and were on par with each other and significantly superior over RNR 15048 and KNM 1638 varieties. It might be due to more relative growth rate and biomass production resulted in intra plant competition among the tillers for growth resources lead to more mortality of tillers when compared to short and extra short duration cultivars [10]. The results are in close conformity with Birhane [11], who also found highly significant difference in number of effective tiller plant<sup>-1</sup> due to variety, method of planting and their interaction.

##### **3.3. Total dry matter per square meter:**

There was no significant difference in dry matter accumulation noticed from 20<sup>th</sup> November to 20<sup>th</sup> December and a significant reduction was observed only in the crop sown early on 20<sup>th</sup> November. Among the genotypes, the genotype of KNM 1638 accumulated more dry matter m<sup>-2</sup> (1730 gm<sup>-2</sup>) and significantly superior over rest of the varieties respectively. The difference in dry matter accumulation among the genotypes might be due to their genetic potential and differential plant height. Similar results were obtained by Dileep et al. [12].

#### **3.4. Phenology:**

The days taken to reach flowering and harvest varied significantly among the sowing dates (Table 1). The significantly higher number of days was taken by 20<sup>th</sup> November sown crop, while the significantly lowest number of days was taken by 20<sup>th</sup> December sown crop. Sowing date primarily influences the length of vegetative period of rice with early sown rice requiring a greater number of days to accumulate the same number of degree days units compared with later sown rice. A linear negative correlation between sowing dates and growth period, in the later sowing dates was also reported by Peng-fei et al. [13]. Wani et al. [14], reported the days taken to reach flowering and harvest varied significantly among the sowing dates.

#### **Yield parameters & Yield**

#### **3.5. Yield attributes:**

The crop sown on 20<sup>th</sup> December produced maximum number of panicles m<sup>-2</sup> (479) and it is on par with the crop sown on 20<sup>th</sup> November and significantly difference with the 5<sup>th</sup> December date sown crop. Among the short duration genotypes, the genotype KNM 733 registered higher number of panicles m<sup>-2</sup> (450) and there was no significant difference was observed among the genotypes. Similarly these findings were supported by Jagtap et al [15]. The crop sown on 20<sup>th</sup> November produced more number of filled grains panicles<sup>-1</sup> (240) and which was significantly difference with the 5<sup>th</sup> December and 20<sup>th</sup> December sown crop (Table 2). Late sowing shortened the growth period of the plant which reduced the number of grains panicle<sup>-1</sup> than early sown crop under aerobic culture [16]. Among the genotypes the RNR 15048 was recorded significantly the highest number of filled grains panicle<sup>-1</sup> (278) and significantly superior to other genotypes. The number of spikelets panicle<sup>-1</sup> is basically genetic feature of a variety.

These results are in close conformity with Balaji naik et al. [17], who also observed significantly differences in number of filled spikelets panicle<sup>-1</sup> due to varieties and sowing times in aerobic rice system.

### **3.6. Test weight (1000 grain weight):**

The crop sown on 5<sup>th</sup> December recorded more 1000 grain weight (17.05 g) and was significantly superior to 20<sup>th</sup> November and 20<sup>th</sup> December sown crop. Higher test weight obtained from the crop sown in 5<sup>th</sup> December might be attributed optimum photoperiod available for crop growth and development. These results are in line with the findings of Muhammad et al. [16] and Jagtap et al. [15]. Among the genotypes, the genotype KNM 118 recorded significantly the highest 1000 grain weight (25.3g) over the other genotypes KNM 733, KNM 1638 and RNR 15048. These results clearly indicated that, the 1000 grain weight is a varietal feature which might be affected least with the environmental conditions.

### **3.7. Grain yield (Kg ha<sup>-1</sup>):**

Significantly more grain yield (8138 kg ha<sup>-1</sup>) was realized from the crop sown on 20<sup>th</sup> December and which is on par with the crop sown on 5<sup>th</sup> December and was significantly difference with the grain yield of crop sown on 20<sup>th</sup> November. The higher grain yield in 20<sup>th</sup> December was due to higher effective tillers and spikelets panicle<sup>-1</sup> compared to rest of the dates (Table 2). Chendge et al. [18], reported that the increased yield might be due to result of optimum growth and development parameters and yield contributes associated with favorable weather condition responsible for more growth and development resulted in more storage of photosynthates in the grain. The grain yield showed no significant difference among the different genotypes. Among the four genotypes tested the genotype KNM 733 produced more grain yield (8024 kg ha<sup>-1</sup>) and followed by other genotypes KNM 1638 (7912 kg ha<sup>-1</sup>) and check genotypes RNR 15048 (7884 kg ha<sup>-1</sup>) and KNM 118 (7791 kg ha<sup>-1</sup>). The genotype KNM 733 was performed better due to more conversion of photo synthetics into economic produce, which resulted in higher yield attributing characters in the respective varieties. Similar findings have been reported by Hussain et al. [19], Suresh et al. [20] and Walia et al. [21]. So, it would be better to choose short duration genotype KNM 733 and follow optimum sowing dates from 5<sup>th</sup> December to 20<sup>th</sup> December under irrigated conditions of Northern Telangana Zone.

### **Interaction effect**

The interaction effect among the different sowing dates and genotypes for grain yield found to be significant in the pooled mean of the data (Table 3). The short duration long slender genotype KNM 118 sown during 20<sup>th</sup> December recorded higher grain yield (8438 kg ha<sup>-1</sup>) and followed by crop sown on 20<sup>th</sup> November With the genotype KNM 733 (8301 kg ha<sup>-1</sup>). Similar findings were reported by Manjunath et al. [22] and Reddy et al. [23].

### **3.8. Harvest index (%):**

The harvest index was significantly influenced by different dates of sowing and genotypes. The crop sown on 20<sup>th</sup> December registered higher harvest index (60.0) and which was significant difference with other dates of sowing. The harvest index showed no significant difference among the genotypes. Among the genotype RNR 15048 registered highest harvest index value (59.67) and followed by other genotypes. Similar findings was observed by Hossain et al. [24] and Salahuddin et al. [25], concluded that high harvest index was mainly due to higher grain yield which was achieved through better performance in most of the yield, attributing traits.

### **3.9. Economic analysis:**

On the basis of economic analysis (Table 4), among the treatment combinations tested, it is observed that the highest net returns (Rs.91,165 ha<sup>-1</sup>) and B:C (2.47) ratio were obtained when rice crop was sown during 20<sup>th</sup> December with the genotype KNM 118 and followed by crop sown on 20<sup>th</sup> November with the genotype KNM 733.

## **4. CONCLUSION:**

In Northern Telangana zone, rice genotypes be sown in rabi during 20<sup>th</sup> December for obtaining higher grain and straw yield with higher net returns and B:C ratio. Similarly the rice genotype KNM 733 be grown for obtaining higher grain and straw yield, thus It would be concluded that rabi rice in Northern Telangana Zone be sown during 20<sup>th</sup> December with genotype KNM118 or KNM 733 sown during 20<sup>th</sup> November, so as to obtained higher yield and economic returns.

**Table. 1. Growth and yield attributes of pre released rice cultures as influenced by different dates of sowing during rabi season (Pooled data of 2 years).**

Treatments	Plant height (cm)	No. of tillers m <sup>-2</sup>	Dry matter production (g m <sup>-2</sup> )	Maximum tillering	No. of days to		Maturity
					Panicle Initiation	50% Flowering	
<b>Main plot: Date of sowing</b>							
20 <sup>th</sup> November	92.0	454.0	1497	65	93	105	134
5 <sup>th</sup> December	95.0	397.0	1586	59	89	101	129
20 <sup>th</sup> December	103.0	457.0	1687	60	93	100	127
SEm ±	0.28	4.17	60.0	0.16	0.53	0.18	0.26
CD (P = 0.05)	1.10	16.39	NS	0.62	2.06	0.70	1.02
<b>Sub-plot: Pre Released Cultures</b>							
KNM 733	96.11	438.17	1588	58	89	100	127
RNR 15048	107.30	432.11	1585	63	92	103	132
KNM 1638	88.87	425.83	1730	66	95	107	136
KNM118	94.17	447.39	1458	58	89	99	125
SEm ±	0.42	3.62	21.6	0.23	0.45	0.16	0.32
CD (P = 0.05)	1.45	12.54	75.0	0.78	1.55	0.57	1.11
<b>Interaction ( D X V )</b>							
SEm ±	SEm ±	0.38	8.16	81.79	0.39	1.0	0.44
CD (P = 0.05)	CD (P = 0.05)	1.16	NS	252.0	1.20	NS	1.35
<b>Interaction ( V X D )</b>							
SEm ±	0.38	7.30	82.45	0.33	0.90	0.37	0.53
CD (P = 0.05)	1.33	NS	286.0	1.15	NS	1.29	1.85

**Table. 2. Yield attributes of pre released rice cultures as influenced by different dates of sowing during rabi season (Pooled data of 2 years).**



Treatments	No. of panicles m <sup>-2</sup>	Panicle length (cm)	No. of filled grains panicle <sup>-1</sup>	Grain yield (kg ha <sup>-1</sup> )	Straw yield (kg ha <sup>-1</sup> )	Test weight (g)	Harvest index (%)	Net returns	B:C Ratio
<b>Main plot: Date of sowing</b>									
20 <sup>th</sup> November	459	22.54	240.16	7550	5617	16.77	57.67	75,680	2.22
5 <sup>th</sup> December	406	23.37	227.50	7993	5778	17.05	58.49	83,059	2.34
20 <sup>th</sup> December	479	23.44	200.33	8138	5472	16.64	60.15	85,597	2.38
SEm ±	4.85	0.10	2.76	1.02	32.01	0.05	0.31	1,412.0	0.02
CD (P = 0.05)	19.02	0.40	10.85	305.45	125.69	0.18	1.22	5544.0	0.09
<b>Sub-plot: Pre Released Cultures</b>									
KNM 733	450	23.51	224.41	8024.34	5765	15.11	58.40	83,642	2.35
RNR 15048	449	23.65	278.44	7884.93	5469	12.38	59.67	81,112	2.31
KNM 1638	443	22.65	254.10	7912.40	5666	14.43	58.49	81,610	2.32
KNM118	449	22.64	133.70	7791.62	5590	25.35	58.53	79,418	2.28
SEm ±	5.01	0.07	2.84	112.0	107.3	0.05	0.66	2,034.0	0.03
CD (P = 0.05)	NS	0.26	9.81	NS	NS	0.18	NS	NS	NS
<b>Interaction ( D X V )</b>									
SEm ±	6.74	0.18	6.34	162.0	137.6	0.09	1.02	2,941.0	0.05
CD (P = 0.05)	20.77	0.57	NS	499	424.0	0.27	NS	9061.0	0.15
<b>Interaction ( V X D )</b>									
SEm ±	6.75	0.17	5.46	142	109.7	0.08	0.83	2590.0	0.04
CD (P = 0.05)	23.38	0.58	NS	494	380.3	0.27	NS	8972.0	0.14

**Table.3. Interaction effect of dates of sowing & varieties on grain yield (kg ha<sup>-1</sup>) of rice cultures (Pooled data).**

Dates of sowing	Varieties				Mean
	KNM 733	RNR 15048	KNM 1638	KNM 118	
20 <sup>th</sup> November	8301	7727	7562	6752	7586
5 <sup>th</sup> December	7877	7872	8035	8183	7992
20 <sup>th</sup> December	7894	8055	8139	8438	8132
Mean	8024	7885	7912	7791	
SEm ±	162.0				
CD (P = 0.05)	499.0				

**Table. 4. Yield and economics of rice cultures as affected by different treatment combinations.**

Treatments	Grain yield (kg ha <sup>-1</sup> )			Gross Returns (Rs. ha <sup>-1</sup> )			Net Returns (Rs. ha <sup>-1</sup> )			B:C Ratio		
	2018	2019	Mean	2018	2019	Mean	2018	2019	Mean	2018	2019	Mean
D1V1	8910	7693	8301	1,61,714	1,39,619	1,50,667	99,714	77,619	88,667	2.61	2.25	2.43
D1V2	8365	7089	7727	1,51,832	1,28,662	1,40,247	89,832	66,662	78,247	2.45	2.08	2.26
D1V3	7767	7357	7562	1,40,965	1,33,536	1,37,250	78,965	71,536	75,250	2.27	2.15	2.21
D1V4	6968	6537	6752	1,26,467	1,18,646	1,22,557	64,467	56,646	60,557	2.04	1.91	1.98
D2V1	7609	8146	7877	1,38,097	1,47,854	1,42,975	76,097	85,854	80,975	2.23	2.38	2.31
D2V2	6874	8870	7872	1,24,764	1,60,996	1,42,880	62,764	98,996	80,880	2.01	2.60	2.30
D2V3	6827	9244	8036	1,23,913	1,67,785	1,45,849	61,913	1,05,785	83,849	2.00	2.71	2.35
D2V4	7880	8487	8184	1,43,026	1,54,038	1,48,532	81,026	92,038	86,532	2.31	2.48	2.40
D3V1	7752	8037	7894	1,40,696	1,45,871	1,43,283	78,696	83,871	81,283	2.27	2.35	2.31
D3V2	7685	8426	8056	1,39,486	1,52,929	1,46,207	77,486	90,929	84,207	2.25	2.47	2.36
D3V3	7157	9122	8139	1,29,896	1,65,567	1,47,731	67,896	1,03,567	85,731	2.10	2.67	2.38
D3V4	8633	8244	8439	1,56,695	1,49,635	1,53,165	94,695	87,635	91,165	2.53	2.41	2.47

## REFERENCES

1. Anonymous. United States Department of Agriculture, 2019.
2. Anonymous. Agricultural statistics at a glance, Directorate of Economics and Statistics, Government of India, 2018.

3. Bhat FM, Riar CS. Effect of amylase, particle size and morphology on the functionality of starches of traditional rice varieties. *International journal of biological macromolecules*. 2016;92:637-644.
4. Bashir MU, Akber A, Iqbal, Zaman H. Effect of different sowing dates on yield and yield components of direct seeded coarse rice (*Oryza sativa* L.) *Pakistan Journal of Agricultural Science*. 2010; 47(4):361-365.
5. Vange T, Obi IU. Effect of planting date on some agronomic traits and grain yield of upland rice varieties at Makurdi, Benue state, *Nigerian Journal of Sustainable Development of Agriculture Environment*. 2006; 2(1):1-9.
6. Annie P, Swain P, Rav KS. Agro-physiological parameter of rice (*Oryza sativa*) hybrids as affected by different data of planting under costal Orissa. *Indian Journal of Agricultural Science*. 2009;79(1): 25-28.
7. Safdar ME, Sarwar AASMG, Awan TH. Effect of transplanting dates on paddy yield of fine grain rice genotypes. *Pakistan Journal of Botany*. 2008;40(6): 2403-2411.
8. Sharma A., Dhaliwal LK, Sandhu SK, Singh S. Effect of plant spacing and transplanting time on phenology, tiller production and yield of rice (*Oryza sativa* L.). *International Journal of Agricultural Science*. 2011;7:249-253.
9. Osman KA, Mustafa AM, Elsheikh YMA, Idris A E. Influence of different sowing dates on growth and yield of direct seeded rice (*Oryza sativa* L.) in semi-arid zone. *International Journal of Agronomy and Agricultural Research*. 2015;6: 38-48.
10. Gill MS, Ashwani kumar, Pradeep kumar. Growth and yield of rice cultivars under various methods and times of sowing. *Indian Journal of Agronomy*.2006; 51(2): 123-127.
11. Birhane A. Effect of planting methods on yield and yield components of rice (*Oryza Sativa* L.) varieties in Tahtay Koraro Wereda, Northern Ethiopia. *International Journal of Technonolgy Enhancement and Emerging Engineeriong Research*. 2013;1(5): 2347-4289
12. Dileep K, Pasupalak S, Baliarsingh A. Effect of establishment methods and sowing time on growth and yield of rice varieties (*Oryza sativa* L.). *The Pharma Innivation Journal*. 2018;7(4): 904-907.

13. Peng-fei L, Zhi-hua Z, Yan-xia Z, Shao-hua X, Mingzhu J, Wen-jing H, Xu-hu Z. Effects of sowing date, basic seedling, nitrogen rate and field water on early rice "Liangyou9168". *Advance Journal of Food Science and Technology*. 2013;5(4): 414-417.
14. Wani SA, Qayoom S, Bhat MA, Lone BA, Nazir A. Influence of sowing dates and nitrogen levels on growth yield and quality of scented rice cv. Pusa Sugandh-3 in Kashmir valley. *Journal of Applied and Natural Science*. 2016;8(3): 1704-1709.
15. Jagtap DN, Chavan VG, Mahadkar UV, Chavan SA. Effect of sowing times on the yield and yield attributes of rice (*oryza sativa* L.) varieties. *Journal of Indian Society of Coastal Agricultural Research*. 2016;34(2): 27-30.
16. Bashir MU, Akbar N, Asifqbal, Haroonzaman. Effect of different sowing dates on yield and yield components of direct seeded coarse rice (*Oryza sativa* L.). *Pakistan Journal of Agricultural Science*. 2010; 47(4): 361-365.
17. Balaji naik B, Raji reddy D, Sreenivas G, Leela Rani P. Effect of sowing dates and varieties on growth, yield and economics of aerobic rice (*Oryza sativa* L.) during kharif season. *The journal of research PJTSAU*. 2015; 43(1&2): 18-24.
18. Chengde PD, Chavan SA, Ashwini patil, Shalu Kumar. Effect of sowing times on yield and economics of different rice genotypes under climatic condition of Konkan. *Journal of Pharmacognosy and Phytochemistry*. 2017; 6(5): 2462-2466.
19. Hussain A, Bhat MA, Ganai MA, Hussain T. Influence of planting dates and spacing schedules on performance of Basmati rice Pusa Sugandh 3 under Kashmir valley condition. *Environment and Ecology*. 2009; 27(1A):396-398.
20. Suresh K, Balaguravalah D, Ramalu V, Rao CSH. Evaluation of rice varieties under different management practices for late planting situation of Nagarjune Sagar left canal command area Andhra Pradesh, India. *International Journal of Plant Animal and environmental Science*. 2013; 3(2):258-260.
21. Walia US, Walia SS, Sindhu Nayyar S. Production of direct-seeded rice in relation to different dates of sowing and varieties in central Punjab. *Journal of crop weed*. 2014;10(1):126-129.

22. Manjunatha BN, Basavarajappa R, Pujari BT. Effect of age seedling on growth, yield and water requirement by different system of rice intensification. *Karnataka Journal of Agricultural Science*. 2010;23(2): 231-234.
23. Reddy SN, Narayana P. Pattern of dry matter accumulation and N uptake by rice as influenced by age of seedling and date of planting. *Andhra Agriculture Journal*. 1984;32:155-56.
24. Hossain MF, Salam M, Uddin M, Sarkar ZP. A comparative study of direct seeding versus transplanting method on the yield of *aus* rice. *Pakistan Journal of Agronomy*. 2002; 1(2-3): 86-88
25. Salahuddin KM, Chowhdury SH, Munira S, Islam MM, Parvin S. Response of nitrogen and plant spacing of transplanted aman rice. *Bangladesh Journal of Agricultural Research*. 2009; 34(2): 279-285.

UNDER PEER REVIEW