

DETERMINATION OF MERCHANTABILITY OF MAIZE (*Zea mays* L.) AS EPIS, SPATHES AND GRAINS STOCKED IN THE PRODUCTION AREAS OF CÔTE D'IVOIRE

ABSTRACT

Aims : This study aims to determine merchantability of maize produced and stocked as grains, epis and spathes of maize in five departments of Côte d'Ivoire.

Study Design : A total of 375 samples were collected at rate of 75 samples by department (Botro, Korhogo, Katiola, Agniblékro and Bondoukou). Then, the perceived samples was sent to the laboratory in order to analyse their merchantability.

Place and Duration of Study : The collection of samples was carried out on grains, epis and spathes maize from February 2013 to January 2014. Then, the analyzes were carried out at the Biochemistry and Food Sciences Laboratory of the Félix Houphouët-boigny University.

Methodology : The humidity was determined by drying until constant weight when the losses and damages of grains have been quantified by counting and weighing.

Results : The results show that for maize grains, humidity levels vary from $8.59\pm 0.11\%$ to $14.18\pm 0.52\%$. Concerning epis of maize, humidity levels are between $9.85\pm 0.23\%$ to $13.02\pm 0.32\%$. For maize spathes, humidity levels fluctuate between $11.16\pm 0.29\%$ to $14.17\pm 0.32\%$. The damages varies from $5.01\pm 2.64\%$ to $19.35\pm 2.00\%$ for maize grains, $9.05\pm 5.27\%$ to $21.29\pm 7.17\%$ for epis and $17.95\pm 1.13\%$ for spathes. As regards weigh losses, proportions of grains, epis and spathes are between respectively $1.53\pm 1.17\%$ and $5.10\pm 0.97\%$, $2.43\pm 1.70\%$ and $6.54\pm 1.79\%$ and $2.76\pm 1.18\%$ and $5.11\pm 2.65\%$.

Conclusion : A significant variability from one department to another can be noticed at level of maize quality regardless the type of maize. The merchantability of maize seems to be tied to post-harvest treatments (drying), type of storage (epis, grains and spathes) and structure of storage.

Keywords: *Merchantability, maize grains, maize epis, maize spathes, production area, Zea mays L., Côte d'Ivoire*

1. INTRODUCTION

Legumes and cereals are the mainstay of many African people's diet [1]. Among the cereals, maize holds a very important place for human and animal food. Native to Central America, this cereal is largely cultivated in many parts of the world for its starchy grains which represents an excellent source of energy. Thus, maize is the first cereal crop with the most significant production volume before soft wheat and rice [2]. Its world production was estimated at around 843 million tonnes in 2014 [3]. America contributes for about one half of that production with a volume estimated to 419 million tonnes, about 48% of production. As for Asia, Europe and Africa, maize provides respectively 28%, 7,5% and 4 % of world production. It guarantees almost half of the caloric intake in southern and eastern Africa and also one-fifth in western Africa [4].

In Côte d'Ivoire, maize holds the seventh rank of agricultural food-crop production and the second in cereal production after rice [5]. It is cultivated alone or in combination with other cultures in diverse agro-ecological area. The national production is estimated at around 760 000 tonnes in 2016 [5]. Maize represents the mainstay of many Ivorian people's diet. It is also used in animal food (poultry, pigs and bovines) and serves as raw material in some industries (brewery, soap and oil factory). Long regarded as a subsistence crop, maize today benefits from a strong support of agricultural research institutions [6-7]. The average yields of traditional varieties in rural area are of the order of 0.8 tonne per hectare versus 2 to 5 tonnes per hectare in controlled environment planted with varieties selected by research [8].

Despite its various uses, maize remains a seasonal crop in many area production. Therefore, its availability during offseason is systematically linked to conditions of its preservation. However, a number of constraints are regularly observed during its preservation. Most of them are linked to bad post-harvest treatments [1, 9-12]. These treatments include many activities such as drying and storage methods (attics, polypropylene and PICS bags). These activities are essential for a good preservation of the harvest quality [1, 12-13]. In fact, some studies revealed an increased insect activity when maize are insufficiently dried (a water content above 13%), stored in poor conditions (air humidity and high temperature) and in inappropriate structures. This contributes to creating a favorable environment for development of molds of genera *Aspergillus*, *Penicillium* and *Fusarium* responsible for production of mycotoxins [2, 9-11, 14-15]. These pests cause tremendous damages in stocks that can lead to qualitative and quantitative losses of grains [9-10, 11, 15]. In Côte d'Ivoire, many studies have been

made on conservation techniques of maize in areas production [9-10, 11, 15]. But few studies tackle the issue of the quality of maize produced and sold in areas productions. The main objective of this study is to determine the merchantability of maize coming from the areas production of Côte d'Ivoire.

2. MATERIAL AND METHODS

2.1. Material

2.1.1. Biological material

The biological material is composed of dry maize in the form of grains, epis and spathes deriving from the major areas production of this resource in Côte d'Ivoire.

2.1.2. Study sites

The samples were collected from the localities of Botro (Center, Gbèkè), Korhogo (North, Poro), Katiola (North Center, Hambol), Agnibilekro (Northeast, Indenié-Djuablin) and Bondoukou (East, Gountougo). Each of these departments has a geographical specificity and climatic characteristics which influence the seasons of maize production. Indeed, the localities of Botro (7°50'nord 5°18'west), Katiola (8°10'nord 5°40'west), Agnibilekro (7°02'nord 3°12'west) and Bondoukou (8°30'N 3°20'West) are characterized by a humid tropical climate (*Baouléen* climate). It has four seasons including two rainy seasons favoring maize production twice a year and two dry seasons [16]. Except the other four departments, the climate of department of Korhogo (9°27' nord 5°38' west) is of Sudanese type characterized by a rainy season favorable to maize production and a dry season [16-17].

2.2. Methods

2.2.1. Sampling

The samples were collected in 5 departments of production areas including Botro, Korhogo, Katiola, Agnibilekro and Bondoukou. This collection of samples was carried out on grains, epis and spathes maize from February 2013 to January 2014. A total of 375 samples were collected for each form of maize (125 grains, 125 epis, and 125 spathes, Table I). Then, the perceived samples was sent to the laboratory in order to analyse their merchantability.

Table I : Number of samples collected according to maize variety and department

Departments	Grains	Epis	Spathes	Total
Botro	25	25	25	75
Korhogo	25	25	25	75
Katiola	25	25	25	75
Agnibilékrou	25	25	25	75
Bondoukou	25	25	25	75
Total	125	125	125	375

2.2.2. Determination of humidity content

After shattering, 5 g (m_1) of maize broyat have been collected in an capsule of initial mass m_0 . The all capsule and sample was placed to dry at $105 \text{ }^\circ\text{C} \pm 2 \text{ }^\circ\text{C}$, until constant mass in an oven brand Memmert. Then, the all dried capsule and sample was weighed (m_2) and the humidity (H) was expressed as a percentage of dry mass on the basis of following formula [18] :

$$H (\%) = \frac{(m_1 - m_2) \times 100}{(m_1 - m_0)}$$

H : humidity of sample ; m_0 : capsule mass ; m_1 : the total mass before shattering (capsule + powder sample) ; m_2 : the total mass after shattering (capsule + powder sample).

Next, determined humidity levels were compared to Codex Alimentarius standard [19].

2.2.3. Evaluation of damages and weight losses

Based on counting and weighing method, the assessment of resulting damage by insects was carried out [20-21]. A 1 kg mass of grains maize containing approximately 3500 grains has been collected. After sieving and the withdrawal of foreign material, grains were weighed again and sorted in order to separate damaged grains (attacked and starved) from sound grains. Then the two fractions have been separately counted and weighed. Grains damages and weight losses are estimated on basis of below formulas :

$$D (\%) = (NGA / NTG) \times 100$$

D = Damages ; *NGA* = Number of attacked grains ; *NTG* = Total number of grains

$$W (\%) = \frac{[(NGA \times PGS) - (NGS \times PGA)]}{(PGS \times NTG)} \times 100$$

W= weight losses ; NGA = Number of attacked grains ; NGS = Number of sound grains ; NTG = Total number of grains ; PGA=Weight of attacked grains ; PGS = Weight of sound grains.

2.2.4. Statistical analysis

All the analyses were carried out in threefold test and data processed with software Statistical Program for Social Sciences (SPSS version 20.0, SPSS for Windows, USA). For each characteristic, the results were expressed in form of averages followed by their standard deviations as parameters of data spread. A two-way analysis of variance (ANOVA 2) was also made in order to test the impact of locality and the ways of preserving maize on assessed characteristics to 5% significant threshold statistical. For the statistically different averages, the Student-Newman-Keuls test served for the classification. Furthermore, the correlation between data and samples was estimated on basis of main components analysis (MCA), thanks to STATISTICA 7.1 software.

3. RESULTS AND DISCUSSION

3-1- Results

3.1.1. Humidity level

The obtained humidity levels statistically vary ($P=0.031$) depending on the departments and for all maize types (grains, epis and spathes) as shown in table II. For the maize grains, the humidity average levels are between $8.58 \pm 0.48\%$ and $14.18 \pm 2.63\%$ with a high proportion observed at Bondoukou and the lower proportions at Botro and Korhogo.

The humidity of samples collected at the level of maize epis is also lower at Botro ($9.85 \pm 1.19\%$) and Korhogo ($10.61 \pm 1.70\%$) as compared with Agnibilékrou ($13.20 \pm 1.63\%$), Bondoukou ($14.18 \pm 2.63\%$) and Agnibilékrou ($13.20 \pm 1.63\%$). With maize spathes, table II shows that humidity levels of samples at Botro, Korhogo and Bondoukou are statistically different (respectively $11.17 \pm 1.58\%$, $11.16 \pm 1.45\%$ and $11.53 \pm 1.49\%$), in comparison with Katiola ($12.43 \pm 1.53\%$) and Agnibilékrou ($14.17 \pm 1.61\%$).

Table II : Maize humidity levels sold in studied production areas

Départments	Grains	Epis	Spathes	F-value	P-value
Botro	8.92±0.59 ^{aA}	9.85±1.19 ^{aB}	11.17±1.58 ^{aC}	22.49	<0.001
Korhogo	8.58±0.48 ^{aA}	10.61±1.70 ^{aB}	11.16±1.45 ^{aB}	27.23	< 0.001
Katiola	9.94±0.99 ^{ba}	11.90±1.34 ^{aB}	12.43±1.53 ^{bB}	46.33	< 0.001
Agnibilékrou	13.20±1.63 ^{cA}	13.02±1.63 ^{ba}	14.17±1.61 ^{cB}	3.62	< 0.031
Bondoukou	14.18±2.63 ^{dA}	12.41±1.90 ^{bB}	11.53±1.49 ^{aC}	10.75	< 0.001
F-value	73.66	17.17	17.16	nd	nd
P-value	< 0.001	< 0.001	< 0.001	nd	nd

nd : not determined

By column and line the averages bearing the same letters are statistically identical. Lowercase letters are representative of columns and uppercase letters represent lines.

3.1.2. Proportion of samples that do not conform to commercial standard

On grains moisture, the maximum moisture limit to ensure good marketability is set at 15%. Thus, the departments of Botro, Korhogo and Katiola do not present any sample (0%) with a moisture content higher than 15%, whether the maize is in grains, epis or spathes. On the other hand, Agnibilékrou has 8% of maize grain, 8% of maize epi and 28% of maize spathe with water contents greater than 15%. Concerning Bondoukou, 40% and 8% of non-conformity of maize grain and maize epi (respectively) are also observed (Table III).

Table III : Samples proportion (%) of maize having a humidity level not in accordance at standard

Départments	Grains	Epis	Spathes
Botro	0	0	0
Korhogo	0	0	0
Katiola	0	0	4

Agnibilékrou	8	8	28
Bondoukou	40	8	0

3.1.3. Maize damages level

For each type of maize, the average level of damages in localities ($P < 0,05$) also differ. For grains the values fluctuate between $5.01 \pm 2.64\%$ and $19.35 \pm 2,00\%$ while epis and spathes respectively shows $9.05 \pm 5.27\%$ to $21.29 \pm 7.17\%$ and $10.06 \pm 3,06\%$ to $17.95 \pm 9.13\%$ of damages. The statistical analysis reveals lower percentages of damages of maize grains in the department of Botro ($5.81 \pm 3.08\%$), Korhogo ($5.01 \pm 2.64\%$) and Katiola ($8.81 \pm 1.49\%$) as compared with départements of Bondoukou ($13,41 \pm 7,83\%$) and specially Agnibilékrou ($19.35 \pm 2.00\%$). The grains collected from epis and spathes maize are also more damaged at Bondoukou ($19.20 \pm 1.50\%$ and $17.95 \pm 9.13\%$) and Agnibilékrou ($21.29 \pm 7.17\%$ and $13.20 \pm 1.13\%$) but in the other localities the averages vary from $9.05 \pm 5.27\%$ to $12.34 \pm 2.93\%$. Furthermore, among all the analysed samples, maize grains coming from Botro, Korhogo et Katiola is the least damaged (Table IV).

Table IV : Level of maize damages sold in production areas

Départements	Grains	Epis	Spathes	F-value	P-value
Botro	5.81 ± 3.08^{aA}	12.34 ± 2.93^{bC}	10.06 ± 3.06^{aB}	25.29	<0.001
Korhogo	5.01 ± 2.64^{aA}	9.05 ± 5.27^{aB}	10.86 ± 2.22^{aB}	17.09	< 0.001
Katiola	8.81 ± 1.49^{bA}	9.85 ± 1.44^{aB}	10.46 ± 1.39^{aC}	8.08	< 0.001
Agnibilékrou	19.35 ± 2.00^{dB}	21.29 ± 7.17^{cC}	13.20 ± 1.13^{bA}	58.93	< 0.001
Bondoukou	13.41 ± 7.83^{cA}	19.20 ± 1.50^{cA}	17.95 ± 9.13^{cA}	1.98	0.15
F-value	28.83	20.65	7.83	nd	nd
P-value	< 0.001	< 0.001	< 0.001	nd	nd

nd : not determined

By column and row, the averages covering the same letters are statistically identical. The lower case letters are representative of columns and capital letters are representative of rows.

3.1.4. Maize weight losses

The analysed level of maize weight losses are presented in table V. These losses are statistically invariable ($P>0.05$) for the three types of maize in the localities of Botro ($2.14\pm1.65\%$ to $3.02\pm1.55\%$), Katiola ($2.65\pm0.88\%$ to $3.02\pm1.99\%$) and Bondoukou ($4.22\pm2.17\%$ to $6.24\pm4.72\%$). However, as compared with epis and spathes of maize, the weight loss is significantly lower ($P<0.05$) at Korhogo and Agnibilékrou when maize is in grains form (respectively $1.53\pm1.17\%$ and $5.10\pm0.87\%$). Based on comparison of losses in localities, the results indicate fewer losses at level of maize grains and epis in the departments of Botro ($2.14\pm1.65\%$ to $3.02\pm1.55\%$) Korhogo ($1.53\pm1.17\%$ to $2.43\pm1.70\%$) and Katiola ($2.65\pm0.88\%$ to $2.87\pm1.50\%$) as compared with Agnibilékrou ($5.10\pm0.90\%$ to $6.54\pm1.79\%$) and Bondoukou ($4.22\pm2.17\%$ to $6.24\pm4.72\%$). Concerning the maize spathes, there is an important weight losses ($P<0.05$) at Bondoukou ($5.11\pm2.65\%$) in comparison with other localities ($2.76\pm1.18\%$ to $3.20\pm0.87\%$).

Table V : Level of maize weight losses sold in production areas

Départements	Grains	Epis	Spathes	F-value	P-value
Botro	2.14 ± 1.65^{abA}	3.02 ± 1.55^{aA}	2.94 ± 0.59^{aA}	2.48	0.09
Korhogo	1.53 ± 1.17^{aA}	2.43 ± 1.70^{aB}	2.76 ± 1.18^{aB}	5.32	0.006
Katiola	2.65 ± 0.88^{bA}	2.87 ± 1.50^{aA}	3.02 ± 1.99^{aA}	0.36	0.55
Agnibilékrou	5.10 ± 0.97^{cA}	6.54 ± 1.79^{cbB}	3.20 ± 0.87^{aC}	13.85	< 0.001
Bondoukou	4.22 ± 2.17^{cA}	6.24 ± 4.72^{bA}	5.11 ± 2.65^{bA}	1.20	0.31
F-value	13.50	9.68	5.09	nd	nd
P-value	< 0.001	< 0.001	0.001	nd	nd

nd : not determined

By column and row, the averages covering the same letters are statistically identical. The lower case letters are representative of columns and capital letters are representative of rows.


3.1.5. Variability of the characteristics of maize types

Table VI shows that analysis of F1 and F2 factors of main components reveal 99.17% of total variability of studied parameters. The F1 factor with a proper value of 2.63 expresses 87.76% of total variability while F2 factor of proper value of 0.34 reveals 11.41%. The projection of analysed variables in factorial design F1-F2 shows strong positive correlation between humidity, damages and weight


losses level of maize with F1 factor (Figure A). Based on the projection of samples in the same design, they are organised in two groups. Group 1 is composed of seven individuals presenting low humidity, weight losses and damages contents. It deals with spathes, epis and spathes maize coming from Korhogo and Botro but also grains maize from Katiola, Korhogo and Botro. Group 2 includes individuals having high humidity, weight losses and damages contents. It deals with spathes, epis and grains maize coming from Katiola, Bondoukou and Agnibilékrou.

Table VI: Proper values of parameters

Components	Component 1	Component 2	Component 3
Proper values	2.63	0.34	0.02
Expressed variability (%)	87.76	11.41	0.82
Accumulation of expressed (%)	87.76	99.17	100



A



B

Figure 1 : Projection of merchants parameters (A) and individuals(B) of grains, epis and spathes of maize in factorial plan 1-2 of the analysis of main components.

BOT : Botro ; KOR : Korhogo ; KAT : Katiola ; AGN : Agnibilékrou ; BON : Bondoukou ; G : Grains ; E : Epis ; SP : Spathes ; HUM : Humidity ; DOM : Damages ; PERT: Weight losses.

3.2. Discussion

The average humidity levels are all less than 15%, according to commercial standard of maize issued by Codex Alimentarius [19]. However, there is variability between the types of maize stored (grains, epis, spathes) as well as the departments visited. In fact, maize at grains remain the less humid type in most of the departments mainly in Botro, Korhogo and Katiola. Such data could result to the good practice of the drying of grains due to their easy handling. Some authors think that maize grains dry more easily and quickly and are more suitable for application of insecticides, compared to maize epis and spathes [22-23]. In the three localities mentioned above and even in Bondoukou, the humidity levels of samples taken from epis and spathes of maize are also less than 13%. This could make easier their conservation since the maximum humidity limit recommended for better maize conservation is 13% [22, 13-14]. Furthermore, some samples of maize spathes collected in the departments of Katiola, as well as the three types of maize collected in Agnibilékrou and Bondoukou provided water contents which are above 15%. An ineffective drying associated with inappropriate storage conditions of harvests could have caused rehumidification of maize. In fact, the production and the post-harvest treatment of agricultural products are influenced by environmental conditions mainly climate, temperature and rainfall [24-25]. Bad environmental conditions can create high humidity and activity of water and therefore favour development of pests (insects, fungi) in stores [26-27]. As corollary of development of these living beings, the attacked products experiences serious damages summed up by weight losses and damages of grains.

Concerning these damages induced by insects, a variability is more observed at different forms of maize and in departments of maize production. In fact, maize grains coming from the departments of Botro, Korhogo and Katiola are less attacked by insects than epis and spathes maize. Furthermore, the grains, epis and spathes of these three departments could benefit better conditions (structures and methods) of conservation than those coming from Bondoukou and Agnibilékrou. The investigations led by Niamketchi et al. [10] demonstrate a variability at level of structures of storage in area of production. According to these authors, the Botro's producers prefer polypropylene bags for storage of epis and grains maize in brick or clay stores. The palisades or palm leaf huts are used for storage of the maize spathes. As for the producers of korhogo, the conservation of grain, epi and spathe maize is done with cylindrical attic in clay. The use of trees whose branches serve as supports to hang spathes maize in the form of cluster. As for those of Katiola, they use polypropylene bags for the conservation of grains and palisades to hang the maize spathes. The producers of Agnibilékrou and Bondoukou store maize either in Chinese bamboo granaries with shelf or polypropylene bags exposed in store or a room of house [10].

Other studies revealed that the storage of harvests in inappropriate structures also favours the growth of insects and mould, especially when the humidity is high [1, 28-29]. Indeed, the increase of humidity strongly coincides with the amplification of damages leading to the significative losses of the harvests weight [9-10, 30-31]. Sankara *et al.* [12] as well as Danso *et al.* [29] revealed that the multiplication of insects pests is due to the storage conditions (grouped or isolated storage). They have also observed a diversity at the level of the number and the specie of insects pests according to the sites of studies.

Maizes coming from Bondoukou and Agnibilékrou are of lowest merchantability whatever the types of maize (epis, spathes, grains) : they are more humid and has strong rates of damages and weight losses. Consequently, their germination capacity and health quality mainly mycotoxin contamination (ochratoxin A, patulin, aflatoxins, zearalenone, fumonisin, deoxynivalenone), could be unsatisfactory. Some authors have shown that the attack by insects is source of weight loss, reduction of nutritive and merchantability and sometimes loss of the germination capacity of grains [1, 7, 9, 31-34]. According to them, the activity of the insects is also correlated with the development of moulds such as *Penicillium*, *Fusarium* et *Aspergillus*, responsible for the production of mycotoxins in stocks. The maizes stored in the form of grains or epis in the departments of Korhogo, Botro and Katiola are of better merchantability than those stored in spathes form. In fact, the grains and naked epis are better dried, less damaged by insects and have less weight losses.

4. CONCLUSION

This study investigated the merchantability of harvested and stored maize in five areas of production in Côte d'Ivoire. It shows that this merchantability is better in the departments of Botro ; Korhogo and Katiola regardless of the maize form (grains, epis , spathes) except for the maize spathes in Katiola. In these localities, maize grains and epis have also better merchantability than in the form of spathe. It would be important to sensitize producers on good post-harvest practices and the use of structures suitable for storing different forms of maize in order to help improve the profitability of their agricultural production and insure food security.

REFERENCES

1- Guèye MT, Seck D, Wathel J-P., LognayG. Pest control of cereal and legume stocks in Senegal and West Africa: bibliographic summary. *Biotechnol. Agron. Soc.* Approximately, 2012, 15 (1): 183-194.

- 2- Alene AD., Manyong VM., Tollens E., Abele S. The economic and poverty impacts of maizeresearch in West and Central Africa. *Agricultural Economics*, 2009, 40: 535-550.
- 3- FAOSTAT. Statistical databases on African countries “food commodities” trade, production, consumption, and utilization. FAO, Rome, Italy, available at: <http://www.uneca.org/sites/default/files/uploads/ice17-report-secu-aliment-industrie-fr-ecawa-2014-04-final.pdf>, 2014 .
- 4- Smale M., Byerlee D., Jayne T. Maize revolution in sub-Saharan Africa: Policy Research Working Paper No. 5659. Washington DC: World Bank, 2011.
- 5- FAOSTAT. Statistical databases on African countries “food commodities” trade, Production, consumption, and utilization. FAO, Rome, Italy, available at: <http://www.uneca.org/sites/default/files/uploads/ice17-report-secu-aliment-industrie-fr-ecawa-2014-04-final.pdf>, 2016.
- 6- Kouakou K., Akanvou L., Konan A., Mahyao A. Peasant strategies for maintaining and managing the biodiversity of maize (*ZeamaysL.*) In the department of Katiola, Côte d'Ivoire. *Journal of Applied Biosciences*, 2010, 33: 2100-2109.
- 7- Johnson F., N'Zi K., Seri-Kouassi, Foua-Bi K. Overview of storage problems and the impact of insects on the conservation of rice and maize in rural areas: the case of the Bouaflé - Coast region Ivory. *European Journal of Scientific Research*, 2012, 83: 349-363.
- 8- CNRA (National Center for Agronomic Research). Meteorological data from the Food Crops Research Station of Bouaké, 2014.
- 9- Niamketchi L., Biego G H., Sidibe D., Coulibaly A., Konan Y., Chatigre O. Changes in Aflatoxins Contents of the Maize (*Zea Mays L.*) Stored in Clay Granaries with use of Biopesticides from Rural Conditions and Estimation of their Intake”, *International Journal of Environmental & Agriculture Research*, 2016, Vol 2, Issue 5, pp 198-211.

- 10- Niamketchi L., Chatigre O., Coulibaly A., Konan Y., Biego M. H. Changes in the Quality of Maize (*Zea mays* L.) Post-Harvest Stored in Granaries with the Biopesticides from Rural Conditions in Côte d'Ivoire. *Global Journal of Biology, Agriculture & Health Sciences*; 2016, 5 (2): 74-87.
- 11- Pierre E., Adama C., Ysidor K., Daouda S., Olivier K. C., Godi HMB. Efficacy of *Lippia multiflora* (Verbenaceae) and *Hyptis suaveolens* (Lamiaceae) Leaves on Merchant Quality of Stored Maize Grain (*Zea mays* L.) in Côte d'Ivoire. *Journal of Agriculture and Ecology Research International*, 2017, 11(3): 1-10.
- 12- Sankara F., Sanou AG., Waongo A., Somda M., Toé P., Somda I. Farming practice after harvesting maize in the Hauts-Bassins region of Burkina Faso *Journal of Animal & Plant Sciences*, 2017, 33 (1): 5274-5288.
- 13- FAO. Assessment of land degradation in Senegal. FAO and Degradation Assessment Project. Preliminary report. April 2003. 59 p.
- 14- Mohale S., Medina A., Rodriguez A., Sulyok M., Magan N. Mycotoxigenic fungi and mycotoxins associated with stored maize from different regions of Lesotho. *Mycotoxin Research*, 2013, 29 : 209-219.
- 15- Pierre E., Constant K., Adama C., Daouda S., Leonce N., Ysidor K., Didier A., Olivier KC., Godi HMB. Daily Intake of Aflatoxin B1 and Ochratoxin a from Maize Grain (*Zea mays* L.) during the Storage with *Lippia multiflora* (Verbenaceae) and *Hyptis suaveolens* (Lamiaceae) Leaves in Côte d'Ivoire. *Asian Journal of Advances in Agricultural Research*, 2017, 3(4) : 1-13.
- 16- Amani MK., Koffi FK., Yao BK., Kouakou BD., Jean EP., Sekouba O. Analysis of climate variability and its influences on seasonal rainfall patterns in West Africa: case of the watershed N'zi (Bandama) in Ivory Coast. *European Journal of Geography*, 2010, 12: 200-222.

- 17- Kouassi AM., Kouamé KF., Goula BTA., Lasm T., Paturel JE., Biémi J. "Influence of climatic variability and modification of land use on the rain-flow relationship from 'a global modeling of the N'zi (Bandama) watershed in Ivory Coast', Ivoirian Science and Technology Review, 2008, vol. 11, 207-229.
- 18- AOAC. Official methods of analysis. Association of Official Analytical Chemists. Washington. DC. USA, 2000, 376 - 384.
- 19- Codex Alimentarius. Cereals, pulses, pulses and protein materials, 1st edition, Food and Agriculture Organization of the United Nations, Rome, Italy. ISBN 978-92-5-205842-7, 2007, 128 p.
- 20- Harris KL., Lindblad CJ. Post-harvest grain loss assessment methods- American Association of Agricultural Chemists, St Paul, Minnesota, 1978, 193 p.
- 21- Boxall R. A. A critical review of the methodology for assessing farm level grain losses after harvest. Report of the Tropical development and Research Institute, 1986, 191-139 p.
- 22- Gilliquet. Cereals in hot regions. AUPELF-UREF, Eds John LibbeyEurotext, Paris, 1989, 3-8.
- 23- IFDC (Developing Agriculture From the Ground Up). Training on Storage and Conservation of agricultural products. Training manual, 2016, 289p.
- 24- Comelade E. Technology and food hygiene: nutrients. Paris: Editions Jacques Lanore, 1990, 144 p.
- 25- Yoka J., Loumeto JJ., Djego JG., Houinato M., Aouango P. Adaptation of a cowpea cultivar (*Vigna unguiculata* L. (Walp.) To the pedoclimatic conditions of the Boundji area (Republic of Congo). *Afrique Science*, 2014, 10 (1) 217-225.
- 26- FAO. Packaging of cereal seeds and pulses. Rome: FAO, 1985, n ° 21. 17 p.
- 27- Kayombo MA., Mutombo TJM., Somue MA., Muka MP., Wembonyama OM., Kebe K., Sembène M. Cowpea (*Vigna unguiculata* (L.) Walp) field infestation by the bruchids

- (Coléoptera : Bruchidae) in the northern Senegal : preliminary biological and ecological data. *Journal of Applied Biosciences*. 2011, 41: 2788 – 2796.
- 28- Gwinner J, Harnisch R., Mück O. Manuel sur la manutention et la conservation des grains après récolte, GTZ, Eschborn, 1996, 368 p.
- 29- Danso JK., Osekre EA., Opit GP, Manu N., Armstrong P., Arthur FH., Campbell JF., Mbata G., McNeill. SG. Post harvest insect infestation and mycotoxins levels in maize markets in the middle belt of Ghana. *Journal of stored products research*, 2018, Vol 77. 9-15.
- 30- Chatta HS., Che MH., Teang SL., Benish NM., Muhammad RM. A study on the quality of wheat grain stored in straw-clay bin. *Journal of Biodiversity and Environmental Sciences*, 2015, 6 (1) : 428-437.
- 31- Konan KC., Fofana I., Coulibaly A., Koffi NE., Chatigre KO., Biego GHM. Optimization of Storage Methods of Cowpea (*Vigna unguiculata* L. Walp) Bagged PICS Containing Biopesticide (*Lippia multiflora*) By Central Composite Experimental Design in Côte d'Ivoire. *International Journal of Environmental & Agriculture Research*, 2016, 2(7) : 45-56.
- 32- Darfour B., Rosentrater KA. Maize in Ghana : an overview of cultivation to processing . In paper no162460492 from the 2016 ASABE annual international meeting. 11-16P. ASABE ST JOSEPH, MO. USA, 2016.
- 33- Niamketchi L., Biego G. H., Chatigre O., Didier A., Emmanuel K., Augustin A. Optimization of Post-Harvest Maize Storage using Biopesticides in Granaries in Rural Environment of Côte d'Ivoire. *International Journal of Science and Research*, 2015, 4 (9) : 1727-1736.
- 34- Kumar D., Kolita P. Reducing postharvest losses during storage of grain crops to strengthen food security in the developing countries. *Food Security*, 2017, 6 : 8-11.