

**Influence of prostate cancer on erectile dysfunction in Northern Cameroon and its management**

**Abstract**

Due to the lack of hospitals with available technical platform and diagnostic costs that are unaffordable by the majority of the population, the incidence of prostate cancer in Cameroon has increased with great impact on people's health.

**Aim:** This work was undertaken with the objective to determine the impact of prostate cancer on erectile dysfunction and its management.

**Methodology :** Sampling of the population was done in a comprehensive and non-probabilistic manner. Biopsies were taken from these patients to confirm and determine the form and stage of cancer followed by PSA assays. After the diagnosis was revealed, the testosterone assay was carried out in order to evaluate erectile function in the patients who equally completed a survey form made available to them in order to get an idea of their health history, the type of treatment followed and their lifestyle.

**Results:** Of the 75 patients received, 50 were participated in this study. The mean age of the patients was 67 years, with a predominance in the 60-70 age range. 85% of the patients had a Gleason score greater than or equal to 8. Of these patients, 42% had low testosterone levels (< 2.3 ng/mL), resulting to lack of morning erection (66.6%), loss of sexual desire (43.9%), difficulty having a spontaneous erection (88%). On the other hand, erectile dysfunction was revealed in the prostate cancer patients with low testosterone levels, with a history of hypertension (16.6%), diabetes (28.5%) alcohol consumption (44%), tobacco smoking (41%) and having undergone as prostate cancer treatment involving transurethral resection of the prostate (80%) and orchiectomy (20%).

**Conclusion:** The erectile dysfunction observed in patients suffering from prostate cancer in Northern Cameroon is mainly due to the evolution of the disease, as well as the health history of the patients (diabetes, hypertension).

**Key Words:** Prostate Cancer, Testosterone level, Erectile Dysfunction, Health history

## 1.Introduction

Cancers are pathologies mainly resulting from both the inability of cells to control their divisions and by the loss of mechanisms of programmed cell death. Modern lifestyle exposes us to various factors that may contribute to the activation of oncogenes [1–4], which in turn may induce disturbances in cell growth, differentiation, and apoptosis, thus, cancers [5,6].

The development of prostate cancer goes through four stages, namely stage I, localized without symptoms; stage II, locally advanced, characterized by a local tumor compressing in certain neighboring organs causing difficulty in urinating, frequent need to urinate, and sometimes hematuria; stage III, caused damage to neighbouring lymph nodes, marked by compression in the rectum and invasion of a urethra causing lumbar pain; stage IV, metastasis, is the most dangerous, as the tumour invades the whole body and is characterised by bone pain and/or alteration of the general condition [7].

Treatment varies according to the severity of the tumour, its location, the extent of proliferation, the tumour volume and the patient's life expectancy. It is known that in many cases, a prostate tumour is slow to grow, so there are different treatment options, including prostatectomy, external radiotherapy, chemotherapy, brachytherapy, and cryotherapy. Each of these first-line therapeutic modalities has advantages but are often associated with significant, life-altering side effects including urinary, bowel, and sexual dysfunction [8–10].

Erection is a physiological phenomenon in which the two *corpus cavernosum* dilate and become filled with blood, making the penis hard. However, as a result of certain physiological disturbances, erectile dysfunction may occur. It can be defined as the inability to achieve or maintain an erection of sufficient quality to have satisfactory sexual intercourse [11]. This problem is most often encountered in patients with evolving prostate cancer and manifests itself as sexual impotence, decreased libido, loss of erection [12]. As a result, out of 60% of men newly diagnosed with prostate cancer, 40% already have erectile dysfunction [8].

Fighting prostate cancer also include the identification of its impact on men's life. It is with this in mind that this study was initiated with the aim of determining the impact of prostate cancer on erectile dysfunction and its management in Northern Cameroon.

## **2- Materials and methods**

### **2.1- Participants and ethical considerations**

The present retrospective, cross-sectional and descriptive study was performed using historical data from the Urology Department of Ngaoundere Islamic Hospital, Ngaoundere, Cameroon, obtained between June 2018 and November 2019. The study included all the patients whose biological assessment mentioned the performance of a prostate biopsy, with the exception of patients whose diagnosis after biopsy was a non-malignant prostate tumour. The data collected were anonymized to protect the privacy of participants.

The study procedures were approved by the Ethics Committee of the Faculty of Medicine of the University of Ngaoundere and by the Institutional Review Board of Ngaoundere Islamic Hospital.

### **2.2- Procedures and data collection**

Information on patients' socio-demographic characteristics and history, including exposure to prostate cancer risk factors (Lifestyle, patient clinical history and treatment types ...) was obtained from the referral forms. Findings from DRE performed following clinical standard procedures (assessing the volume, consistency and regularity of the prostate) were also obtained from patient files, as well as other clinical data relevant for the study such as the PSA (Prostate Specific antigen) level, the Gleason score and testosterone level. Then, data were anonymized. The Urology Department of Ngaoundere Islamic Hospital obtained the Gleason scores of patients by applying Gleason classification to the anatomopathological findings of H&E stained prostate biopsies. The weight of the prostate determined using ultrasound was also available.

### **2.3- PSA (Prostate Specific antigen) assay**

Available data on the serum PSA level were obtained using the Chemiflex flexible assay protocol in the ARCHITECT i2000SR automatic immunoassay analyzer (Abbott Diagnostics, Lake Forest, IL, USA), as recommended by the manufacturer. Briefly, 75 uL of serum was pipetted and transferred into a microcentrifuge tube containing paramagnetic microparticles coated with anti-PSA antibodies. The tube was closed and shaken vigorously for 5 min with an agitator to allow an optimal binding of the serum PSA to the anti-PSA coated microparticles. Then, after rinsing, 75 uL of acridinium-labelled anti-PSA antibody conjugate was added to the mixture and incubated at room temperature for 15 minutes. Subsequently the pre-activation and activation solutions were added to the

reaction mixtures and the resulting chemiluminescent reaction was measured in relative units of URL light using ARCHITECT i2000SR.

#### **2.4- Testosterone assay**

It is an immunological assay that whose aim is to determine the total testosterone concentration in the human serum using the competitive immunodetection method. Initially, 30  $\mu\text{L}$  of displacement reagent was piped and transferred to the sample mixing tube and 75 $\mu\text{L}$  of serum piped and transferred to a mixing tube containing the displacement reagent. After closing the tube, the sample was homogenized by tube agitation and the mixture was incubated at room temperature for 3 minutes. 75  $\mu\text{L}$  of this mixture was piped and loaded into the detection buffer tube containing the fluorescence-labelled antibody (FL). 75  $\mu\text{L}$  of a sample mixture (complex) was piped and transferred to the sampling well on the cartridge. This complex is responsible for migrating to the nitrocellulose matrix, where the covalent pair of testosterone and bovine serum albumin (BSA) is immobilized on a test strip, and interferes with the binding of the target material and the marked antibody (FL.). The resulting reaction was measured in relative units of light.

#### **2.5- Statistical analysis**

Data were collected, organized, and presented as frequency distributions. Data analysis was performed using XLStat Version 2019. The  $\chi^2$  test of independence was used to determine the existence of significant relationships between the Gleason score and: (i) patients' lifestyle; (ii) patients' history of risk factors; (iii) treatment types. Statistical significance was set at  $P < 0.05$

### **3- Results**

The population included in this study consists of 50 prostate cancer patients who are married and over 50 years of age. Before the diagnosis, we carried out a number of check-ups, including PSA assay.

#### **3.1- PSA, Gleason Score and testosterone level of patients**

Table 1 below shows the distribution of the study population as a function of PSA, Gleason score, testosterone level. The figure shows that 76% of patients have a PSA level  $\geq 20$  ng/mL and 10% have a PSA level between 10-20 ng/mL. Indeed, 40% of patients have a Gleason score  $\leq 6$ , and 40% of patients have a Gleason score  $\geq 8$  and 20% have a

Gleason score 7. In addition, 42% of patients have a testosterone level  $\geq 3.5$ ng/mL, and 16% have a testosterone level between 2.3-3.5ng/mL.

**Table 1:** PSA, Gleason Score and testosterone level of patients

<b>Variables</b>	<b>Characteristics</b>	<b>%</b>
<b>PSA levels (ng/mL)</b>	< 10	14
	[10-20[	10
	>20	76
<b>Gleason Score</b>	<6	40
	7	20
	>8	40
<b>Testosterone level (ng/mL)</b>	< 2.3	42
	[2.3-3.5[	16
	> 3.5	42

### 3.2- Influence of Gleason score, PSA level on testosterone level

Table 2 below illustrates the influence of the Gleason score and PSA level on testosterone levels. The table shows that of the patients with low testosterone levels, 85% had a Gleason score of 8 or higher and 50% had a PSA level greater than 20 ng/mL. A significant association was found between Gleason score and testosterone level and between Gleason score and PSA level.

**Table 2:** Influence of Gleason score, PSA level on testosterone level

<b>Variable</b>	<b>Characteristics</b>	<b>Testosterone levels (ng/mL)</b>			<b>P-Value</b>
		<b>Low [1 - 2.3](%)</b>	<b>Intermediary [2.3 – 3.5](%)</b>	<b>Normal [3.5 – 8](%)</b>	
<b>Gleason Score</b>	6	5	20	75	<b>0.0001</b>
	7	30	20	50	
	> 8	85	10	5	
<b>PSA</b>	[0-10[	25	12.5	62.5	<b>0.0018</b>
	[10-20[	16.6	16.7	66.7	
	>20	50	16.6	33.4	

**PSA:** Prostate Specific antigen

### 3.3- Influence of lifestyle on testosterone levels

Table 3 below illustrates the lifestyle of the patients, showing that 41% of prostate cancer patients with low testosterone levels were tobacco smokers and 42% of patients were alcohol consumers. No significant association was recorded between tobacco

consumption and testosterone levels ( $P = 0.449$ ) or between alcohol consumption and testosterone levels ( $P = 0.228$ ).

**Table 3:** Influence of lifestyle on testosterone levels

Variable		Testosterone levels (ng/mL)			P-Value
		Low [1 – 2.3[ (%)	Intermediary [2.3 – 3.5[ (%)	Normal [3.5 – 8](%)	
Smoking	Yes	41	18	41	<b>0.449</b>
	No	42	29	29	
Alcoholism	Yes	44	18	38	<b>0.228</b>
	No	40	20	40	

### 3.4- Influence of patient clinical history on testosterone levels

Table 4 illustrates the influence of patient health history on testosterone levels, showing that 16.6% of patients with low testosterone levels had high blood pressure and 28.5% had diabetes. There was no correlation between these pathologies and testosterone levels ( $P=0.752$  and  $P=0.365$ ).

**Table 4:** Influence of patient clinical history on testosterone levels

Variable		Testosterone levels (ng/mL)			P-Value
		Low [1 – 2.3[ (%)	Intermediary [2.3 – 3.5[ (%)	Normal [3.5 – 8] (%)	
Hypertension	Yes	16.6	16.7	66.7	<b>0.752</b>
	No	54.4	18.1	27.5	
Diabetes	Yes	28.5	14.3	57.2	<b>0.365</b>
	No	44.2	16.3	39.5	

### 3.5- Characteristics of erectile dysfunction in prostate cancer patients

Table 5 below illustrates the characteristics of erectile function of patients with prostate cancer. It shows that 66% patients with low testosterone levels had no morning erection, 43% had loss of sexual desire and 88% had no spontaneous erection. Apart from the loss of sexual desire ( $P= 0.843$ ), a correlation was noted between the other parameters and the testosterone level.

**Table 5:** Erectile dysfunction in prostate cancer patients

Erectile dysfunction		Testosterone level (ng/mL)			p-value
		Low [1 – 2.3] (%)	Intermediary [2.3 – 3.5] (%)	Normal [3.5 - 8] (%)	
<b>Morning erection</b>	Yes	11.1	42.2	46.7	<b>0.004</b>
	Often	14.8	55.1	28.5	
	No	66.6	14.9	18.5	
<b>Sexual desire</b>	Yes	30	10	60	<b>0.843</b>
	No	43.9	19.5	36.6	
<b>Erection</b>	Absent	88.1	3.1	8.8	0.043

### 3.6- Influence of treatment types on testosterone levels

Table 6 illustrates the influence of treatment types on testosterone levels. It shows that 32.5% of prostate cancer patients who underwent transurethral resection of the prostate as treatment have low testosterone levels. 80% of prostate cancer patients who underwent orchiectomy as treatment have low testosterone levels. There was no correlation between treatment and decreased testosterone levels ( $P = 0.524$ ).

**Table 6:** Influence of treatment types on testosterone levels

Treatments	Testosterone levels (ng/mL)			P-Value
	Low [1-2.3] (%)	Intermediary [2.3 – 3.5] (%)	Normal [3.5 - 8] (%)	
<b>TURP</b>	32.5	20	47.5	<b>0.524</b>
<b>Orchidectomy</b>	80		20	

**TURP** : Transurethral resection of the prostate

#### 4- Discussion

Despite the slow and silent progression of prostate cancer, it causes changes in several areas: urinary, intestinal, vitality and sexual function [10]. Amongst these, erectile dysfunction (ED) appears as the most common impact on sexual function and is often accompanied by a loss of sexual desire or difficulty reaching orgasm [12] with a high impact on their quality of life [13]. Managing prostate cancer includes the identification of its impact on men's life. It is with this aim that this study was undertaken with the objective to determine the influence of prostate cancer on the sexual life of men especially on erectile function.

Biopsy results revealed that a large majority of patients (40%) had invasive, aggressive lesions represented by a Gleason score greater than 8. This could be justified by the fact that the most affected age group is in the range [60-70[ made up of people considered as elderly in Cameroon. Indeed, cellular ageing would seem to disrupt the physiology and integrity of prostate tissue, resulting in the accumulation of DNA damage and loss of control of the cell cycle by the transformation of proto-oncogenes into oncogenes or by the inactivation of tumour suppressor genes [14]. A statistically significant association is observed between the Gleason score and testosterone level ( $p < 0.0001$ ) showing that the more aggressive the tumour, the lower the testosterone level. Indeed, testosterone enters the prostate cell where it is largely metabolized in the cytoplasm into dihydrotestosterone by the enzyme 5-reductase [15]. Once fixed on the androgenic stromal cell receptor, it leads to the secretion of growth factors that will directly act on the proliferation and differentiation of epithelial cells. In addition, the stroma produces FGF7 and FGF10 factors which directly stimulate epithelial cell proliferation and activate the production and secretion of PSA, which goes directly into the nucleus of the prostate cell where it is able to bind directly to the DNA, thus exerting its action on cell proliferation, leading to an increase in prostate volume [15–17]. This increase in prostate volume will block the passage of testosterone which is produced by the testicles (Leydig cell), hence the low testosterone level observed in patients with a more aggressive tumour, leading to erectile dysfunction in most of them.

Although no statistical association was observed in this study between tobacco and alcohol consumption and a drop in testosterone levels ( $P = 0.449$  and  $P = 0.228$ ), several cancer patients who use these products have low testosterone levels (41% and 42% respectively for tobacco and alcohol). Indeed, there is accumulating evidence that smoking



increases the risk of fatal prostate cancer [18]. This is because the chemical compounds contained in tobacco smoke (nicotine, irritating substances, benzopyrenes, carbon monoxide) bind preferably in place of oxygen in the red blood cells. This leads to poor oxygenation of the penile muscles in the *corpus cavernosum*, which must become filled with blood during erection, thus leading to muscle weakness, which disturbs the erection process [19,20].

Patients with health history of diabetes and high blood pressure are more likely to develop erectile dysfunction although no significant statistical association is observed in this study. Indeed, 16.6% of patients with low testosterone levels had high blood pressure and 28.5% had diabetes. Diabetes induces a generalized alteration of endothelial cell function, responsible for the decrease in nitric oxide activity [21], which acts as an activator of guanylate cyclase to cyclic guanosine monophosphate (cGMP). cGMP is a nucleotide that promotes relaxation of the smooth musculature of the penile arteries and intracavernosal tissue, contributing to penile *corpus cavernosum* engorgement and erection [22]. Its absence or decrease will thus result in the difficulty of the penis to become filled with blood in order to achieve an erection. Also, diabetes is at the origin of several cases of neuropathies which constitutes a greater risk of erectile dysfunction [23]. On the other hand, high blood pressure is an independent risk factor of erectile dysfunction and even in treated hypertensives, probably due to the conjunction of several neurogenic (noradrenaline), hormonal (angiotensin II) and hemodynamic factors that determine endothelial dysfunction. All of these phenomena lead to a decrease in oxygen supply to the *corpus cavernosum*, inducing fibrosis of the *corpus cavernosum*. This irreversibly limits the possibilities of smooth muscle relaxation and expansion of the sinusoid spaces. We are thus witnessing a failure of the veno-occlusive mechanisms that makes any rigid erection impossible [21].

In our study all patients underwent transurethral resection of the prostate which consists in reducing the size of the prostate to treat advanced cancer in patients with a life expectancy of less than 10 years. 80% of the patients underwent transurethral resection of the prostate only and did not experience erectile dysfunction a few months after the resection but rather a small amount of ejaculate, as the prostate having decreased, there will be less production of sperm fluid. The orchiectomy was performed in patients with metastatic prostate cancer. It is a curative treatment which consists in the total stop of production of androgenic hormones which will lead to erectile dysfunction in the long run [24]. The association between the stage of symptomatic development of prostate cancer in

the body, age, and/or the presence of a co-morbidity related to smoking or the patient's lifestyle may be the reason for erectile dysfunction in prostate cancer patients.

### **Conclusion**

Results in this study showed that the erectile dysfunction observed in patients suffering from prostate cancer is due to the evolution of the disease as well as the lifestyle of the patients and their clinical history.

### **Consent**

Written informed consent was obtained from all the patients.

### **Ethical Approval**

The study procedures were approved by the Ethics Committee of the Faculty of Medicine of the University of Ngaoundere and by the Institutional Review Board of Ngaoundere Islamic Hospital.

### **Reference**

1. Djiogue S, Nwabo Kamdje AH, Vecchio L, Kipanyula MJ, Farahna M, Aldebasi Y, et al. Insulin resistance and cancer: the role of insulin and IGFs. *Endocr Relat Cancer*. févr 2013;20(1):R1-17.
2. Kipanyula MJ, Seke Etet PF, Vecchio L, Farahna M, Nukenine EN, Nwabo Kamdje AH. Signaling pathways bridging microbial-triggered inflammation and cancer. *Cell Signal*. févr 2013;25(2):403-16.
3. Makarem N, Bandera EV, Lin Y, Jacques PF, Hayes RB, Parekh N. Carbohydrate nutrition and risk of adiposity-related cancers: results from the Framingham Offspring cohort (1991-2013). *Br J Nutr*. juin 2017;117(11):1603-14.
4. Nair-Shalliker V, Yap S, Nunez C, Egger S, Rodger J, Patel MI, et al. Adult body size, sexual history and adolescent sexual development, may predict risk of developing prostate cancer: Results from the New South Wales Lifestyle and Evaluation of Risk Study (CLEAR). *Int J Cancer*. 1 févr 2017;140(3):565-74.
5. Nwabo Kamdje AH, Takam Kamga P, Tagne Simo R, Vecchio L, Seke Etet PF, Muller JM, et al. Developmental pathways associated with cancer metastasis: Notch, Wnt, and Hedgehog. *Cancer Biol Med*. mai 2017;14(2):109-20.

6. Seke Etet PF, Vecchio L, Nwabo Kamdje AH. Interactions between bone marrow stromal microenvironment and B-chronic lymphocytic leukemia cells: any role for Notch, Wnt and Hh signaling pathways? *Cell Signal.* juill 2012;24(7):1433-43.
7. Zerbib M, Zelefsky MJ, Higano CS, Carroll PR. Conventional Treatments of Localized Prostate Cancer. *Urology.* 1 déc 2008;72(6):S25-35.
8. Choo R, Long J, Gray R, Morton G, Gardner S, Danjoux C. Prospective survey of sexual function among patients with clinically localized prostate cancer referred for definitive radiotherapy and the impact of radiotherapy on sexual function. *Support Care Cancer.* juin 2010;18(6):715-22.
9. Donovan JL, Hamdy FC, Lane JA, Mason M, Metcalfe C, Walsh E, et al. Patient-Reported Outcomes after Monitoring, Surgery, or Radiotherapy for Prostate Cancer. *N Engl J Med.* 13 oct 2016;375(15):1425-37.
10. Watson E, Shinkins B, Frith E, Neal D, Hamdy F, Walter F, et al. Symptoms, unmet needs, psychological well-being and health status in prostate cancer survivors: implications for redesigning follow-up. *BJU International* [Internet]. 27 mars 2015 [cité 23 sept 2020]; Disponible sur: [https://www.research.ed.ac.uk/portal/en/publications/symptoms-unmet-needs-psychological-wellbeing-and-health-status-in-prostate-cancer-survivors\(29b17f29-9e68-47e3-b6ab-3f1f9dfed20c\)/export.html](https://www.research.ed.ac.uk/portal/en/publications/symptoms-unmet-needs-psychological-wellbeing-and-health-status-in-prostate-cancer-survivors(29b17f29-9e68-47e3-b6ab-3f1f9dfed20c)/export.html)
11. Pan F, Zhang J, Liu Y, Lu L, Qiu X, Lv K, et al. Intracavernosal Pressure Recording to Evaluate Erectile Function in Rodents. *J Vis Exp.* 06 2018;(136).
12. Schover LR, Fouladi RT, Warneke CL, Neese L, Klein EA, Zippe C, et al. Defining sexual outcomes after treatment for localized prostate carcinoma. *Cancer.* 15 oct 2002;95(8):1773-85.
13. Chambers SK, Chung E, Wittert G, Hyde MK. Erectile dysfunction, masculinity, and psychosocial outcomes: a review of the experiences of men after prostate cancer treatment. *Transl Androl Urol.* févr 2017;6(1):60-8.
14. Nicot C. Tumor Suppressor Inactivation in the Pathogenesis of Adult T-Cell Leukemia. *J Oncol.* 2015;2015:183590.
15. Tindall DJ, Rittmaster RS. The Rationale for Inhibiting 5 $\alpha$ -Reductase Isoenzymes in the Prevention and Treatment of Prostate Cancer. *J Urol.* avr 2008;179(4):1235-42.
16. Lasselin J, Drouin SJ, Champy CM, Léon P, Casenave J, Cussenot O, et al. [Influence of plasmatic testosterone during natural history of prostate cancer: a review]. *Prog Urol.* juin 2013;23(7):438-43.
17. Tu H, Gu J, Meng QH, Kim J, Strom S, Davis JW, et al. Low serum testosterone is associated with tumor aggressiveness and poor prognosis in prostate cancer. *Oncol Lett.* mars 2017;13(3):1949-57.
18. Islami F, Moreira DM, Boffetta P, Freedland SJ. A systematic review and meta-analysis of tobacco use and prostate cancer mortality and incidence in prospective cohort studies. *Eur Urol.* déc 2014;66(6):1054-64.

19. Safavy S, Kilday PS, Slezak JM, Abdelsayed GA, Harrison TN, Jacobsen SJ, et al. Effect of a Smoking Cessation Program on Sexual Function Recovery Following Robotic Prostatectomy at Kaiser Permanente Southern California. *Perm J.* 2017;21:16-138.
20. Sighinolfi MC, Mofferdin A, De Stefani S, Micali S, Cicero AFG, Bianchi G. Immediate improvement in penile hemodynamics after cessation of smoking: previous results. *Urology.* janv 2007;69(1):163-5.
21. Lue TF, Giuliano F, Montorsi F, Rosen RC, Andersson K-E, Althof S, et al. Summary of the recommendations on sexual dysfunctions in men. *J Sex Med.* juill 2004;1(1):6-23.
22. Choi BR, Kim HK, Park JK. Penile Erection Induced by Scoparone from *Artemisia capillaris* through the Nitric Oxide-Cyclic Guanosine Monophosphate Signaling Pathway. *World J Mens Health.* déc 2017;35(3):196-204.
23. Weinberg AE, Eisenberg M, Patel CJ, Chertow GM, Leppert JT. Diabetes Severity, Metabolic syndrome, and the risk of Erectile Dysfunction. *J Sex Med.* déc 2013;10(12):3102-9.
24. Krajewski W, Halska U, Poletajew S, Piszczek R, Bieżyński B, Matyjasek M, et al. Influence of Transurethral Resection of Bladder Cancer on Sexual Function, Anxiety, and Depression. In: Pokorski M, éditeur. *Clinical Medicine Research* [Internet]. Cham: Springer International Publishing; 2018 [cité 23 sept 2020]. p. 37-50. (Advances in Experimental Medicine and Biology). Disponible sur: [https://doi.org/10.1007/5584\\_2018\\_264](https://doi.org/10.1007/5584_2018_264)