

Original Research Article

COMPARING POULTRY MANURE TO INORGANIC FERTILIZER AND THEIR COMBINATION ON THE GROWTH AND YIELD PERFORMANCE OF MAIZE. (OMANKWA)

ABSTRACT

Aims: Major effect of soil fertility decline is the observed reduced food production in most African countries, including Ghana. In order to sustain soil and crop productivity, it is necessary to explore alternative soil fertility replenishment strategies, which are effective and affordable to farmers, especially the smallholder farmer.

Study design: A RCBD in three replications with each block with dimension 4m×3m using spacing of 0.75m and 0.4m inter and intra and inter-row.

Place and Duration of Study: The research work was conducted at the CSIR-Crops Research Institute, Kumasi between April 2016 - December 2016 using Omankwa a released maize variety from CSIR-Crops Research Institute, Kumasi, Ghana.

Methodology: There were six treatments per rep and each rep was randomized. Treatments were as follows; control (no fertilizer or poultry manure); T1 (100% Poultry manure two weeks after planting of maize), T2 (100% (N.P.K) two weeks after planting of maize), T3 (25% Poultry manure and 75% NPK fertilizer), T4 (50% Poultry manure and 50% NPK fertilizer) and T5 (75% Poultry manure and 25% NPK fertilizer).

Results: Results from the study showed that Omankwa performed better under treatment T1 better than other treatments T4 and T5 even though T4 and T5 perform better in other parameters measure for the experiment but did not results in grain yield for Omankwa.

Conclusion: The use of poultry manure still remains to be the best source as alternative to soil nutrient used to often better yield of crop production but if the total or required amount are not applied. NPK can be applied to obtain optimum grain yield in maize production.

Keywords: Omankwa, treatment, optimum, fertility, productivity

1. INTRODUCTION

On smallholder farms, soil fertility decline has been recognized as one of the major biophysical constraints affecting agriculture, particularly nitrogen (N) and phosphorus (P) deficiencies (Mokwunye et al., 1996). However, soil fertility has seldom been considered a critical issue by the development community who, until very recently have focused primarily on other biophysical constraints, such as soil erosion, drought, and the need for improved crop germplasm, especially in Africa (Lele, 1981; Eicher, 1982; Davis and Schirmer, 1987).

Most food crops in Ghana are produced by subsistent farmers, who over the years, practiced slash and burn and shifting cultivation to sustain yield even though at a low level. However,

increasing human population has led to intensive cultivation without adequately replenishing soil nutrients. The result has been the decline in crop yields and depletion of the resource base. The soils become fragile, quickly lose organic matter and nutrients when exposed to harsh environmental conditions or intensive cultivation.

Generally, Ghanaian soils are of low inherent fertility and therefore require external inputs to improve fertility. The use of mineral fertilizers is the most effective and convenient way to improve soil fertility. However, fertilizer use in Ghana has dropped to $< 5\text{kg NPK ha}^{-1}$ (MOFA, 1998), due mainly to the high cost of mineral fertilizers (Bumb, 1994; Gerner et al., 1995). Consequently, there is presently a serious negative balance in nutrient budgets of soils in the country posing a major constraint to sustainable soil management for increased crop growth and yield. Ironically, there is a large amount of organic waste that can be turned into fertilizers for crop production at low cost. Everywhere in the world people settle first in areas with high potential fertile soils, adequate rainfall and mild temperatures. As populations grow, soil nutrient capital is gradually depleted when farmers are unable to sufficiently compensate losses by returning nutrients to the soil via crop residues, manures and mineral fertilizers.

Increasing pressures on agriculture result in much higher nutrient outflows and the subsequent breakdown of many traditional soil fertility maintenance strategies. These traditional fertility maintenance strategies such as fallowing, intercropping cereals with legume crops, manure producing mixed crop livestock farming and opening new lands have not been replaced by an effective fertilizer supply (Sanders et al., 1996). Several decades of nutrient depletion have transformed originally fertile lands that yielded about 2 to 4 t ha⁻¹ of cereal grain into infertile ones where cereal crops yield less than 1 t ha⁻¹. The bulk of the food in Africa is produced on smallholder farms (Cleaver and Schreiber, 1994; Gladwin et al., 1997). One of the major problems affecting food production in Africa is the rapid depletion of nutrients in smallholder farms (Badiane and Delgado, 1995). This is because the smallholder farmer is poorly resourced and unable to invest in soil fertility inputs, particularly mineral fertilizers. This is not surprising since about half of Africa's population is classified as "absolute poor" subsisting on per capita incomes of less than 1 US\$ per day (Badiane and Delgado, 1995). The situation is critical especially when the poor farmer has to bear the full cost of production owing to the removal of subsidies on mineral fertilizers.

Therefore, the use of inorganic fertilizers results in high cost of maize production, destruction of the soil and aquatic life when leached into water bodies. Salt accumulation which develops as a result of repeated application of inorganic fertilizers forces the plant to expend more energy to draw water from the soil and can cause them to appear wilted or dried out. The general objective of the study is to compare poultry manure, inorganic fertilizer N.P.K (15-15-15) and their combination on the growth and yield of maize (Omankwa).

2. MATERIAL AND METHODS

2.1 Location of the study area

The research work was conducted at Crops Research Institute Fumesua, Kumasi-Ashanti Region. Fumesua is a Metropolitan Assembly in Kumasi. It is situated on the Accra and Kumasi high way, under Ejisu -Juaben district in the Ashanti region. It is located between latitude 6.8 North and longitude 1.8 west of the equator. The area is a dense forest belt and experiences two types of rain fall pattern which are the major rain and the minor rain pattern,

the dry season from November to January (Meteorological Service Department, Ashanti, 2016).

2.2 Experimental design and Management

Before planting, soil samples from the experimental site were collected at 0-15cm and 15-30cm depth for analysis of organic carbon, total nitrogen using standard methods, extractable P, Ca, Mg, K, Na using Mehlich-1 (M1) extraction method, where P and Mg²⁺ were determined colour metrically in a spectrophotometer and Ca²⁺ and K⁺ were determined using flame photometer. Results of the soil analysis is as showed in Table 1. The field was ploughed and left as such for one-week. Total land area of 16m×12m was used with each plot size measuring 4m×3m using spacing of 0.75m and 0.4m inter and intra and inter-row respectively in a RCBD in three replications. The blocks were further divided into six treatments per plot with 1m alley between them. The treatments were; T0 – Control, T1 – 100% Poultry manure two weeks after planting of maize, T2 - 100% (N.P.K) two weeks after planting of maize, T3 – 25% Poultry manure and 75% NPK fertilizer, T4 – 50% Poultry manure and 50% NPK fertilizer and T5 – 75% Poultry manure and 25% NPK fertilizer constituted from recommended rate of 4 ton/ ha (poultry manure) and 250kg/ha as showed in Table 2.

Tables 1. Compositional soil analysis of the three-study areas

Parameters	Fumesua		Landon (1991) interpretation	
	0-15 cm	15-30 cm	High	Low
pH	3.64	3.47	>6.5	<5.8
Total Nitrogen (%)	0.11	0.07	>0.5	<0.2
Organic Carbon (%)	3.03	2.47	>10.0	<4.0
Organic Matter (%)	5.22	4.26	>10.0	<2.0
Ca (Cmol/Kg)	42.4	30.8	>10.0	<4.0
Mg (Cmol/Kg)	2.6	4.6	>4.0	<0.5
K (Cmol/Kg)	0.36	0.28	>0.6	<0.2
Na (Cmol/Kg)	0.94	0.8	>1.0	<1.0
Al (Cmol/Kg)	0.5	0.67		
H (Cmol/Kg)	0.33	0.33		
P (mg/Kg)	53.47	32.28	>50.0	<15.0
SAND (%)	80	84		
CLAY (%)	10	10		
SILT (%)	10	6		
TEXTURAL CLASS	Sandy Loam	Loamy fine sand		

Table 2: Treatment Description

No	Description of Treatment	Abbreviation
1	Control (No Fertilizer)	T0
2	100% Poultry manure two weeks after planting of maize	T1
3	100% (N.P.K) two weeks after planting of maize,	T2

4	25% Poultry manure and 75% NPK fertilizer	T3
5	50% Poultry manure and 50% NPK fertilizer	T4
6	75% Poultry manure and 25% NPK fertilizer	T5

2.3 Data Collection

Data for growth was collected after two weeks of planting and continue after every two weeks based on the following parameters.

- measuring the stem girth
- leave length,
- Number of leaves per plant
- Plant height
- Ear height

Data on yield will be analysis after harvesting based on

- Cob girth/width
- Number of cob/plants
- Grains weight
- Grain moisture
- Field weight

2.4 Data Analysis

Data of maize yield and growth parameters were analyzed using GenStat to run analysis of variance (ANOVA) and result will be presented in tables and at LSD of ($p=0.05$).

3. RESULTS AND DISCUSSION

The analysis of variance for grain yield and yield related traits for Omankwa under various soil amendments at Fumesua are given in Table 3 and Table 4. Analysis of variance showed significant differences for leaf length (cm), Stem girth (cm), grain yield (ton/ha), field weight (kg) and the number of leaves at 4WAP at the experimental site. The significance difference among the treatment indicates the positive responses of the various treatments used for grain yield which could be exploited for improvement of the respective treatments. Results from the experiment showed that leaf length were significantly difference even though T1 recorded the highest leaf length and the treatment T0 recorded the least length as showed in Table 3 and 4.

Stem girth of Omankwa showed that stem girth was not significantly different. T5 recorded the highest stem girth and lowest T0. In this study the increase in available nitrogen levels influenced positively the stem girth and ear length of maize plant. These findings were also reported by Okumura et al. (2011) and Santos et al. (2002). The ear height of Omankwa under treat T4 recorded the highest and the lowest T0. Ear height recorded no significant difference among the treatments used for the experiment. Treatment T5 recorded the highest height among the treatment and T0 the lowest height as showed in Table 1. No significant difference was recorded among various plants under different treatments as showed in Table 3 and 4.

Treatment T4 recorded the highest weight at the field and the lowest weight at T0 treatment showing no significant difference among the treatments used for the experiment. Increase in grain weight and yield was promoted by adequate nitrogen supply, because nitrogen absorbed by plants in responsible by fixation of carbon skeletons to amino acids

synthesis (Marschner et al., 1995), which results in several protein that have specific functions in plant metabolism. On the average the chlorophyll content index of T1 recorded the highest value compared with T0 which recorded less showing significant difference among treatments. T4 treatment recorded the highest girth compared with T0 treatment showing no significant different among the treatments used for the experiment. In addition, increase in chlorophyll content index and stem girth are needed during grain filling period these carbon compounds previously fixed are broken down, transported and stored in form of proteins and amino acids (Okumura et al., 2011). Gul et al., (2015) reported that maize grain yield was linearly influenced by nitrogen level applied. Bashir et al. (2012), Okumura et al. (2011), Deparis et al. (2007), Cruz et al., (2008) and Bastos et al. (2008) showed also linear behaviour linked to yield in maize induced by increase in nitrogen level as showed in Table 3 and 4.

No significant difference was recorded when the number of cobs per plant were taken during the experiment. All treatments used for the experiment the same number of cobs per plant. T2 treatment had 50% tasselling latest and the earliest been T4 showing no significant different among the treatments used for the experiment. Treatments T1, T4 and T5 had its silk appearing earlier than T2 showing significant difference among the treatments as showed in Table 3 and 4.

There was no significant difference when the number of leaves were determined at 2 WAP but slight difference in number of leaves were observed when they were counted after 4WAP with the least recorded for T0 with no significant difference with the minimum count of 8 and maximum 11. No significant difference was recorded when taken after 6WAP but T4 recorded the highest of 13 leaves and T0 recording the lowest as showed in Table 3 and 4. Biomass (leaf and stem dry weight) and yield components (ear weight, ear length, ear diameter and 100 seed weight) were also significant increased with the application of poultry manure which resulted in an overall increase in grain yield per hectare

Table 3. Analysis of variance of yield and yield related performance of Omankwa maize variety.

Sources of variation	DF	Length of Leaf	Stem Girth	Ear Height	Plant Height	CCI	Grain Yield	Field Wt.	Cob Girth	DAS	D
Rep	3	29.6871	0.1737	45.0771	132.7217	35.0828	19.2708	0.1526	0.0897	6.1111	1
Treatment	5	76.3368*	1.4964*	84.9227	246.9444	51.4157	22.4195*	0.2594*	0.1786*	4.5667	1
Error	15	19.1514	0.3471	39.121	117.6065	84.6301	7.5687	0.0703	0.0503	3.7444	2
Total	23										
<i>Descriptive Data</i>											
Min		66.1	4.8	46.36	104.64	5.2	1.39	0.6	3.07	53	5
Max		89.2	8.2	68.18	147.82	39.5	13.18	1.9	4.33	61	5
Mean		80.14	6.5	56.5	128.21	16.06	6.33	1.2	3.77	56.67	5
Std Dev		5.74	0.7578	7.06	12.15	8.42	3.51	0.3495	0.2886	2.06	1
CV (%)		5.46	9.06	11.07	8.46	57.29	43.44	22.02	5.95	3.41	2

Table 4: Means of various yield and yield related performance of Omankwa maize variety.

Treatment	N	Length Leaf	Stem Girth	Ear Height	Plant Height	CCI	Grain yield	Field Wt.	Cob Girth	DAS	DAT	No Leaves of 2WAP	No Leaves of 4WAP	No Leaves of 6WAP
T ₀	4	73.7b	5.55b	50.12	115.82	10.75	2.72a	0.85b	3.46b	57.25	53.25	5.25	9b	11.5
T ₁	4	84.1a	6.85ab	58.86	131.55	16.1	6.41a	1.27ab	3.88ab	56.25	52.75	5.25	9.75ab	12.25
T ₂	4	76.28ab	6.05ab	52.16	126.16	21.82	5.07a	1.03ab	3.62ab	58.5	53.75	5	9.5ab	11.75
T ₃	4	79.67ab	6.45ab	55.91	123.82	14.57	5.99a	1.2ab	3.81ab	56.5	52.75	5	9.5ab	12
T ₄	4	83.65ab	7a	61.5	135.59	16.82	8.88a	1.6a	4.08a	55.5	52	5.25	10.75a	12.75
T ₅	4	83.42ab	7.12a	60.46	136.34	16.27	8.93a	1.27ab	3.76ab	56	53.25	5.75	10.25ab	12.5

Means with the same letter is not significantly different

4. CONCLUSION

Based on the findings of this study, it can be concluded that maize growth parameters (stem girth and ear length) increase significantly with increase in nitrogen level. Maize yield characters, namely ear weight and 100 grain weight increased significantly with increased in nitrogen level. Maize grain yield was increased by nitrogen rates evaluated. It was possible to visualize relationships amongst maize growth characters, yield parameters and maize grain yield. The use of poultry manure still remains to be the best source of soil nutrient used to often better yield of crop production but if the total or required amount are not applied. NPK can be applied to obtain optimum grain yield in maize production.

REFERENCES

- Badiane, O. and Delgado, C. L. (1995). A 2020 vision for food, agriculture and the environment, Discussion Paper 4. International Food Policy Research Institute, Washington, DC
- Bashir, S. (2012). Response of brown sarson to NPK application under early, normal and late sown conditions (Ph.D thesis), Division of Agronomy, SKUAST-K. 223p
- Bastos, E. A., Cardoso, M. J., Melo, F. B., Ribeiro, V. Q. and Andrade Junior, A. S. (2008). Rates and timing of nitrogen application for obtaining the economic grain yield, under no tillage, *Revista Ciencia Agronomica*. Vol: 39:275-280.
- Bumb, B. I. (1994). World nitrogen supply and demand: An overview. In Bacon, P. E. (ed.) *Nitrogen Fertilization and the Environment*. Marcel Dekker Inc. New York, USA.
- Cleaver, K. M. and Schreiber, G. A. (1994). Reserving the spiral; the population, agriculture and environment nexus in sub-Saharan Africa. World Bank, Washington, DC.
- Cruz, S. C. S., Pereira, F. R. S., Santos, J. R., Albuquerque, A. W. and Pereira, R. G. (2008). Nitrogen fertilization for corn cultivated under a no tillage system in the state of Alagoas, Brazil. *Revista Brasileira de Engenharia Agricola e Ambiental*. Vol: 12: 62-68.
- Davis, T. J. and Schirmer, I. A. (ed.). (1987). Sustainability issues in agricultural development. Proc. Of the Seventh Agriculture Sector Symposium. World Bank, Washington, DC.
- Eicher, C. K. (1982). Facing up to Africa's food crisis. *Foreign Affairs* 61(1): 154-174.
- Gerner H., Asante E. O., Owusu-Bennoah E. and Marfo K. (1995). Ghana fertilizer privatization scheme: Private Sector Roles and Public Sector responsibilities in meeting needs of farmers. IFDC-Africa, Lome, Togo
- Gladwin, C. H., Buhr, K. L., Goldman, A., Hiebsch, C., Hildebrand, P. E., Kidder, G., Langham, M., Lee, D., Nkedi-Kizza P. and Williams, D. (1997). Gender and soil fertility in Africa. In: R. J. Buresh, P. A. Sanchez and F. Calhoun (eds.): *Replenishing soil fertility in Africa*. ASA-SSSA Special Publication 51: 219. American Society of Agronomy, 107Madison, W1, USA
- Gul, S., Khan, M. H., Khanday, B. A., and Nabi, S. (2015). Effect of sowing methods and NPK levels on growth and yield of rainfed maize (*Zea mays* L.) *Scientifica*. Vol 2015, Article ID 198575, 6p.
- Lele, U. (1981). Rural Africa: modernization, equity and long-term development. *Science* 211(4): 557-583.
- Marshner, H. (1995). *Mineral nutrition of higher plant* 2nd edn. Academic press, New York, 889p.
- Meteorological Service Department, Ashanti.
- Mokwunye, A. U., de Jager, A. and Smaling, E. M. A. (1996). Restoring and maintaining the productivity of West Africa Soils: Key to sustainable development. International Fertilizer Development Center (IFDC), Muscle Shoals, Alabama. 94p
- Okumura, R. S, Takahashi, H. W., Santos, D. G. C., Lobato, A.K. S., Mariano, D.C., Marques. O. J., Silva, M. H. L. S., Neto, C. F. O., and Lima Junior, J. A. (2011). Influence of different nitrogen levels on growth and production parameters in maize plants. *Journal of food, agriculture & Environment*. Vol 9 (3 & 4): 510-514. www.world-food.net.

Sanders, J. H., Shapico, B. I. and Ramaswamy, S. (1996). The economics of agricultural technology in semi-arid sub-Saharan Africa. John Hopkins University Press, Baltimore, MD, USA. 303p

Santos, P. G., Juliatti, F. C., Buiatti, A. L and Hamawaki. O. T. (2002). Evaluation of agronomic performance of corn hybrids in Uberlandia, MG, Brazil. Pesquisa Agropecuaria Brasileira. Vol. 37: 597-602.

UNDER PEER REVIEW

APPENDIX

APPENDIX 1. RAINFALL FIGURES FROM EJURA, WENCHI AND FUMESUA (2016)

LOCATION	MONTH	JANUARY	FEBRUARY	MARCH	APRIL	MAY	JUNE	JULY	AUGUST	SEPTEMBER	OCTOBER	NOVEMBER	DECEMBER
FUMESUA	2016	2.8	0	90.1	85.7	149.3	90.9	134.9	496.5	214.31	194	67.6	15.6

APPENDIX 2. MAXIMUM AND MINIMUM TEMPERATURE FROM EJURA, WENCHI AND FUMESUA (2016)

		JANUARY		FEBRUARY		MARCH		APRIL		MAY		JUNE		JULY		AUGUST		SEPTEMBER		OCTOBER		NOVEMBER		DECEMBER	
LOCATION	YEAR	MAX	MIN	MAX	MIN	MAX	MIN	MAX	MIN	MAX	MIN	MAX	MIN	MAX	MIN	MAX	MIN	MAX	MIN	MAX	MIN	MAX	MIN	MAX	MIN
FUMESUA	2016	32.7	21.1	35.4	23.3	34.4	23.5	32.8	24.1	28.1	23	30.3	22.2	27.9	22	27.4	21.7	30	22.1	30.7	22.3	31.6	22.4	32.5	21.5