

TYPE OF MANUSCRIPT: An in vitro study

Anti-Microbial Efficacy Of Commercially Available Herbal Dentifrices On Oral Microbial Flora- An In vitro Study.

Running title: Anti-Microbial Efficacy Of Commercially Available Herbal Dentifrices On Oral

ABSTRACT:

Dental caries is a multi-factorial disease. Various chemically derived antimicrobial agents are incorporated to dentifrices to curb cariogenic microorganisms. There has been renewed interest in naturally occurring products in recent years, the side effects encountered with the use of synthetic formulations led to the surge for novel alternatives. To compare the efficacy of herbal dentifrices on oral microflora using antibiotic susceptibility tests. Two strains of microorganisms- Streptococcus mutans, and Candida albicans were taken and incubated in Mutans media, Sabouraud Dextrose agar respectively. Different dilutions (1:5, 1:10, 1:15) of several brands of toothpastes with different compositions were made. Sterile discs were incorporated with equal amounts of prepared toothpaste formulations. Then these discs were placed equidistant to each other followed by plates were incubated for 24 hours. Zone of inhibition is found to be higher in Ayush (30 mm), followed by Dant kanti, Colgate vedshakti, Meswak, Dantajeevan and Dabur red showing inhibition zones of 26 mm, 24 mm, 17 mm and 14 mm respectively. Results show that herbal toothpaste formulations were effective in controlling the oral microflora. The antimicrobial properties of the six dentifrices concluded that almost all the commercially available herbal dentifrices had antimicrobial activity to some extent.

Keywords: Antibacterial activity, fluoride, herbal, micro-organisms, toothpaste,

INTRODUCTION:

Dental health has been defined by the WHO, as a state of complete normality and functional efficiency of the teeth and supporting structures, and that of the surrounding part of the oral cavity, and the various structures related to mastication and the maxillofacial complex. Dental caries is a multifactorial disease and is the outcome of a multiple complex process involving factors like diet, microorganisms, trace elements, saliva, genetic predisposition and tooth morphology.

Oral cavity harbours wide variety of microorganisms, these are considered crucial for the initiation and progression of dental diseases[1]. Over 400 species of microbes inhabit as commensals in the oral cavity of a healthy adult. An aberration to this ecology due to dietary habits, improper oral hygiene or systemic factors lead to increased cariogenic microorganisms[1,2,3].

Cariogenic microorganisms like *Streptococcus mutans* and *Lactobacillus acidophilus* are the primary causative microorganisms for the development of dental caries. These cariogenic microorganisms encourage the accumulation and adherence of plaque biofilm by metabolizing sucrose into sticky glucan. The microorganisms in dental plaque degrade the dietary

carbohydrates producing lactic acid leading to localized demineralization and the eventual formation of dental caries[3–5] . Plaque – induced caries is a local disease, therefore, the local use of antimicrobial agents is more efficient than their systemic use[6]

Poor oral hygiene is one of the reasons for accumulation of microbes and their harmful activities [6–8]. Dental diseases are effectively avoidable by simple inexpensive and easy to practice personal hygiene habits [9,10]. Tooth paste is used to promote oral hygiene. Triclosan is a common antibacterial agent in the tooth paste.

Besides, the use of chemical agents has been proposed as a potential prophylactic method of reducing plaque-mediated disease[11]. Most effective of them is the tooth brushing habit. Of various factors of this practice, antibacterial efficacy of the tooth paste has a major role to play in the outcome [12][13].

Recently, there has been increased interest in the various medicinal plants which have demonstrated anti-cariogenic activity in various studies. Herbal products contain antimicrobial agents and these formulations may be more appealing as they do not require alcohol, artificial preservatives, flavors or colours for their activity.

Considering the overflowing number of brands of numerous toothpastes in the market, the efficacy of these toothpastes in controlling the bacterial count has to be scientifically analyzed. Rising popularity of herbal and natural products has mandated dental professionals to evaluate the effectiveness of these products and provide evidence-based suggestions to their patients to make a better choice[14] Also there has been an increasing interest in the study of medicinal plants and their traditional use in different parts of the world over the last decade.

Antibiotic susceptibility testing is the method employed for testing the effectiveness of agents against specific microorganisms [15]. A variety of laboratory methods can be used to evaluate or screen the in vitro antimicrobial activity of an extract or a pure compound. The most known and basic methods are the disk-diffusion and broth or agar dilution methods. Disk-diffusion assay offers many advantages over other methods such as simplicity, low cost, the ability to test enormous numbers of microorganisms and antimicrobial agents, and the ease to interpret results provided. The minimum inhibitory concentration is the minimum (lowest) concentration of an antibiotic that will inhibit the growth of a bacterial strain. According to the epidemiological triad for the dental caries causes, there is a requirement to evaluate the host factors such as oral health knowledge, oral hygiene practices, dental visits, and eating habits among the school going children. We have successfully completed numerous epidemiological studies [16–22][23] and in vitro studies [24][25][26][27] for the betterment of our community [28][29][30][31][32][33]. Hence, an in vitro study was done to compare the antimicrobial efficacy of herbal dentifrices on *Streptococcus mutans* and *Candida albicans*.

MATERIALS AND METHODS:

Study type: In-vitro study.

Study organisms: *Streptococcus mutans*, *Candida albicans*.

Streptococcus mutans is a facultative anaerobic, gram-positive coccus (round bacterium) commonly found in the human oral cavity and is a significant contributor to tooth decay. This bacterium, along with the closely related species *Streptococcus sobrinus*, can co-habit the mouth. Both contribute to oral disease, and the expense of differentiating them in laboratory

testing is often not clinically necessary. Therefore, for clinical purposes they are often considered together as a group, called the mutans streptococci.

Candida albicans causes oral thrush in which curd-like white patches form inside the mouth, on the tongue and palate and around the lips[15,34]. It may also cause cracked, red, moist areas of skin at the corner of the mouth. *Candida* species causes infections in immuno-compromised patients, individuals on drug therapy, and the chronically ill[35].

Prior to the start of the study ethical clearance was obtained from the institutional ethics committee, Saveetha university. The study was scheduled in the month of October 2017.

Preparation of dentifrices:

Solutions of selected dentifrices were made by mixing 1 gram of dentifrices in 4 ml of distilled water to give 1:5 dilutions, in a sterile container. Further dilutions were made by mixing 1 gram of tooth paste with 9 ml and 14 ml to give 1:10 and 1:15 dilutions respectively.

Anti-microbial assay:

Two strains of microorganisms –*Streptococcus mutans* and *Candida albicans* were taken. The media used were Mutans media and Sabouraud dextrose agar media respectively for each microorganism respectively. These bacteria were inoculated in their respective medium by swab method.

Disc diffusion method:

Sterile discs were incorporated with equal amounts of toothpaste formulations using micropipette. Then the discs were placed equidistant to each other following which these plates were incubated for 24 hours.

Measurement of zone of inhibition:

The test plates were held in front of a desk lamp, and the zones were measured with a ruler held against the back of the Petri plate. The diameters of the zones of inhibited growth were measured to the nearest whole millimeter.

RESULTS AND DISCUSSION:

Results show that the zone of inhibition of various types of herbal dentifrices against *Streptococcus mutans*. Zone of inhibition is found to be higher in Ayush (30 mm), followed by Dant kanti, Colgate vedshakti, Meswak, Dantajeevan and Dabur red showing inhibition zones of 26 mm, 24 mm, 17 mm and 14 mm respectively. Only Dantajeevan shows inhibition at 1:10 and 1:15 dilutions respectively. It also shows the zone of inhibition of various types of herbal dentifrices against *Candida albicans*. Zone of inhibition is found to be higher in Ayush and Dant kanti with 18 mm zone of inhibition followed by Colgate vedshakti, Meswak, Dantajeevan and Dabur red showing inhibition zones of 12mm and 10 mm respectively. Only Dant kanti shows inhibition at 1:10 dilution respectively.

Microbial biofilms are the cause for the dental caries and periodontal disease; therefore it is very important to control these biofilms by mechanical debridement which can be achieved with the use of adjunctive antimicrobials in the tooth pastes. Poor oral hygiene is solely responsible for

prevalence of dental caries in our community[36]. Key to the prevention of dental diseases is maintenance of good oral hygiene. Herbal extracts have received special attention because of being non-chemical and non-synthetic in nature, and have been used in traditional medicine [36,37][38][39]. Interest in natural-based toothpastes has increased recently[40].

As there is a paucity of literature regarding the efficacy of different types of herbal dentifrices against oral microflora, this in-vitro study was conducted to test the antimicrobial efficacy of different types of herbal dentifrices against *Streptococcus mutans* and *Candida albicans*.

Antimicrobial efficacy of six different herbal dentifrices against *Streptococcus mutans* and *Candida albicans* were tested. Three different dentifrice formulations 1:5, 1:10 and 1:15 were made by mixing distilled water with the dentifrice. The media used were Mutans media and Sabouraud dextrose agar media respectively for each microorganism respectively by swab method. The antimicrobial efficacy is then tested using a disc diffusion method.

In our study, antimicrobial efficacy formulations of dentifrice formulations against *Streptococcus mutans* were measured as zones of inhibition. The zone of inhibition is found to be higher in Ayush (30 mm), followed by Dant kanti, Colgate vedshakti, Meswak, Dantajeevan and Dabur red showing inhibition zones of 26 mm, 24 mm, 17 mm and 14 mm respectively. Only Dantajeevan shows inhibition at 1:10 and 1:15 dilutions respectively. Similar results were obtained in the study done by Sabiha Shaheen et al[41] in which herbal toothpastes such as Dant kanti show maximum antimicrobial efficacy. Also, few studies show that herbal toothpastes have similar and better antimicrobial activity. During the last few decades there has been an increasing interest in the study of medicinal plants and their traditional use in different parts of the world[42]

Zone of inhibition of various types of herbal dentifrices against *Candida albicans* shows that Zone of inhibition is found to higher in Ayush and Dant kanti with 18 mm zone of inhibition followed by Colgate vedshakti, Meswak, Dantajeevan and Dabur red showing inhibition zones of 12mm and 10 mm respectively. Only Dant kanti shows inhibition at 1:10 dilution respectively. Similar results were obtained in the study conducted by Farogh Gabraiel et al[7]; in which herbal toothpastes gave maximum zones of inhibition against *Candida albicans*. Okpalugo et al[43] reported that the toothpaste brands considered for their study were not effective in reducing the oral microorganisms whereas the results of our study showed inhibition by all brands.

Herbal dentifrices show antimicrobial activity against the oral microflora hence it becomes necessary to preserve this traditional system of medicine by proper documentation and identification of species[44].

CONCLUSION:

The present study was based on in-vitro experiments. Results of this study have shown that herbal toothpaste formulations were effective in controlling the oral microflora. The antimicrobial properties of the six dentifrices concluded that almost all the commercially available herbal dentifrices had antimicrobial activity to some extent to benefit dental health or antiplaque action.

Based on the results of the present study, it can be concluded that all herbal dentifrices exhibited antimicrobial activity against the selected oral microorganisms, with Ayush and Dant Kanthi (DK) being the most effective. Hence, it can be inferred that herbal dentifrices can also be recommended like the conventional formulations. However, future research efforts are needed for the evaluation of quality and efficacy of the herbal dentifrices for its regular use in the oral

hygiene products. Further studies on the safety and efficacy of such products need to be established. Thus turning to nature can be as good as the latest advances.

UNDER PEER REVIEW

REFERENCES:

[1]Nayak SS, Ankola AV, Metgud SC, Bolmal UK. An in vitro Study to determine the

Effect of Terminalia chebula Extract and Its Formulation on Streptococcus mutans. The Journal of Contemporary Dental Practice 2014;15:278–82. <https://doi.org/10.5005/jp-journals-10024-1528>.

[2]Agarwal G, Chatterjee A, Saluja M, Alam M. Green tea: A boon for periodontal and general health. Journal of Indian Society of Periodontology 2012;16:161. <https://doi.org/10.4103/0972-124x.99256>.

[3]Subramaniam P, Reddy KRM, Eswara U. Effect of different types of tea on Streptococcus mutans: An in vitro study. Indian Journal of Dental Research 2012;23:43. <https://doi.org/10.4103/0970-9290.99037>.

[4]Bedrood Z, Rameshwaram, Hosseinzadeh H. Toxicological effects of Camellia sinensis (green tea): A review. Phytotherapy Research 2018;32:1163–80. <https://doi.org/10.1002/ptr.6063>.

[5]Taylor PW, Hamilton-Miller JMT, Stapleton PD. Antimicrobial properties of green tea catechins. Food Science Technology Bulletin: Functional Foods 2005;2:71–81. <https://doi.org/10.1616/1476-2137.14184>.

[6]Lobo PLD, de Carvalho CBM, Fonseca SGC, de Castro RSL, Monteiro AJ, Fonteles MC, et al. Sodium fluoride and chlorhexidine effect in the inhibition of mutans streptococci in children with dental caries: a randomized, double-blind clinical trial. Oral Microbiology and Immunology 2008;23:486–91. <https://doi.org/10.1111/j.1399-302x.2008.00458.x>.

[7]Anushree DB, Anushree B. Comparison of Antimicrobial Efficacy of Triclosan-Containing, Herbal and Homeopathy Toothpastes- An Invitro Study. JOURNAL OF CLINICAL AND DIAGNOSTIC RESEARCH 2015. <https://doi.org/10.7860/jcdr/2015/11984.6626>.

- [8] Smiline G, Hariprasad P, Pandi S, Raguraman R. A preliminary study on the screening of emerging drug resistance among the caries pathogens isolated from carious dentine. *Indian Journal of Dental Research* 2012;23:26. <https://doi.org/10.4103/0970-9290.99033>.
- [9] Balakrishnan M, Simmonds RS, Tagg JR. Dental caries is a preventable infectious disease. *Australian Dental Journal* 2000;45:235–45. <https://doi.org/10.1111/j.1834-7819.2000.tb00257.x>.
- [10] Kaur R, Kataria H, Kumar S, Kaur G. Caries Experience among Females aged 16–21 in Punjab, India and its Relationship with Lifestyle and Salivary HSP70 Levels. *European Journal of Dentistry* 2010;04:308–13. <https://doi.org/10.1055/s-0039-1697844>.
- [11] S S, Shetty S. Comparative Evaluation of the Effect of a Herbal Dentifrice and a Regular Dentifrice on Beneficial Oral Microflora – A Clinico Microbiological Study. *International Journal of Dentistry and Oral Health* 2016;1. <https://doi.org/10.16966/2378-7090.117>.
- [12] Menendez A, Li F, Michalek SM, Kirk K, Makhija SK, Childers NK. Comparative analysis of the antibacterial effects of combined mouthrinses on *Streptococcus mutans*. *Oral Microbiology and Immunology* 2005;20:31–4. <https://doi.org/10.1111/j.1399-302x.2004.00189.x>.
- [13] Ozaki F, Pannuti CM, Imbronito AV, Pessotti W, Saraiva L, de Freitas NM, et al. Efficacy of a herbal toothpaste on patients with established gingivitis: a randomized controlled trial. *Brazilian Oral Research* 2006;20:172–7. <https://doi.org/10.1590/s1806-83242006000200015>.
- [14] Lee SS, Zhang WU, Li Y. The antimicrobial potential of 14 natural herbal dentifrices. *The Journal of the American Dental Association* 2004;135:1133–41.

<https://doi.org/10.14219/jada.archive.2004.0372>.

[15] Balouiri M, Sadiki M, Bensouda SK. Methods for in vitro evaluating antimicrobial activity: A review. *Journal of Pharmaceutical Analysis* 2016;6:71–9.

<https://doi.org/10.1016/j.jpha.2015.11.005>.

[16] Prabakar J, John J, Arumugham IM, Kumar RP, Sakthi DS. Comparing the Effectiveness of Probiotic, Green Tea, and Chlorhexidine- and Fluoride-containing Dentifrices on Oral Microbial Flora: A Double-blind, Randomized Clinical Trial. *Contemp Clin Dent* 2018;9:560–9.

[17] Prabakar J, John J, Arumugham IM, Kumar RP, Sakthi DS. Comparative Evaluation of the Viscosity and Length of Resin Tags of Conventional and Hydrophilic Pit and Fissure Sealants on Permanent Molars: An Study. *Contemp Clin Dent* 2018;9:388–94.

[18] Prabakar J, John J, Arumugham IM, Kumar RP, Srisakthi D. Comparative Evaluation of Retention, Cariostatic Effect and Discoloration of Conventional and Hydrophilic Sealants - A Single Blinded Randomized Split Mouth Clinical Trial. *Contemp Clin Dent* 2018;9:S233–9.

[19] Shenoy RP, Salam TAA, Varghese S. Prevalence and Clinical Parameters of Cervical Abrasion as a Function of Population, Age, Gender, and Toothbrushing Habits: A Systematic Review. *World Journal of Dentistry* 2019;10:470–80.

[20] Manchery N, John J, Nagappan N, Subbiah G, Premnath P. Remineralization potential of dentifrice containing nanohydroxyapatite on artificial carious lesions of enamel: A comparative in vitro study. *Dent Res J* 2019;16:310.

[21] Vishnu Prasad S, Kumar M, Ramakrishnan M, Ravikumar D. Report on oral health status and treatment needs of 5-15 years old children with sensory deficits in

Chennai, India. *Spec Care Dentist* 2018;38:58–9.

[22] Khatri SG, Madan KA, Srinivasan SR, Acharya S. Retention of moisture-tolerant fluoride-releasing sealant and amorphous calcium phosphate-containing sealant in 6-9-year-old children: A randomized controlled trial. *J Indian Soc Pedod Prev Dent* 2019;37:92–8.

[23] Prabakar J, John J, Srisakthi D. Prevalence of dental caries and treatment needs among school going children of Chandigarh. *Indian Journal of Dental Research* 2016;27:547. <https://doi.org/10.4103/0970-9290.195683>.

[24] Samuel SR, Acharya S, Rao JC. School Interventions-based Prevention of Early-Childhood Caries among 3-5-year-old children from very low socioeconomic status: Two-year randomized trial. *J Public Health Dent* 2020;80:51–60.

[25] Mohapatra S, Pradeep Kumar R, Meignana Arumugham I, Sri Sakthi D, Jayashri P. Assessment of Microhardness of Enamel Carious Like Lesions After Treatment with Nova Min, Bio Min and Remin Pro Containing Toothpastes: An in Vitro Study. *Indian Journal of Public Health Research & Development* 2019;10:375. <https://doi.org/10.5958/0976-5506.2019.02832.8>.

[26] Pratha AA, Ashwatha Pratha A, Prabakar J. Comparing the effect of Carbonated and energy drinks on salivary pH- In Vivo Randomized Controlled Trial. *Research Journal of Pharmacy and Technology* 2019;12:4699. <https://doi.org/10.5958/0974-360x.2019.00809.6>.

[27] Kumar RP, Pradeep Kumar R, Vijayalakshmi B. Assessment of Fluoride Concentration in Ground Water in Madurai District, Tamil Nadu, India. *Research Journal of Pharmacy and Technology* 2017;10:309. <https://doi.org/10.5958/0974-360x.2017.00063.4>.

[28] Kumar RP, Pradeep Kumar R, Preethi R. Assessment of Water Quality and

Pollution of Porur, Chembarambakkam and Puzhal Lake. *Research Journal of Pharmacy and Technology* 2017;10:2157. <https://doi.org/10.5958/0974-360x.2017.00380.8>.

[29] Neralla M, Jayabalan J, George R, Rajan J, P SKM, Haque AE, et al. Role of nutrition in rehabilitation of patients following surgery for oral squamous cell carcinoma. *International Journal of Research in Pharmaceutical Sciences* 2019;10:3197–203. <https://doi.org/10.26452/ijrps.v10i4.1622>.

[30] Harini G, Leelavathi L. Nicotine Replacement Therapy for Smoking Cessation- An Overview. *Indian Journal of Public Health Research & Development* 2019;10:3588. <https://doi.org/10.5958/0976-5506.2019.04144.5>.

[31] Pavithra RP, Preethi Pavithra R, Jayashri P. Influence of Naturally Occurring Phytochemicals on Oral Health. *Research Journal of Pharmacy and Technology* 2019;12:3979. <https://doi.org/10.5958/0974-360x.2019.00685.1>.

[32] Kannan SSD, Kumar VS, Rathinavelu PK, Indiran MA. AWARENESS AND ATTITUDE TOWARDS MASS DISASTER AND ITS MANAGEMENT AMONG HOUSE SURGEONS IN A DENTAL COLLEGE AND HOSPITAL IN CHENNAI, INDIA. *Disaster Management and Human Health Risk V* 2017. <https://doi.org/10.2495/dman170121>.

[33] Mathew MG, Samuel SR, Soni AJ, Roopa KB. Evaluation of adhesion of *Streptococcus mutans*, plaque accumulation on zirconia and stainless steel crowns, and surrounding gingival inflammation in primary molars: randomized controlled trial. *Clin Oral Investig* 2020. <https://doi.org/10.1007/s00784-020-03204-9>.

[34] Katirae F. In vitro antifungal activity of essential oils against fluconazole-susceptible and-resistant *Candida albicans* n.d.

<https://doi.org/10.26226/morressier.5ac39995d462b8028d899f6c>.

- [35] Frawley D, Ranade S. Ayurveda, Nature's Medicine. Motilal Banarsidass Publ.; 2004.
- [36] Pereira WD, Geetha RV, Thangavelu L. Anti-inflammatory activity of Punica granatum extract on oral microbes - In vitro. International Journal of Research in Pharmaceutical Sciences 2019;10:1019–22. <https://doi.org/10.26452/ijrps.v10i2.375>.
- [37] Mukunda DA, Dept of Oral & Maxillofacial Pathology, Royal Dental College, Chalissery, Kerala, India. Efficacy of Psidium guajava leaf extract on Streptococcus mutans and Enterococcus faecalis – an in vitro study. Journal of Medical Science And Clinical Research 2019;7. <https://doi.org/10.18535/jmscr/v7i5.121>.
- [38] Kamal R, Dahiya P, Kumar M, Tomar V. Probiotics in oral health – A new tool in pharmaceutical science. Indian Journal of Pharmaceutical and Biological Research 2013;1:168–73. <https://doi.org/10.30750/ijpbr.1.4.27>.
- [39] Shaikh S, Manoj Kumar S. Beneficial effects of specific natural substances on oral health. Saudi Medical Journal 2017;38:1181–9. <https://doi.org/10.15537/smj.2017.12.20516>.
- [40] Gopinath NM, John J, Senthilkumar E, Nagappan N. Knowledge Awareness and Attitude about Research Ethics among Dental Faculties in India. The Journal of Contemporary Dental Practice 2014;15:608–13. <https://doi.org/10.5005/jp-journals-10024-1587>.
- [41] Shaheen SS, Sabiha Shaheen S. Antimicrobial Efficacy of Ten Commercially Available Herbal Dentifrices against Specific Oral Microflora – In Vitro Study. JOURNAL OF CLINICAL AND DIAGNOSTIC RESEARCH 2015.

<https://doi.org/10.7860/jcdr/2015/11127.5829>.

[42] Hakeem KR, Abdul WM, Hussain MM, Razvi SSI. Oral Health Care Products Obtained from Medicinal Plants. *Oral Health and Herbal Medicine* 2019;29–31.

https://doi.org/10.1007/978-3-030-04336-0_6.

[43] Okpalugo J, Ibrahim K, Inyang US. Toothpaste formulation efficacy in reducing oral flora. *Tropical Journal of Pharmaceutical Research* 2009;8.

<https://doi.org/10.4314/tjpr.v8i1.14714>.

[44] Singh R, Sharma A. Medicinal Plants Used for Diarrhoea by Tribals from Majhgawan Block of District Satna, Madhya Pradesh, India. *Studies on Ethno-Medicine* 2011;5:205–8. <https://doi.org/10.1080/09735070.2011.11886410>.

UNDER PEER REVIEW




Figure 1: Zone of inhibition of various types of herbal dentifrices against *Streptococcus mutans*. Zone of inhibition is found to be higher in Ayush (30 mm), followed by Dant kanti, Colgate vedshakti, Meswak, Dantajeevan and Dabur red showing inhibition zones of 26 mm, 24 mm, 17 mm and 14 mm respectively. Only Dantajeevan shows inhibition at 1:10 and 1:15 dilutions respectively.



Figure 2: Zone of inhibition of various types of herbal dentifrices against *Candida albicans*.

Zone of inhibition is found to be higher in Ayush and Dant kanti with 18 mm zone of inhibition followed by Colgate vedshakti, Meswak, Dantajeevan and Dabur red showing inhibition zones of 12mm and 10 mm respectively. Only Dant kanti shows inhibition at 1:10 dilution respectively.

Table 1 depicts the zone of inhibition of various types of herbal dentifrices against *Streptococcus mutans*. Zone of inhibition is found to be higher in Ayush (30 mm), followed by Dant kanti, Colgate vedshakti, Meswak, Dantajeevan and Dabur red showing inhibition zones of 26 mm, 24 mm, 17 mm and 14 mm respectively. Only Dantajeevan shows inhibition at 1:10 and 1:15 dilutions respectively.

S.no	Dentifrices	Zones of inhibition(mm)		
		1:5 dilution	1:10 dilution	1:15 dilution
1.	Dantajeevan	14	12	10
2.	Dant kanti	26	Nil	Nil
3.	Colgate cibaca vedshakti	24	Nil	Nil
4.	Meswak	17	Nil	Nil
5.	Ayush	30	Nil	Nil

6.	Dabur red	14	Nil	Nil
----	-----------	----	-----	-----

UNDER PEER REVIEW

S.no	Dentifrices	Zones of inhibition(mm)		
		1:5 dilution	1:10 dilution	1:15 dilution
1.	Dantajeevan	10	Nil	Nil
2.	Dant kanti	18	8	Nil
3.	Colgate cibaca vedshakti	12	Nil	Nil
4.	Meswak	12	Nil	Nil
5.	Ayush	18	Nil	Nil
6.	Dabur red	10	Nil	Nil

Table 2 depicts the zone of inhibition of various types of herbal dentifrices against *Candida albicans*. Zone of inhibition is found to be higher in Ayush and Dant kanti with 18 mm zone of inhibition followed by Colgate vedshakti, Meswak, Dantajeevan and Dabur red showing inhibition zones of 12mm and 10 mm respectively. Only Dant kanti shows inhibition at 1:10 dilution respectively.

UNDER PEER REVIEW

UNDER PEER REVIEW