

Original Research Article

Elevated red blood cell counts are associated with hormonal contraceptive usage among women in the Cape Coast Metropolis, Ghana

I suggest this title

hematological influence of the prolonged use of contraceptive methods in Ghanaian women: case of those in Cape Metropolis,

Formatted: Font: 18 pt, Bold, Not Italic, Not Superscript/ Subscript

Formatted: Centered

Formatted: Font: 18 pt, Bold, Not Italic, Not Superscript/ Subscript

Formatted: Font: 18 pt, Bold, Not Italic

Formatted: Font: 18 pt, Bold

Formatted: Font: 18 pt, Bold

Formatted: Font: 18 pt, Bold

UNDER PEER REVIEW

ABSTRACT

Background: The widespread use of hormonal contraceptives gives grounds for assessing their influence on various biochemical parameters of the human system since its safety has become controversial. This study assessed ~~the~~ effects of hormonal contraceptives on ~~haematological~~~~hematological~~ parameters among users in the Cape Coast Metropolis.

Methods and Material: A simple randomized case-control study approach was used to recruit 94 healthy women of which 54 were hormonal contraceptive users and 40 healthy age-matched non-contraceptive users served as controls. Venous blood samples were taken for full blood count (FBC) analyses using an automated ~~haematology~~~~hematological~~ ~~analyser~~~~analyzer~~. Data was ~~analysed~~~~analyzed~~ using SPSS (V.22.0). Data was expressed in means (Mean \pm SD) for the different variables. T-test statistic was used to compare the mean scores of two groups whilst one-way ANOVA was used to compare more than two groups. Pearson correlation was used to determine association between the various parameters. *P*-value <0.05 was considered statistically significant.

Results: The study observed mean red blood cell (RBC) count to be significantly higher among hormonal contraceptive users compared to non-users ($P= 0.030$). Additionally, the duration of contraceptive usage had an influence on the blood cell parameters in various ways, with a significant negative correlation between duration of contraceptive use and red cell distribution width (RDW) ($r= -0.303$, $P= 0.026$).

Conclusion and Recommendation: Hormonal contraceptives cause significant increase in red blood cell (RBC) count among users. Its effects also depend largely on duration of contraceptive use. Additionally, the main limitation to this study was the fact that blood films were not prepared to actually confirm the results presented by the ~~haematology~~~~hematological~~ ~~analyser~~~~analyzer~~, hence further study is recommended with the incorporation of blood films to the test panel.

Warning!!! A case-control study is a retrospective observational statistical study

this type of study alone cannot prove a causal link between a factor and the variation observed

Keywords: Hormonal contraceptives, haematological parameters, full blood count.

1. INTRODUCTION

Hormonal contraceptives are birth control methods that influence the endocrine system. Their purpose is to block ovulation by inhibiting the secretion of follicle stimulating hormone and luteinizing hormone. They also thicken the cervical mucus, making it a barrier to sperm transport, thereby blocking fertilization.^[1]

Hormonal contraceptives may comprise a combination of estrogen and progesterone or progesterone-only and can as well be administered via different routes (oral, intramuscular, vaginal, transdermal implants and associated with the intra-uterine device).

Over the years, there has been growing interests in the safety of hormonal contraceptives among users.

More concern has been about the impact of ~~oestrogen~~~~estrogen~~ and progesterone components of hormonal contraceptives on various biochemical and physiological processes in the bodies of users.^[2]

Available information indicates that hormonal contraceptives are associated with alterations in some trace elements and vitamins which influence blood production in humans.^[3-7] Further, some studies have

reported various haematological changes associated with the use of hormonal contraceptives.^[2, 8, 9] These changes include high platelet count which increases the risk of thromboembolism, myocardial infarction, arterial disease and carcinogenicity.^[10-17] These side effects may range from mild to severe but without a doubt may greatly influence the health status of users.^[18] The potential risk these types of contraceptives pose on the health of users raises some degree of health concern. Since the effects of these contraceptives vary from population to population, continual research is very vital. We examined the effect of hormonal contraceptive use on the haematological profile of females in the Cape Coast metropolis.

[English should be revised](#)

[material and methods](#) **SUBJECTS AND METHODS:**

STUDY DESIGN/STUDY SITE

This simple ~~randomised~~randomized case-control study was carried out within Cape Coast Metropolis in the Central Region of Ghana. Participants were recruited from the family planning units of the University of Cape Coast Hospital, Planned Parenthood Association of Ghana (PPAG) Clinic and Cape Coast Metropolitan ~~Hospital, all within the Cape Coast Metropolis. Laboratory analyses of samples were done at the University of Cape Coast Hospital laboratory in Cape Coast. Cape Coast is the administrative capital of Cape Coast Metropolitan Assembly. It forms part of the twenty (20) Metropolitan, Municipal and District Assemblies in the Central Region of Ghana. The Cape Coast Metropolis is bounded on the south by the Gulf of Guinea, west by Komenda/ Edina/ Eguafo/ Abrem Municipal assembly, east by the Abura/ Asebu/ Kwamankese District and north by the Twifu/ Hemang/ Lower Denkyira District. The Metropolis covers an area of 122 square kilometres and it is the smallest metropolis in the country~~^[19]

[English should seriously be revised](#)

STUDY POPULATION/ ELIGIBILITY CRITERIA

~~Ninety-four~~Ninety-four (94) non pregnant, non-lactating healthy women within the age range of 16-45 years comprising 54 on hormonal contraceptives (cases) and 40 women not on any hormonal contraceptives (controls) were recruited for the study. The fifty-four (54) women on hormonal contraceptives comprised 36 injectable Depo provera users, 11 Jadelle implant users and 7 combined oral

Formatted: Font color: Auto

Formatted: Font color: Red

contraceptives pills users. Apparently healthy women with no known haematological conditions were recruited as controls. Particularly, only those who had used hormonal contraceptives for at least six months were included as cases. The following categories of women were excluded; women with any known haematological disorder, menopausal and post-menopausal women, pregnant women, lactating mothers, those menstruating, those who were not within the age range of interest, and women who had used hormonal contraceptives for less than six months.

ETHICAL CONSIDERATION

Ethical clearance was obtained from the Institutional Review Board of the University of Cape Coast (IRB/UCC) concerning this research and approval was sought from the authorities of the various health facilities that were involved in this research. Informed consent was obtained from of all participants by enlightening them on the nature and relevance of the study. They were also informed of their right to participate or not in the study.

[Present the registration number and any proof of this ethical aspect](#)

BLOOD SAMPLE COLLECTION

With the aid of syringe and needle, three millilitres (3ml) of venous blood was taken from each participant into labelled tri-potassium ethylene diamine tetra-acetic acid (K₃EDTA) tubes according to standard protocol.^[20]

HAEMATOLOGICAL TEST

Full blood count (red blood cells, haemoglobin, haematocrit, mean corpuscular volume, mean corpuscular haemoglobin concentration, mean corpuscular haemoglobin, red cell distribution width, platelet count, white blood cell, lymphocytes, monocytes, neutrophils, eosinophils and basophils) analysis was done on each blood sample using automated haematology [analyseranalyzer](#) according to the principles of operation of the [analyseranalyzer](#) and their respective results were recorded. The haematology [analyseranalyzer](#) used was ABX Pentra 60 (Horiba ABX Diagnostics, France).

DATA ANALYSIS

Data was entered into Microsoft Excel (v. 15.0) and ~~analysed~~analyzed with SPSS (V.22.0). Data was expressed in means (Mean \pm SD) and percentages for the different variables. T-test statistic was used to compare the mean scores of two groups whilst one-way ANOVA was used to compare more than two groups. Additionally, Pearson correlation was used to determine association between the various parameters. P -value <0.05 was considered statistically significant.

RESULTS

[You should present socio-demographic data in order to better appreciate your sample
Not only ages and the marital status](#)

Table 1 shows the general characteristics of participants and compares the haematological parameters of study participants. Additionally, red blood cells (RBC) ($P=0.030$) was particularly higher in hormonal contraceptive (HC) users compared to non-users.

Table 2 compares haematological parameters of the various hormonal contraceptive users (injectable Depo provera, combined oral contraceptive pills and Jadelle implant). The study observed that in spite of fluctuations in the blood parameters among the various users, there was generally no significant difference in haematological parameters among users of the various types of hormonal contraceptives ($P>0.05$).

Table 3 compares haematological parameters of different groups of hormonal contraceptive users based on duration of use. MCV and MCH were higher among women who had used HC for more than 2 years compared to those who had used HC for 2 years or less. The major difference in these parameters (MCV and MCH) were observed between those who had used for 2 years or less and those who had used for 3 to 6 years were statistically significant ($P=0.031$ and $P=0.033$ respectively). RDW was higher among those who had used for 2 years or less compared to those who had used for higher years, with the difference between those had used for 2 years or less and those who had used for 3 to 6 years showing statistical significance ($P=0.011$).

Table 4 compares haematological parameters of hormonal contraceptive users according to their age groups. Monocyte count was significantly higher ($P=0.013$) among contraceptive users at the age of 40 or above as compared to those within the age range of 25 to 29.

Table 5 shows correlation between duration of contraceptive use and haematological parameters among HC users. There was a significant negative correlation between duration of contraceptive use and RDW ($r = -0.303$, $P = 0.026$).

[Results will be more relevant if you calculate the odd ratio](#)

Table 1: Demographic and haematological profile of study participants

Variable	Cases (n = 54)	Controls (n = 40)	P-value
Age (years)	30.04 ± 5.76	29.18 ± 8.15	0.549
Marital status			
Single	21 (38.9)	20 (50.0)	0.283
Married	33 (61.1)	20 (50.0)	
Haematological profile			
RBC × 10¹²/mm³	4.35 ± 0.48	4.14 ± 0.42	0.030
HGB (g/dl)	12.32 ± 0.81	11.93 ± 1.20	0.061
HCT (%)	37.64 ± 2.46	36.52 ± 3.56	0.075
MCV (fl)	87.04 ± 5.92	88.03 ± 4.56	0.382
MCH (pg)	28.73 ± 2.16	28.47 ± 1.76	0.534

MCHC (g/dl)	32.71 ± 0.90	32.55 ± 0.93	0.382
RDW (%)	13.05 ± 1.35	12.89 ± 1.61	0.597
PLT ×10 ⁹ /l	206.28 ± 39.08	196.65 ± 35.12	0.221
WBC ×10 ⁹ /l	6.07 ± 1.29	6.19 ± 1.33	0.651
LYM (%)	32.69 ± 3.83	32.81 ± 4.11	0.880
MON (%)	8.04 ± 1.63	8.30 ± 1.82	0.464
NEU (%)	56.12 ± 3.70	55.67 ± 4.14	0.579
EOS (%)	2.35 ± 1.10	2.16 ± 0.83	0.362
BAS (%)	0.91 ± 0.25	1.00 ± 0.34	0.126

RBC= Red blood cells, HGB= Haemoglobin, HCT= Haematocrit, MCV= Mean corpuscular volume, MCH= Mean corpuscular haemoglobin, MCHC= Mean corpuscular haemoglobin concentration, RDW= Red cell distribution width, PLT= Platelets, WBC= White blood cells, LYM= Lymphocytes, MON= Monocytes, NEU= Neutrophils, EOS= Eosinophils, BAS= Basophils

Table 2: Haematological profile of hormonal contraceptive users based on the types of contraceptives

Parameter	Type of HC			P-value
	Depo Provera (n = 36)	COCP (n = 7)	Jadelle (n = 11)	
RBC × 10 ¹² /mm ³	4.41 ± 0.47	4.25 ± 0.40	4.18 ± 0.51	0.297
HGB (g/dl)	12.44 ± 0.79	12.01 ± 0.74	12.10 ± 0.91	0.274
HCT (%)	38.07 ± 2.50	36.59 ± 2.07	36.92 ± 2.38	0.192
MCV (fl)	86.58 ± 5.74	86.57 ± 7.30	88.82 ± 5.83	0.544
MCH (pg)	28.28 ± 2.19	28.57 ± 2.67	29.03 ± 1.78	0.604
MCHC (g/dl)	32.69 ± 1.01	32.83 ± 0.35	32.72 ± 0.81	0.935
RDW (%)	13.24 ± 1.35	13.37 ± 1.29	12.22 ± 1.16	0.070
PLT ×10 ⁹ /l	203.72 ± 41.07	214.86 ± 49.22	209.18 ± 25.59	0.766
WBC ×10 ⁹ /l	5.91 ± 1.19	5.89 ± 0.97	6.70 ± 1.68	0.195
LYM (%)	33.36 ± 4.03	31.66 ± 3.09	31.14 ± 3.20	0.183
MON (%)	7.92 ± 1.27	8.56 ± 1.82	8.09 ± 2.49	0.642
NEU (%)	55.55 ± 3.91	57.04 ± 4.40	57.42 ± 1.92	0.269
EOS (%)	2.39 ± 1.20	1.90 ± 0.59	2.51 ± 1.01	0.494
BAS (%)	0.91 ± 0.27	0.84 ± 0.21	0.94 ± 0.23	0.750

HC= Hormonal contraceptive, COCP= Combined oral contraceptive pills

Table 3: Haematological profile of hormonal contraceptive users stratified by the duration of contraceptive use

Parameter	Duration of use (years)			P-value
	≤ 2 (n = 30)	3-6 (n= 21)	≥ 6 (n = 3)	
RBC × 10 ¹² /mm ³	4.45 ± 0.48	4.41 ± 0.46	4.20 ± 0.48	0.176
HGB (g/dl)	12.27 ± 0.88	12.30 ± 0.73	12.93 ± 0.71	0.411
HCT (%)	37.71 ± 2.83	37.32 ± 1.86	39.27 ± 2.25	0.438
MCV (fl)	85.17 ± 5.01	89.43 ± 6.52*	89.00 ± 4.58	0.031
MCH (pg)	27.79 ± 1.97	29.33 ± 2.26*	29.20 ± 0.98	0.033
MCHC (g/dl)	32.53 ± 1.05	32.94 ± 0.66	32.93 ± 0.31	0.269
RDW (%)	13.52 ± 1.34	12.52 ± 1.20*	12.00 ± 0.36	0.011
PLT ×10 ⁹ /l	200.70 ± 41.22	213.00 ± 34.48	215.00 ± 53.84	0.510
WBC ×10 ⁹ /l	5.93 ± 1.24	6.27 ± 1.41	6.03 ± 1.31	0.671
LYM (%)	32.13 ± 3.97	33.01 ± 3.68	35.90 ± 2.15	0.240
MON (%)	7.95 ± 1.86	8.25 ± 1.31	7.43 ± 1.34	0.655
NEU (%)	56.72 ± 3.54	55.67 ± 3.80	53.27 ± 4.05	0.239
EOS (%)	2.30 ± 1.08	2.24 ± 0.91	3.63 ± 2.06	0.112
BAS (%)	0.93 ± 0.28	0.89 ± 0.23	0.77 ± 0.06	0.543

P < 0.05, *significantly different from groups on contraceptive for ≤ 2 years

UNACCEPTED FOR PUBLICATION
 REVIEW


Table 4: Haematological profile of hormonal contraceptive users stratified by age groups

Parameter	Age group					P-value
	<25 (n = 10)	25-29 (n = 17)	30-34 (n = 13)	35-39 (n = 12)	≥ 40 (n = 2)	
RBC ×10 ¹² /mm ³	4.33 ± 0.33	4.47 ± 0.42	4.33 ± 0.59	4.16 ± 0.31	4.65 ± 1.40	0.412
HGB (g/dl)	12.28 ± 0.82	12.46 ± 0.84	12.27 ± 1.00	12.16 ± 0.62	12.80 ± 0.85	0.824
HCT (%)	37.33 ± 2.30	37.99 ± 2.29	37.28 ± 2.82	37.22 ± 1.55	41.10 ± 6.08	0.283
MCV (fl)	86.60 ± 4.45	85.12 ± 4.27	86.92 ± 7.99	89.67 ± 4.94	90.50 ± 13.44	0.303
MCH (pg)	28.53 ± 1.25	27.85 ± 1.73	28.32 ± 2.71	29.38 ± 1.67	28.80 ± 7.07	0.469
MCHC (g/dl)	32.84 ± 0.55	32.74 ± 1.14	32.84 ± 0.45	32.66 ± 0.77	31.35 ± 2.62	0.287
RDW (%)	12.94 ± 1.16	13.54 ± 1.40	13.17 ± 1.24	12.28 ± 1.25	13.25 ± 2.62	0.177
PLT ×10 ⁹ /l	198.60 ± 46.64	201.29 ± 35.10	217.23 ± 41.41	206.42 ± 30.02	215.00 ± 91.92	0.789
WBC ×10 ⁹ /l	5.70 ± 1.06	6.19 ± 1.64	6.04 ± 1.05	6.05 ± 1.26	7.15 ± 1.06	0.682
LYM (%)	31.14 ± 3.22	32.86 ± 4.04	32.35 ± 4.84	33.62 ± 2.76	35.45 ± 2.76	0.489
MON (%)	8.49 ± 1.44	7.21 ± 1.07	8.35 ± 2.15	8.02 ± 1.09	10.90 ± 1.98*	0.013
NEU (%)	56.72 ± 2.71	56.98 ± 3.86	55.81 ± 4.27	55.50 ± 3.35	51.60 ± 3.82	0.337
EOS (%)	2.86 ± 1.37	2.19 ± 1.00	2.52 ± 1.36	2.14 ± 0.49	1.30 ± 0.71	0.283
BAS (%)	0.85 ± 0.18	0.85 ± 0.22	0.98 ± 0.32	0.98 ± 0.26	0.75 ± 0.21	0.376

P < 0.05, *significantly different from age group (25-29) years

DU= Duration of contraceptive use, PMT=Parameter

UNDER PEER REVIEW



3.2 DISCUSSION

This study assessed the effect of hormonal contraceptives (HC) on full blood count parameters (RBC, HGB, HCT, MCV, MCH, MCHC, RDW, platelets, WBC, lymphocytes, monocytes, neutrophils, eosinophils and basophils) among women within the Cape Coast Metropolis. Our study observed a significantly elevated RBC count among HC users compared to non-users. This finding is however contrary to that of similar studies conducted in Pakistan and Gaza.^[2, 9] Our observation may be due to reduced menstrual blood flow which is known to be common among HC users.^[2, 3]

Additionally, slight increases in RDW were observed among HC users compared to non-users. This result is however contrary to that reported in Gaza.^[18] In their study, they found significantly lower RDW among users of progestogen-only contraceptives compared to controls in Gaza city. The variation in results may be due to oestrogenic effect of the combined oestrogen-progestogen contraceptives included in this study. This may account for RDW values in this present study being slightly higher among combined oestrogen-progestogen users compared to the progestogen-only contraceptives users.

Further, duration of contraceptive use was seen to influence MCV, MCH and RDW. There was a significant increase in MCV and MCH, and a significant decrease in RDW among those who had used HC for 3 to 6 years compared to those who had use for 2 years or less. This was further seen in correlation analysis where there was a significant negative correlation between duration of contraceptive use and RDW. This shows a duration-dependent effect of HC on RDW. From the MCV results, it is possible that long-term use of these contraceptives may have macrocytic effect on the red blood cells, perhaps when the negative effect HC on vitamin B12 and folate suggested by some studies begins to manifest.^[4-6]

Lastly, HC users at the age of 40 and above had significantly higher monocyte count compared to those within the ages of 25 to 29 years. The reason for this observation is however not immediately apparent.

CONCLUSION AND RECOMMENDATION

Hormonal contraceptives cause significant increase in red blood cell (RBC) count among users. Its effects also depend largely on duration of contraceptive use. Additionally, the main limitation to this study was the fact that blood films were not prepared to actually confirm the results presented by the haematology [analyser/analyzer](#), hence further study is recommended with the incorporation of blood films to the test panel.

ACKNOWLEDGEMENTS

A brief acknowledgement section may be given after the conclusion section just before the references. The acknowledgments of people who provided assistance in manuscript preparation, funding for research, etc. should be listed in this section. All sources of funding should be declared as an acknowledgement. Authors should declare the role of funding agency, if any, in the study design, collection, analysis and interpretation of data; in the writing of the manuscript. If the study sponsors had no such involvement, the authors should so state.

COMPETING INTERESTS

Declaration of competing interest should be placed here. All authors must disclose any financial and personal relationships with other people or organizations that could inappropriately influence (bias) their work. Examples of potential conflicts of interest include employment, consultancies, honoraria, paid expert testimony, patent applications/registrations, and grants or other funding. If no such declaration has been made by the authors, SDI reserves to assume and write this sentence: "Authors have declared that no competing interests exist."

CONSENT (WHERE EVER APPLICABLE)

Written informed consent was obtained from participants before the study took place and these documents are preserved by the authors.

ETHICAL APPROVAL (WHERE EVER APPLICABLE)

Ethical clearance was sought from the Institutional Review Board of the University of Cape Coast (IRB/UCC) and administrators of the Hospital before the study was commenced. In addition, all women recruited into the study gave informed consent after thorough explanation of the rationale of the study has been given.

REFERENCES

REFERENCES SHOULD BE REVISED

SOME ARE VERY OLD

USE REFERENCES UNDER 6 YEARS OLD

1. Brito BM, Nobre F, Vieira SC. Contracepção hormonal e sistema cardiovascular. Arq Bras Cardiol, 2011. **96**(4): p. 81-9.
2. Afsar AN, Barakzai Q, Adil NS. Effect Of A 'Progestin Only' Contraceptive On Platelet Aggregation In A Pakistani Set Of Population. J Ayub Med Coll Abbottabad 2005. **17**(3).
3. Akinloye O, Adebayo TO, Oguntibeju OO, Oparinde DP, Ogunyemi EO. Effects of contraceptives on serum trace elements, calcium and phosphorus levels. West Indian Med J, 2011. **60**(3): p. 308-315.
4. Lussana F, Zighetti ML, Bucciarelli P, Cugno M, Cattaneo M. Blood levels of homocysteine, folate, vitamin B6 and B12 in women using oral contraceptives compared to non-users. Thromb Res 2003. **112**: p. 37-41.
5. Palmery M, Saraceno A, Vaiarelli A, Carlomagno G. Oral contraceptives and changes in nutritional requirements. European Review for Medical and Pharmacological Sciences, 2013. **17**: p. 1804-1813.

6. Sütterlin MW, Bussen SS, Rieger L, Dietl J, Steck T. Serum folate and Vitamin B12 levels in women using modern oral contraceptives (OC) containing 20 microg ethinyl estradiol. *Eur J Obstet Gynecol Reprod Biol*, 2003. **107**: p. 57-61.
7. Afrifa J, Essien-Baidoo S, Ofori EG, Abaka-Yawson A. Trace Elements Levels and Blood Pressure in Ghanaian Women Using Depot Medroxyprogesterone Acetate Contraceptive. *British Journal of Medicine and Medical Research*. 2017. **20**(11): p. 1-7.
8. Al-Chalaby SSH, Taib SM, Ahmed AF. The Effect of Oral Contraceptive Pills on Haematological Indices. *Tikrit Medical Journal*, 2006. **12**(1): p. 65-69.
9. Zabut B, Abu-Hani I. Evaluation of some Physiological and Biochemical Parameters of Healthy Women Administrated Oral Contraceptive Pills in Gaza City. *IUG Journal of Natural and Engineering Studies*, 2012. **20**(2): p. 29-44.
10. Bloemenkamp KW, Rosendaal FR, Büller HR, Helmerhorst FM, Colly LP, Vandenbroucke JP. Risk of venous thrombosis with use of current low-dose oral contraceptives is not explained by diagnostic suspicion and referral bias. *Archives of Internal Medicine*, 1999. **159**: p. 65-70.
11. Farmer RD, Lawrenson RA, Thompson CR, Kennedy JG, Hambleton IR. Population-based study of risk of venous thromboembolism associated with various oral contraceptives. *The Lancet*, 1997. **349**: p. 83-88.
12. Lidegaard Ø, Edström B, Kreiner S. Oral contraceptives and venous thromboembolism: A five-year national case-control study. *Contraception*, 2002. **65**: p. 187-196.
13. Parkin L, Skegg DC, Wilson M, Herbison GP, Paul C. Oral contraceptives and fatal pulmonary embolism. *The Lancet*, 2000. **355**: p. 2133-2134.
14. Poulter NR, Meirik O, Chang CL, Farley TM, Kelaghan J, Marmot MG, Debert-Ribeiro M, Medina E, Artigas J, He S, Hui ZY. Acute myocardial infarction and combined oral contraceptives: results of an international multicentre casecontrol study. *WHO Collaborative Study of Cardiovascular Disease and Steroid Hormone Contraception. Lancet*, 1997. **349**: p. 1202–1209.
15. Tanis BC, Rosendaal FR. Venous and arterial thrombosis during oral contraceptive use: risks and risk factors. *Semin Vasc Med*, 2003. **3**: p. 69–84.
16. Kay C. Incidence of arterial disease among oral contraceptive users. *Gen. Pract.*, 1983. **33**: p. 95-92.
17. Marchbanks PA, McDonald JA, Wilson HG, Folger SG, Mandel MG, Daling JR, Bernstein L, Malone KE, Ursin G, Strom BL, Norman SA. Oral contraceptives and

the risk of breast cancer. *New England Journal of Medicine*, 2002. **346**: p. 2025-2032.

18. Sabatini R, Cagiano R, Rabe T. Adverse Effects of Hormonal Contraception. *Journal of Reproductive Medicine and Endocrinology*, 2011. **8**(1): p. 130-156.
19. Ghanadistricts. 2006 [cited 2013 April 13]; Available from: www.ghanadistricts.com/districts/?news&r=3&_id=50.
20. Turgeon ML. *Clinical Hematology: Theory and Procedures*. 5 ed. 2012, Baltimore: Lippincott Williams & Wilkins, a Wolters Kluwer business.

UNDER PEER REVIEW