

## Original Research Article

### Assessment of Knowledge and Compliance with Standard Precautions in Selected Hospitals in Rivers State, Nigeria

#### Abstract

**Background:** Standard precautions (SPs) in the hospital are very important in the reduction of health care worker's exposure to occupational health hazards but most hospitals operate reactive rather than proactive safety practices. The effect of reactive safety practices is overwhelming to health care workers, hospitals and the society. Thus, this study assessed health care worker's knowledge and compliance with standard precautions. **Material and Methods:** Descriptive cross-sectional design and proportionate stratified random sampling method was adopted with a sample size of 391. The study relied on a structured questionnaire with a reliability of 0.87. Descriptive and inferential statistics were used with the aid of statistical package for social science (SPSS) software version 21. **Results:** HCWs have fair knowledge of SPs and poor compliance with SPs. Also, there is significant association between HCW's knowledge of SPs and gender, marital status, years of experience and job category in the hospitals ( $p < 0.05$ ). Consequently, there is significant association between compliance with SPs and gender, years of experience and job categories in the hospitals ( $p < 0.05$ ). **Conclusion:** There is great need for strong advocacy on knowledge of SPs and compliance with SPs in the hospitals to curb increasing occupational health hazards. The government and hospital managements should develop an all-encompassing integrated SPs policies, strategies and procedures consistent with global best practices.

**Key Word:** Knowledge, Compliance, Standard Precautions, personal protective equipment (PPEs), Hospitals

#### I. Introduction

There is absence of obligatory laws and regulations governing the labor environment and this makes a lot of workers susceptible to hazards which may be life threatening. In developing countries, occupational health and safety (OHS) regulations make up only 10% of the populace [1]. Therefore, abandoning quite a lot of key hazardous businesses and professions like the health sector is dangerous [2], [3]. Even where the laws exist, workers are still at risk of being hired into situations that deprive them of their ethics and self-respect and these exposes them to a great number of job-related events and accidents. This has led to worsening workers' health and welfare as some African studies opined that occupationally, health care workers (HCWs) obtained infections like tropical illnesses [2]. There is also the problem of lack of, or incomplete vaccination against hepatitis B virus (HBV) infection, inadequate knowledge of standard precautions (SPs), reduced availability/insufficient use of personal protective equipment (PPE), and absence of preventive training in hospitals which amounts to high possibility for hazard exposure to HCWs [4-7], [5], [6], [7]. These issues have been confirmed by some studies. Knowledge of SPs among HCWs is low. Only 37.0% of HCWs knew that SPs comprises hand washing before and after any contact with a patient, is a component of SPs. 39.0% of the HCWs

Comment [K1]: Workers'?

Formatted: Highlight

Comment [K2]: Workers'?

Comment [K3]: Was used to select 391 study sample?

Comment [K4]: Meaning what?

Comment [K5]: In full first.

Comment [K6]: HCWs'

Comment [K7]: Meaning?

Comment [K8]: Hanging sentence!!

recognized cough protocols and only 40.0% knew about sanitary procedures which comprise infection preventive approaches to curb the dangers of exposure to infection. 50% of the HCWs continually guard themselves against blood and & body fluids (BBFs) of patients, 25% do not recap needles after usage and 28.0% do not always wipe blood spills on time [8]. 63.0% of HCWs opined that using PPEs such as aprons, gloves, gowns and goggles, might cause patients to fright occasionally and adhering to SPs, sometimes obstructs the ability to offer care (38.0%).

**Comment [K9]:** Do not start a sentence with numbers!

**Comment [K10]:** Protect?

**Comment [K11]:** See K9.

This study adopts the Health Belief Model (HBM). It is a psychosocial model that tries to predict health behaviors portrayed by HCWs as they perform health care services. The HBM lays emphases on the attitudes, perception and knowledge of HCWs. It is centered on an understanding that a HCW will take a health-related action if he/she perceives that a health illness can be avoided, or has a rational anticipation that by carrying out a suggested action (compliance with SPs), he/she will avoid a hazard. Occupational exposure of HCWs and the susceptibility of diseases are on the rise. The continuous exposure to threats from infectious ailments has caused numerous HCWs to suffer from different illnesses contracted by handling of patients infected with various transmissible diseases. Likewise, lack of safety practices through the hospitals is a concern to HCWs, administrative bodies and the general public. This is unquestionably a crucial problem in the health sector. The long term effects comprise HCWs'/patients' susceptibility to infections, scarcity of health sector employees, HCWs' apathy and cost effects, even with the SPs scheme. Thus, there is plea for cautionary actions and reaction safety practices in hospitals. Therefore, the objectives of the study are to determine the level of knowledge of standard precautions and assess health care workers' compliance with standard precautions.

**Comment [K12]:** Provide?

**Comment [K13]:** In/within?

**Comment [K14]:** Were?

## II. Material And Methods

**Study Design:** The study employed the descriptive cross-sectional design.

**Study Location:** The study was conducted in the two tertiary in Rivers State, Southern Nigeria, namely Rivers State University Teaching Hospital (RSUTH) and University of Port Harcourt Teaching Hospital (UPTH), and the Zonal Hospitals including Okirika, Degema, Bonny, Isiokpo, Ahaoda and Bori hospitals respectively.

**Comment [K15]:** Hospitals/health facilities?

**Study Duration:** April, 2019-October, 2019.

**Sample Size:** 391 HCWs

**Sample Size Calculation:** proportionate stratified random sampling method with the use of table of random numbers was adopted for the sample selection. A population of 3876 staffs was considered, giving a sample size of 391, which includes 10% non-response rate was used by applying Taro-Yamane formula [9] given as:

$$n = \frac{N}{1 + N(e)^2} \quad (1)$$

where n is the sample size, e is tolerance error (0.05) and N is the study population.

Primary data were gathered using closed-ended structured questionnaire with closed ended which were self- and interviewer-administered, while secondary data was obtained via hospital records of the management of the hospitals.

**Comment [K16]:** Meaning?

**Inclusion Criteria:** The research was limited to direct staffs who have worked in the hospital facility for at least 6 months and above. This includes doctors, nurses, pharmacists, clinical laboratory attendants, mortuary attendants, cleaners/porters, administrative workers and radiographers. All participants are aged 20 to 60 years. Their typical work day starts from 8am to 4pm after which the call duty starts till 8am the following day.

**Comment [K17]:** Were?

**Exclusion Criteria:** HCWs who are on contract, brief internship training, students, those in maternity leave, as well as those who have worked for less than 6 months were excluded from the study.

**Procedure methodology:** The questionnaires copies were administered with the help of a research assistant. Knowledge of SPs score of 1-4 is rated poor, 4-7 is rated fair and 8-10 is rated good. Questions on compliance were structured using a 4-point likert scale.

**Comment [K18]:** Provide a brief description of this tool!

**Statistical analysis:** The data obtained were subjected to statistical analyses such as descriptive (mean, percentages and charts) and inferential (Chi-Square) statistics using statistical package for social sciences (SPSS) version 21. A test retest using Pearson Product Moment of Correlation Coefficient was carried out and a value of 0.87 was obtained. Content validation was done by experts in occupational health and safety.

### **Ethical Approval**

This study was approved by River State Hospital Management Board (RSHMB) and University of Port Harcourt Research Ethical Committee, Port Harcourt, Nigeria. Informed consent was obtained from respondents before they participated in the study. In the process, respondents were educated in a nut shell on hazards they may face as healthcare workers and how to mitigate them. Respondents were informed of their right to opt out of the study if necessary.

## **III. Results**

### **Knowledge of Standard Precautions (SPs)**

**Table 1** assesses the association between knowledge of SPs and social-demographic variables. Result revealed that 41.7% of respondents have good knowledge of SPs, while 34.3% and 24.0% have fair and poor knowledge of SPs respectively. The Chi-Square results showed that knowledge of SPs is associated with gender, marital status, years of experience and job categories of respondents ( $p < 0.05$ ). But knowledge of SPs is not associated with educational level and age of respondents ( $p > 0.05$ ).

**Comment [K19]:** Presents ?

Table 1: Association between Knowledge of SPs and Social-Demographic Variables

Variables	Good (%)	Fair (%)	Poor (%)	Total (%)	df.	Chi-Square (p-value)	Decision
<b>Gender</b>							
Male	98(48)	70(34.3)	36(17.6)	204(100)	2	11.381 (0.003)	Significant (p<0.05)
Female	65(34.8)	64(34.2)	58(31)	187(100)			
<b>Marital Status</b>							
Single	76(47.8)	56(35.2)	27(17)	159(100)	8	15.758 (0.046)	Significant (p<0.05)
Married	84(38.4)	74(33.8)	61(27.9)	219(100)			
Divorced	3(60)	1(20)	1(20)	5(100)			
Co-habiting	0(0)	2(40)	3(60)	5(100)			
Widow/Widower	0(0)	1(33.3)	2(66.7)	3(100)			
<b>Years of Experience</b>							
0-5yrs	5(14.7)	18(52.9)	11(32.4)	34(100)	10	20.855 (0.022)	Significant (p<0.05)
6-10yrs	66(47.8)	46(33.3)	26(18.8)	138(100)			
11-15yrs	46(37.1)	46(37.1)	32(25.8)	124(100)			
16-20yrs	30(50.8)	16(27.1)	13(22)	59(100)			
21-25yrs	9(45)	3(15)	8(40)	20(100)			
26+yrs and Above	7(43.8)	5(31.2)	4(25)	16(100)			
<b>Job Categories</b>							
Doctors	26(35.6)	31(42.5)	16(21.9)	73(100)	16	35.709 (0.003)	Significant (p<0.05)
Nurses	29(31.9)	34(37.4)	28(30.8)	91(100)			
Pharmacists	23(50)	16(34.8)	7(15.2)	46(100)			
Radiographers	19(61.3)	9(29)	3(9.7)	31(100)			
Administrators	30(45.5)	22(33.3)	14(21.2)	66(100)			
Cleaners/Porters	4(19)	7(33.3)	10(47.6)	21(100)			
Clinical Lab.	12(38.7)	7(22.6)	12(38.7)	31(100)			
Mortuary	9(60)	2(13.3)	4(26.7)	15(100)			
<b>Educational Status</b>							
Post Graduate	50(40.3)	45(36.3)	29(23.4)	124(100)	6	4.989 (0.545)	Significant (p>0.05)
Tertiary	85(43.6)	66(33.8)	44(22.6)	195(100)			
Secondary	27(42.9)	19(30.2)	17(27)	63(100)			
Primary	1(11.1)	4(44.4)	4(44.4)	9(100)			
<b>Age (Years)</b>							
<25 & Below	10(43.5)	8(34.8)	5(21.7)	23(100)	10	7.100 (0.716)	Significant (p>0.05)
26-35	82(46.9)	56(32)	37(21.1)	175(100)			
36-45	49(38.6)	44(34.6)	34(26.8)	127(100)			
46-55	20(36.4)	20(36.4)	15(27.3)	55(100)			
56-65	1(12.5)	5(62.5)	2(25)	8(100)			
66+ and Above	1(33.3)	1(33.3)	1(33.3)	3(100)			
Total	163(41.7)	134(34.3)	94(24)	391(100)			

Comment [K20]: Technicians?

Comment [K21]: Attendants?

## Compliance with Standard Precautions (SPs)

**Table 2** assesses the association between compliance with SPs and social-demographic variables. Results revealed that 14.85% of respondents always comply with SPs, while 21.0%, 44.0% and 20.2% of respondents often, sometimes and rarely comply with SPs respectively. The Chi-Square results showed that compliance with SPs is associated with gender, years of experience and job categories of respondents ( $p < 0.05$ ). But compliance with SPs is not associated with marital status, educational level and age of respondents ( $p > 0.05$ ).

**Comment [K22]:** In that order?


Consequently, Figure 1 revealed HCW's non-compliance with SPs for various job categories; the bar chart revealed that 85.8% of cleaners/porters do not comply with SPs followed by nurses (70.4%), mortuary attendants (66.7%), radiographers (64.5%) and administrators (62.1%) respectively. Also, clinical laboratory attendants (61.3%) followed by pharmacists (60.9%) and lastly doctors (60.2%).

**Table 2: Association between Compliance with SPs and Social-Demographic Variables**

**Comment [K23]:** See changes suggested in Table 1 and incorporate in this Table.

Variables	Always (%)	Often (%)	Sometimes (%)	Rarely (%)	Total (%)	df.	Chi-Square (p-value)	Decision
<b>Gender</b>								
Male	35(17.2)	50(24.5)	89(43.6)	30(14.7)	204(100)	3	10.494 (0.015)	Significant ( $p < 0.05$ )
Female	23(12.3)	32(17.1)	83(44.4)	49(26.2)	187(100)			
<b>Marital Status</b>								
Single	28(17.6)	38(23.9)	72(45.3)	21(13.2)	159(100)	12	17.521 (0.131)	Significant ( $p > 0.05$ )
Married	29(13.2)	43(19.6)	95(43.4)	52(23.7)	219(100)			
Divorced	1(20)	1(20)	2(40)	1(20)	5(100)			
Co-habiting	0(0)	0(0)	2(40)	3(60)	5(100)			
Widow/Widower	0(0)	0(0)	1(33.3)	2(66.7)	3(100)			
<b>Years of Experience</b>								
0-5yrs	2(5.9)	9(26.5)	13(38.2)	10(29.4)	34(100)	15	26.811 (0.030)	Significant ( $p < 0.05$ )
6-10yrs	26(18.8)	25(18.1)	64(46.4)	23(16.7)	138(100)			
11-15yrs	12(9.7)	35(28.2)	53(42.7)	24(19.4)	124(100)			
16-20yrs	13(22)	8(13.6)	26(44.1)	12(20.3)	59(100)			
21-25yrs	5(25)	3(15)	5(25)	7(35)	20(100)			
26yrs and Above	0(0)	2(12.5)	11(68.8)	3(18.8)	16(100)			
<b>Job Categories</b>								
Doctors	6(8.2)	23(31.5)	29(39.7)	15(20.5)	73(100)	24	65.631 (0.000)	Significant ( $p < 0.05$ )
Nurse	13(14.3)	14(15.4)	41(45.1)	23(25.3)	91(100)			
Pharmacist	17(37)	1(2.2)	21(45.7)	7(15.2)	46(100)			

Radiographer	6(19.4)	5(16.1)	17(54.8)	3(9.7)	31(100)			
Administration	10(15.2)	15(22.7)	27(40.9)	14(21.2)	66(100)			
Cleaners/Porters	1(4.8)	2(9.5)	9(42.9)	9(42.9)	21(100)			
Clinical Lab.	0(0)	12(38.7)	12(38.7)	7(22.6)	31(100)			
Mortuary	0(0)	5(33.3)	9(60)	1(6.7)	15(100)			
<b>Education</b>								
Post Graduate	23(18.5)	20(16.1)	55(44.4)	26(21)	124(100)	9	9.787 (0.368)	Significant (p>0.05)
Tertiary	29(14.9)	44(22.6)	86(44.1)	36(18.5)	195(100)			
Secondary	6(9.5)	17(27)	27(42.9)	13(20.6)	63(100)			
Primary	0(0)	1(11.1)	4(44.4)	4(44.4)	9(100)			
<b>Age (Years)</b>								
25 & Below	4(17.4)	5(21.7)	10(43.5)	4(17.4)	23(100)	15	12.234 (0.661)	Significant (p>0.05)
26-35	27(15.4)	46(26.3)	73(41.7)	29(16.6)	175(100)			
36-45	19(15)	21(16.5)	56(44.1)	31(24.4)	127(100)			
46-55	8(14.5)	8(14.5)	26(47.3)	13(23.6)	55(100)			
56-65	0(0)	1(12.5)	6(75)	1(12.5)	8(100)			
66 and Above	0(0)	1(33.3)	1(33.3)	1(33.3)	3(100)			
<b>Total</b>	58(14.8)	82(21)	172(44)	79(20.2)	391(100)			



**Figure 1: Bar Chart of HCWs Non-Compliance with SPs.**

#### IV. Discussion

This study revealed that HCWs have fair (76.0%) knowledge of standard precautions (SPs) in the hospitals. This is in agreement with [10] who opined that HCWs lack knowledge about blood borne pathogens (BBPs) transmission and SPs in China. [8] also reported HCWs had low level of knowledge of SPs among HCWs in Ghana. HCWs are not continually informed on SPs principles, insufficiencies or lack of knowledge and practice of SPs [11-13], [12], [13]. This study is at variance with [13] who stated that 89 % of HCWs (89%) had good knowledge about SPs in health care facilities (HCFs). Some other studies also demonstrated good knowledge of SPs at 86.9% and 81.8% respectively [14-15], [15]. Some scholars also argued that knowledge of SPs is vital to avert and regulate the susceptibility to hospital acquired infections [16-17], [17]. Also, access to information about to SPs will bring about enhancement of knowledge and better adherence to SPs [18]. Apparently, Chi-Square result showed that gender, marital status, years of experience and job category is significantly associated with knowledge of SPs ( $p < 0.05$ ).

Furthermore, this study showed that there is non-compliance with SPs in hospitals. This study reinforces that only few HCWs (35.8%) complied with SPs. The non-compliance with SPs is as a result of poor safety culture in the hospitals, insufficient delivery of PPEs, absence of supervision, absence of incident reporting, irregular trainings on SPs, nonexistence of monitoring, too much workloads, poor time management, nonexistence of punitive actions on non-compliance, prompt handling in emergency situations, assumptions that patient lacks infection, general self-efficacy, nonexistence of checks by hospital management amongst others. This finding agrees with [19] who reported that 75% of HCWs were below acceptable safety levels of practice SPs. [20] also reported median scores for compliance with SPs, HCWs complained of insufficient resources to practice SPs. In developing countries like Nigeria, SPs are either absent, not adhered to, or not enforced [21-22] studies and affirmation of insufficient compliance with SPs by HCWs and they stated that observance of SPs is internationally suboptimal. In Rwanda, there was a fall in adherence to 32.1% between 2015 and 2016 [23]. More previous researchers reported poor compliance with SPs by HCWs [12, 17], [24-25], [26], [12]. Much lower results (12% and 16.1%) on HCWs adherence to SPs have been were reported [23, 27] [23]. This study's disagrees with findings by [28] who stated that 77.9% of HCWs practice SPs and disease control in northern Nigeria.

[15] also opined that HCWs (70.1%) had a good adherence to SPs. Other studies also opined that hand washing, use of PPEs and safer devices (like retractable lancets), reduces the risk of OHHS [29], [17], [29-30], [31]. These measures also help reducing the spread of infectious agents in the HCFs [30]. [32] also advocated that adherence to SPs, is the most effective way of preventing transmission of BBPs. Yet, even with published procedures on disease control and adverse health concerns following noncompliance, major issues still exists around HCWs' adherence to SPs. More so, the Chi-Square result showed that gender, years of experience and job categories is significantly associated with compliance with SPs ( $p < 0.05$ ). Female HCWs are more likely not to comply with standard precaution than male HCWs and HCWs with job experience less than 10 years are more likely not to comply with SPs HCWs with job experience 11 year and above. This finding agrees with [33] who opined that the level of awareness of SPs rises with increased

Comment [K24]: Write last name of author first.

Comment [K25]: See K24.

Comment [K26]: Source? As they are not all from study findings.

Formatted: Highlight

Formatted: Highlight

Comment [K27]: Findings differ from?

Formatted: Highlight

Comment [K28]: Any explanation?

Formatted: Highlight

Comment [K29]: In full first.

Formatted: Highlight

Comment [K30]: Concur?

Formatted: Highlight

years of experience at work. Also, [15] reported that HCWs gender and years of work experience is significantly associated with compliance with SPs.

Formatted: Highlight

## V. Conclusion

This study revealed that the Health Belief Model (HBM) is not obtainable or practiced in the hospitals in River State. Even with high perception of susceptibility to infection, or high perception of the seriousness of the problem, HCWs will still not comply with SPs. The non-compliance with SPs reported in this study poses devastating consequences in the hospitals, such as increase in occupational health hazards (OHHs), decline in sustainable development and growth in the health sector as well as a healthy society at large. Hence this study recommends the following:

Comment [K31]: But not presented in the discussion.

Comment [K32]: Make specific (eg. to HCWs, Healthcare system in Nigeria) and feasible recommendations.

Comment [K33]: A paragraph would be better than bullets/separated ideas.

Comment [K34]: Not clear!

- i. Laws and policies on SPs should be made by the government through the help of their legislation.
- ii. The hospital management should enforce a system of surveillance for OHHs. This will assist in improving HCWs knowledge of SPs, as well as encouraging adherence with SPs.
- iii. They should encourage regular provision of PPEs, their maintenance (for those that are not disposable) and their timely replacement when worn out.
- iv. HCWs should also be enlightened that they are obligated to comply with SPs as this will reduce their exposure to OHHs.

## References

- 1) LaDou J. International Occupational Health. *International Journal of Hygiene and Environmental Health* 2003; 206: 303-313.
- 2) Ahasan MR, Partanen T. Occupational health and safety in the least developed countries- a simple case of neglect. *Journal of Epidemiology*, 2001; 11(2), 74-80. <https://doi.org/10.2188/jea.11.74>
- 3) Watterson A, Ladou, J. Health and safety executive inspection of U.K. semiconductor manufacturers. *International Journal of Occupational and Environmental Health*. <https://doi.org/10.1179/oeh.2004.10.3.352>. 2003
- 4) Yassi A, Lockhart K, Copes R, Kerr M, Corbiere M, Bryce E, Team SS. Determinants of healthcare workers' compliance with infection control procedures. *Healthc Q*, 2007; 10(1), 44-52. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/17326369>
- 5) Zungu L, Sengane M, Setswe K. Knowledge and experiences of needle prick injuries (NPI) among nursing students at a university in Gauteng, South Africa. *South African Family Practice*, 2008; 50(5), 48-48c. <https://doi.org/10.1080/20786204.2008.10873762>
- 6) Askarian M, Yadollahi M, Kuoachak F, Danaei M, Vakili V, Momeni M. "Precautions for health care workers to avoid hepatitis B and C virus infection." *The international journal of occupational and environmental Medicine* 2011; 2, no. 4
- 7) Hinkin J, Gammon J, Cutter J. Review of personal protection equipment used in practice. *British Journal of Community Nursing*. <https://doi.org/10.12968/bjcn.2008.13.1.27978>. 2008



- 8) Akagbo SE, Nortey P, Ackumey MM. Knowledge of standard precautions and barriers to compliance among healthcare workers in the Lower Manya Krobo District, Ghana. *BMC Research Notes*, 2017: 10(1). <https://doi.org/10.1186/s13104-017-2748-9>
- 9) Yamane T. Statistics, an Introductory Analysis, 2nd Ed., New York 1967.
- 10) Zhang M, Wang H, Miao J, Du X, Li T, Wu Z. Occupational exposure to blood and body fluids among health care workers in a general hospital, China. *American Journal of Industrial Medicine*, 2009: 52(2), 89–98. <https://doi.org/10.1002/ajim.20645>
- 11) Mahajan A, Bedi R. Knowledge and attitude of dentists in India towards patient safety – A cross-sectional study. *Journal of Patient Safety & Infection Control*, 2015: 3(3), 134–138. <https://doi.org/10.1016/j.jpsic.2015.11.002>
- 12) Ndu AC, Arinze-Onyia, SU. Standard precaution knowledge and adherence: Do doctors differ from medical laboratory scientists? *Malawi Medical Journal*, 2017: 29(4), 294–300. <https://doi.org/10.4314/mmj.v29i4.3>
- 13) Aluko OO, Adebayo AE, Adebisi TF, Ewegbemi MK, Abidoye AT, Popoola BF. Knowledge, attitudes and perceptions of occupational hazards and safety practices in Nigerian healthcare workers. *BMC Research Notes*, 2016: 9(1), 71. <https://doi.org/10.1186/s13104-016-1880-2>
- 14) Hassan M, Awosan KJ, Nasir S, Tunau K, Burodo A, Yakubu A, Oche MO. Knowledge, risk perception and hepatitis B vaccination status of healthcare workers in Usmanu Danfodiyo University Teaching Hospital, Sokoto, Nigeria. *Journal of Public Health and Epidemiology*, 2016: 8(4), 53–59. <https://doi.org/10.5897/JPHE2015.0795>
- 15) Adebimpe WO. Knowledge, Attitude, and Practice of Use of Safety Precautions Among Health Care Workers in a Nigerian Tertiary Hospital, 1 Year After the Ebola Virus Disease Epidemic. *Annals of Global Health*, 2016: 82(5), 897–902. <https://doi.org/10.1016/j.aogh.2016.07.004>
- 16) Garcia-Zapata MR.-C, Silva AC, Valadares GJ, Veiga TAF, Aparecida PM, Garcia-Zapata MTA. Standard precautions: knowledge and practice among nursing and medical students in a teaching hospital in Brazil. *International Journal of Infection Control*, 2010: 6(1). <https://doi.org/10.3396/ijic.v6i1.005.10>
- 17) Abdulraheem IS, Amodu MO, Saka MJ, Bolarinwa OA, Uthman MMB. Knowledge, Awareness and Compliance with Standard Precautions among Health Workers in North Eastern Nigeria. *Journal Community Medical Health Education* 2012: 2, no. 131: 2.
- 18) Vaz, K., D. McGrowder, R. Alexander-Lindo, L. Gordon, P. Brown, and R. Irving. (2010). "Knowledge, awareness and compliance with universal precautions among health care workers at the University Hospital of the West Indies, Jamaica." *The international journal of occupational and environmental medicine* 1, no. 4: 171-181.
- 19) Askarian M, Mirzaei K, Mundy LM, McLaws, ML. Assessment of Knowledge, Attitudes, and Practices Regarding Isolation Precautions among Iranian Healthcare Workers. *Infection Control & Hospital Epidemiology*, 2005: 26(01), 105–108. <https://doi.org/10.1086/502495>
- 20) Ogoina D, Pondei K, Adetunji B, Chima G, Isichei C, Gidado S. Knowledge, attitude and practice of standard precautions of infection control by hospital workers in two tertiary hospitals in Nigeria. *Journal of Infection Prevention*, 2015: 16(1). <https://doi.org/10.1177/1757177414558957>
- 21) Okaro A, Eze C, Ohagwu C. Awareness, Knowledge and Attitude of Radiographers towards HIV/AIDS Patients Attending Radiology Clinics in Enugu State, Nigeria.

- European Journal of Scientific Research*, 2010: 39(3), 440–447. Retrieved from [http://www.acrobatplanet.com/go/Knowledge\\_and\\_Attitude\\_of\\_Radiographers.pdf](http://www.acrobatplanet.com/go/Knowledge_and_Attitude_of_Radiographers.pdf)
- 22) Gammon J, Morgan-Samuel H, Gould D. A review of the evidence for suboptimal compliance of healthcare practitioners to standard/universal infection control precautions. *Journal of Clinical Nursing*, 2008: 17(2), 157–167. <https://doi.org/10.1111/j.1365-2702.2006.01852.x>
  - 23) Haile TG, Engeda, EH, Abdo AA. Compliance with Standard Precautions and Associated Factors among Healthcare Workers in Gondar University Comprehensive Specialized Hospital, Northwest Ethiopia. *Journal of Environmental and Public Health*, 2017. <https://doi.org/10.1155/2017/2050635>
  - 24) Fadeyi A, Fowotade A, Abiodun MO, Jimoh AK, Nwabuisi C, Desalu OO. Awareness and practice of safety precautions among healthcare workers in the laboratories of two public health facilities in Nigeria. *The Nigerian Postgraduate Medical Journal*, 2011: 18(2), 141–146. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/21670783>
  - 25) Karim J, Al-Saraji M, Al-Mousawi F, Al-Haddad Z, Al-Sharaf D, Marwan Y Akhtar S. Knowledge and self-reported practice of universal precautions among Kuwait University medical students in their clinical years. *Medical Principles and Practice*, 2012: 21(4), 328–333. <https://doi.org/10.1159/000335278>
  - 26) Musu M, Lai A, Mereu NM, Galletta M, Campagna M, Tidore M, Coppola RC. Assessing hand hygiene compliance among healthcare workers in six Intensive Care Units. *Journal of Preventive Medicine and Hygiene*, 2017:58(3), E231–E237. Retrieved from <http://www.ncbi.nlm.nih.gov/pubmed/29123370>  
<http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC5668933>
  - 27) Okechukwu EF, Motshedisi, C. Knowledge and practice of standard precautions in public health facilities in Abuja, Nigeria. *International Journal of Infection Control*, 2012: 8(3). <https://doi.org/10.3396/IJIC.V8I3.10003>
  - 28) Amoran O, Onwube O. Infection control and practice of standard precautions among healthcare workers in northern Nigeria. *Journal of Global Infectious Diseases*, 2013: 5(4), 156–163. <https://doi.org/10.4103/0974-777X.122010>
  - 29) Centers for Disease Control and Prevention. National Healthcare Safety Network patient safety component manual. [http://www.cdc.gov/nhsn/TOC\\_PSCManual.html](http://www.cdc.gov/nhsn/TOC_PSCManual.html). 2011
  - 30) Huang C, Wenjun M, Susan S. The hygienic efficacy of different hand-drying methods: a review of the evidence. *Mayo Clinic Proceedings*. 2012: Vol. 87. No. 8. Elsevier, .35.
  - 31) Towe L. Standard Precautions Policy. Available on <https://www.ashfordstpeters.info/images/policies/INF75.pdf>. 2016.
  - 32) Powers D, Armellino D, Dolansky M, Fitzpatrick J. Factors influencing nurse compliance with Standard Precautions. *American Journal of Infection Control*, 2016: 44(1), 4–7. <https://doi.org/10.1016/j.ajic.2015.10.001>
  - 33) Mcgrowder D, Vaz K, Alexander-lindo R, Gordon J, Brown P, R, I. Knowledge, Awareness and Compliance with Universal Care Workers at the. *Indian Journal of Occupational and Environmental Medicine*, 2010: 1(4), 171–181.