

Original Research Article

HERBOMINERALS AND ANTIBACTERIAL ACTIVITIES OF *Allium sativum* L EXTRACTS ON PATHOGENIC BACTERIA CAUSING MENINGITIS IN SUB-SAHARAN AFRICA, ZARIA, KADUNA STATE, NIGERIA

Comment [v1]: Sp.check and centre

ABSTRACT

Bacterial Meningitis (BM) is the most common serious infection of the central nervous system (brain and spinal cord). This research aims to determine herbominerals and to evaluate the in vitro antibacterial activity of the extracts (JEAS, EEAS and AEAS). The collected bulbs of A. sativum (600 g) were washed and air dried under shade for 2 hours and the dry scaly outer covering was peeled-off to obtain the fresh garlic cloves which were then divided into three parts of 200 g each. These three portions were crushed separately for cold extraction. The first portion was homogenized and poured into a muslin cloth to squeeze out the juice, while second and third portions were homogenized and submerged into 500 ml of 96% ethanol and 500 ml of distilled water respectively for 24 hours and both filtered after thorough shaking. The antibacterial activity of bulbs of A. sativum juice, ethanolic and aqueous (JEAS, EEAS and AEAS) extracts as folkloric medicine against Neisseria meningitides, Streptococcus pneumoniae, Haemophilus influenzae, Klebsiella pneumoniae and Escherichia coli were determined using Agar well diffusion and broth dilution method. Distilled water, concentrated nitric acid (HNO₃) and hydrochloric acid (HCl) were used to digest the extract, which was then heated in water bath at 90°C and filtered to obtain the filtrate for the analytical studies for A. sativum nutritional composition and zeolite herbominerals. The micro-herbominerals with their proximate values were observed were Nano pharmacologic of Silver, Manganese, Zinc, Iron and Selenium; which exerts biocidal properties as well as stimulates immune system to cushioning the challenges of the BM pathogens. The minimum inhibitory concentration (MIC), minimum bactericidal concentration (MBC) and phytochemical screening of the extracts were evaluated. The results obtained showed that the juice and ethanolic extracts were potent, inhibiting the growth of clinical isolates with zone of inhibition ranging from 14-36mm. The extracts inhibited bacterial isolates in the concentration dependant manner with MICs ranging between 0.02-15mg/ml and MBCs 0.04-5mg/ml. Phytochemical screening of the extracts revealed the presence of alkaloids, flavonoids, anthraquinone, carbohydrates, fats and oils, steroidal ring, saponins and terpenoids. This experimental investigation has provided the scientific validation basis for the ethnomedical use of A. sativum as a remedy to treat bacterial meningitis locally as anti-infectious chemotherapeutic.

Comment [v2]: determine

Comment [v3]: the mineral composition

Comment [v4]: for the first time give the expansion for the extract

Comment [v5]: subscript

Comment [v6]: superscript

Comment [v7]: delete

Comment [v8]: reduce abstract to 250 words.

Keywords: Herbominerals, Antibacterial, *Allium sativum*, Bacterial Meningitis, Sub-Saharan Africa, Kaduna State.

Comment [v9]: herbominerals

1.0 INTRODUCTION

Bacterial Meningitis (BM) is the most common serious infection of the central nervous system (pia mater and arachoid inflammation called leptomeningitis and inflammation of brain parenchyma called encephalitis) and spinal cord (myelitis) [1].

Comment [v10]: Sp.check

Bacterial endotoxins in the blood infect the pia mater, arachnoid and subarachnoid space leads to cerebrospinal fluid (CSF) meningeal irritation and the clinical trial of headache, fever and meningism [2]. Antibiotics and vaccines meant to reduce morbidity and mortality of BM as modern therapy are available in BM radical treatment. Despite all these efforts put in place, BM remains associated with unacceptable rate of high morbidity and mortality in Sub-Saharan Africa, including Nigerian Northern States [3].

Comment [v11]: Sp.check

Aside of the massive side effects associated with the conventional antibiotics; there has also been the emergence of resistant strains of the infecting organisms probably from the frequent irrational use of the available antibiotics to treat conditions of unknown aetiology. And often resistance hampers the control of bacteria in general and tends to increase cost of healthcare services for bacterial chemotherapeutics [4].

Comment [v12]: bacterial

Therefore, bacterial meningitis (BM) is a devastating air-borne infectious disease of humans, and it is of huge health burden as it can easily kill or maim when meningeal pathogens invaded blood stream. The BM causes cellular hemorrhagic sepsis, metabolic, endocrine and neurologic disorders. This burden of the disease is highest in the developing countries and especially in the immunity compromised rural populations [5]. Inflammation of the meninges causes leakage of the infected cerebrospinal fluid (CSF) and alteration of the brain system (cognitive deficit) giving

rises to meningism (stiff neck, severe headache, fever, rashes, shortness of breath or noisy breathing, hydrocephalus (swelling or oedema of the brain microglia and astrocytes due to inappropriate antidiuretic hormone secretion in the brain parenchyma and leptomeninges).

Comment [v13]: rise

High cost of acquiring synthetic antibiotics and the inconveniences of seeking for effective treatment due to using exorbitantly expensive drugs make life difficult for people, considering the economic status of an average Nigerian as a low income earner [3]. Moreover re-emergence of BM persists in epidemic meningitis belt of Africa. Therefore, to overcome BM, new knowledge of natural products against BM pathogens is urgently needed; especially new drugs are required for BM mitigation which differs from existing medicines [4].

As for the prevalence of meningeal bacterial organisms, the potential of epidemic virulence of meningeal bacteria organism in Nigeria varies with the geographical location, time of season, climate and pathogenic serotype strains of the causative organism [3]. Six subspecies serotypes of *meningococci* (A, B, C, W-135, X and Y) have been clinically recognized in Sub-Saharan African countries of which serotypes A and C *meningococci* were found to be more prevalent and predominant among other serotypes which often occur as co-morbid infections [1]. The epidemics of Sudan in 2006 was of serotypes W-135 and X [1], 2009). The most prevalent meningeal bacteria organism in Nigeria is of the serotype A and C strains of *Neisseria meningitides* seen abundantly during the harmattan period when humidity is very low, usually around November-June and which is the breeding time of the organisms. *Neisseria meningitides* is the most implicated bacterial organism that causes meningitis and constitutes about 80% of the epidemic cases in Africa [6]. Other bacterial organisms (*Haemophilus influenzae* type b, *Streptococcus pneumoniae*, *Escherichia coli*, *Klebsiella pneumoniae*) also cause meningitis to a

Comment [v14]: bacterial

Comment [v15]: bacterial organism causing meningitis

Comment [v16]: which is seen

Comment [v17]: which happens to be the main breeding period

lesser extent (about 20% of the epidemic cases) [6, 7]. Thus, these five bacterial organisms are those mostly known to be involved in meningitis, systemic septic-inflammations and hemorrhage in Sub-Saharan Africa including Nigeria [6]. Bacterial meningitis caused by *H. influenzae* and *meningococci* have a better prognosis than of *S. pneumoniae* and *E.coli* that mainly attack neonates and growing children. The group B *streptococci* (subtype III) occur mainly during the first week of life (newborns), while *Escherichia coli* and *Klebsiella pneumoniae* are of both newborns and growing children less than five years of age. *Haemophilus influenzae* type B, *Neisseria meningitides* (meningococci) and *Streptococcus pneumoniae* affect mostly the adults, but people over 50 years have increased risk of *Escherichia coli* [3].

Comment [v18]: affect mostly

1.1 African BM Epidemiology

Comment [v19]: Epidemiology of Bacterial Meningitis

Annual epidemic of BM occurs during the hot dry season in Sub-Saharan African countries while BM strikes during the cold rainy season in Western countries [8]. About 1.2 million cases of BM annually worldwide with 135,000 deaths and is now a 'top10' infectious disease cause death worldwide and half survivors deformed [5]. African BM (epidemic) ranges from 100 to 1000 per 100,000 populations and over 3,000 populations per day were infected during explosive meningitis belt while 20 per 100,000 populations in the western world (USA, UK and Europe) [7]. Thus, African BM (non-epidemic) ranges from 10 to 20 per 100,000 for Sub-Saharan African countries while 1 to 5 per 100,000 for western countries [9].

Comment [v20]: In western countries it strikes.....

Comment [v21]: have been reported annually

Comment [v22]: comma

Comment [v23]: causing

Comment [v24]: with reports of deformity in 50 percent of the survivors

Comment [v25]: individuals

Comment [v26]: in the belt

Comment [v27]: remove bold

1.2 The Pathogenesis and Pathophysiology of Bacterial Meningitis

BM involves a complex interplay between virulent factors of the pathogens and the host immune responses. Therefore, BM develops when virulent factors of the pathogen overcome host defense mechanisms. Meningeal bacteria usually weaken the

functions of the immune defense system which predisposes the body system to damaging opportunistic infections. Bacterial infection in particular often leads to septicaemia or persistent multiplication that traumatizes the blood vessels. Bacteria lyses, immune activator binding, leucocyte infiltration causes glial cells and astrocytes to be inflamed, inflammatory factors as TNF, COX-2, NK, IL, iNO, and Phospholipase A2 which triggered meningeal inflammation. Neuropathologic effect of BM, brain oedema formation due to brain parenchyma inflammation, impairment of CSF hydrodynamics (inappropriate secretion of antidiuretic hormone, IASADH), increased intracranial pressure leads to hydrocephalic head in neonates [10, 3]. BM is pathologically characterised by metabolic, endocrine, neurologic disorders and amnesia as well as dysfunctions of multiple organs such as heart, liver, kidneys, and lungs [1].

Comment [v28]: lyses causes binding of immune activators and release of inflammatory factors like.....and leukocyte infiltration causes glial cells and astrocytes to be inflamed finally leading to meningeal inflammation

Comment [v29]: include

Comment [v30]: leading to

In the extreme conditions involving intravascular coagulations, organ failure and / or sudden death may occur related to the rapid shutdown of the vital organs. Haemorrhage is a severe form of blood loss that can cause general shock as a result of reduced tissue perfusion and / or ischaemia [11, 10]. Meningeal bacteria causing reduction in the levels of blood clotting cells (platelets), this leads to severe erythematous purpura, DIC and uncontrollable bleeding which can be internal haemorrhage in the vital organs of the body [12]. These blood clots can reduce or block blood flow through the body blood vessels due to bacteria coagulases lead to the damage and sudden death of body's organs. In DIC, the increased clotting uses up platelets and clotting factors in the blood, it deregulates normal physiologic mechanisms of the blood. Furthermore, it can cause internal and external bleeding in meningitis patients mainly infants [10]. Thus, internal bleeding occurs inside the body while external bleeding occurs underneath or from the skin or mucosa. These

Comment [v31]: delete

Comment [v32]: expand for the first time

Comment [v33]: bacterial

meningitis pathogens cause sepsis of the blood, fever, diarrhoea and sudden death [10].

Pains (abdominal, chest), swelling, hemoptysis (bloody vomitus), rashes, photophobia (light intolerance), hyperthermia, hypoglycemia, convulsion/seizures, phonophobia (high noise intolerance), drowsiness, confusion, clouded vision, loss of consciousness, **arthralgia**, altered mentation, sight and hearing loss and crippled limbs [12, 6].

Comment [v34]: The associated symptoms include pain.....

Comment [v35]: Sp.check

Comment [v36]: Give a diagrammatic representation




Figure 1: shows the pathogenesis of bacterial meningitis.




Figure 2: shows the anatomical pathogenesis of bacterial meningitis. Adapted from www.anatomyatlas.com

1.3 Aetiologic Factors of Bacterial Meningitis

- Malnutrition
- Climate change usually hot weather
- Overcrowding (as in crowded quarters, or gatherings with infected persons who most times are asymptomatic carriers of the pathogens)
- Deficient immune system
- Poor sanitation (environmental hygiene) that breeds the causative organisms
- Symptoms of inappropriate antidiuretic hormone secretion (SIADHS) in bacterial endotoxin-induced oedema of the body due to endocrine crises [5, 3].

However, *A. sativum* (AS) has a long history of use as food condiment and also a high value of use traditionally for many ailments including antimicrobial effects. However, its specific activity on meningeal bacteria as well as antihemorrhagic activity in condition of meningococcal-induced cellular injuries has actually not been reported in

Comment [v37]: Before telling about Allium tell few points about the treatment for meningitis done and its drawbacks...

literature. Also, a scientific validation of this utilization has not been previously made to the best of my knowledge [3].

Comment [v38]: Who??/author of this paper or author of the third reference.

1.4 Description and Origin of *Allium sativum*

African Traditional medicine constitutes an important source of drugs for ethnopharmacological investigation. Thus, Traditional medicine by WHO definition is the sum total of the knowledge, skills, and practices based on the theories, beliefs and experiences indigenous to different cultures, whether explicable or not, used in maintenance of health in prevention, diagnosis, improvement or treatment of physical and mental illnesses [13]. Various medicinal food-plants and animal products-supplements are available for use in certain immune-deficiency disease conditions related to malnutrition such as hemorrhagic sepsis. Furthermore, phytotherapy still remains a habitual part of health care system wholly or in part especially in rural communities [3]. *Allium sativum* (AS) is one of such medicinal food-plant that has a long history of use as food condiment and it also has a high value of use traditionally for many ailments including typhoid fever, ulcer, cholera, dysentery etc and thus, there is a need to investigate its specific *in vitro* antibacterial activity on meningeal bacterial organisms as well as in condition of meningococcal-induced cellular injuries having been known to boost the immune system [14].



Figure 3: Garlic Plant in its Natural Habitat, Source: Rader and McGuinness, [15]

Scientific name: *Allium sativum* Linn

Family: Liliaceae

Common name: Garlic

Hausa name: *Tafarnuwa*

Yoruba name: *Allubosa ayu*

Igbo name: *Allibasa ayo*

Ebira name: *Allivasa ahono* [3].

1.5 Folkloric and Ethnomedical Uses

A. sativum has an ancient use as antiseptic, antibacterial, carminative, bleeding of the nose, diaphoretic, anthelmintic, antimicrobial, antifatulence, and analgesic. It is also used locally for confluent smallpox, typhoid fever, dropsy or congestive heart failure, pulmonary tuberculosis or phthisis and for eye and ear infections [16]. It has also been reported to have antitumour and anticholesterol-anemia properties whereby it is used to decrease low density lipoprotein and / or increase high density lipoprotein as to lower blood pressure. In China and Arabia, it has a history of use as energizer or palliative and it is believed to have a natural nutritional healing power for cough, hay fever, pile or dysentery and also to promote digestion (dyspepsia) [16,17].

Comment [v39]: Arrest bleeding

Comment [v40]: check

2.0 METHODOLOGY

2.1 Collection and Identification of Plant Material

The bulbs of *A. sativum* were obtained as fresh farm products from Valiki Market, Okene, Kogi State, Nigeria in November, 2014. The plant's identification was authenticated by Mallam Umar S. Gallah at the Herbarium Unit of the Department of Biological Sciences, Ahmadu Bello University, Zaria, and compared with voucher specimen number of 2156 for future reference.

2.2 Extraction of Plant Material

The collected bulbs of *A. sativum* (600 g) were washed and air dried under shade for 2 hours and the dry scaly outer covering was peeled-off to obtain the fresh garlic cloves which were then divided into three parts of 200 g each. These three portions were crushed separately for cold extraction according to the method of Fattouch *et al.*, [18]. The first portion was homogenized and poured into a muslin cloth to squeeze out the juice, while second and third portions were homogenized and submerged into 500 ml of 96% ethanol and 500 ml of distilled water respectively for 24 hours and both filtered after thorough shaking. The first and second portions were freeze dried, while the third portion was evaporated over water bath at 50°C to obtain the powdered yield. The three samples obtained were then stored in separately labeled air-tight container for later use. The percentage (%) extract yield was calculated as follows:

$$\% \text{ yield} = \text{weight of extract (g)} / \text{weight of garlic cloves (200 g)} \times 100$$

2.3 Preparation of the Extracts

Working concentrations to be used for the study were prepared from the obtained extracts (JEAS, EEAS and AEAS) using distilled water as a diluent. The required

various concentrations to be used for each experiment were always freshly prepared from the stock solution of the extracts.

2.4 Test Clinical Bacteria Isolates

Five pathogenic clinical isolates (three anaerobic organisms): *Neisseria meningitides*, *Streptococcus pneumoniae*, *Haemophilus influenzae* and (two aerobic organisms) *Klebsiella pneumoniae* and *Escherichia coli* obtained from the Department of Medical Microbiology, Ahmadu Bello University Teaching Hospital (ABUTH) Shika, Zaria in Kaduna state, Nigeria were cultured in Mueller Hinton Agar (MHA) to obtain colonies of the organisms as described by Monica [19] and Sharma and Aneja [20].

2.5 Phytochemical Screening of *Allium sativum*

The methods of [13] were used to screen for the presence of various phytochemical constituents in the *A. sativum* extracts. Phytochemical constituents screened include alkaloids, anthraquinones, cardiac glycosides, carbohydrates, fat and oils, flavonoids, saponins, steroids, terpenoids and tannins. 2.0g of powders of each of the three extracts were dissolved in 100ml of distilled water for this screening experiment.

2.6 Herbomineral Analysis and Proximate Values of *Allium sativum* Extract (JEAS)

Analytical automated instruments including spectronic 21D / flame photometer, unicam 969 AA spectrometer (AAS), fast sequential atomic spectrometer AA240FS and AA spectrophotometer Buck205 were used to analyze their proximate values in the JEAS according to the methods described by Naturopathy [21]. Distilled water, concentrated nitric acid (HNO₃) and hydrochloric acid (HCl) were used to digest the extract, which was then heated in water bath at 90°C and filtered to obtain the filtrate for the analytical studies for *A. sativum* zeolite herbominerals. The presence of dietary

Comment [v41]: Sp.check

Comment [v42]: subscript

phytonutrients including carbohydrates, proteins, dietary fibres, fats and oils, vitamins as well as the zeolite herbomineral elements and their proximate values were determined. 2.0g of JEAS powder was dissolved in 100ml of DH₂O for the quantitative determination of vitamins, macro and micro-elements and other nutrients. 10ml for vitamin analysis (200mg), 30ml for macro/micro (zeolite) element analysis (600mg) and 30ml for other nutrients analysis (600mg) respectively.

Comment [v43]: subscript

2.7 Determination of Antibacterial Activity of *Allium sativum* Extracts

The bacterial cultures were prepared by transferring with sterile wire loop, each bacterium cell into a Nutrient Broth Medium which was incubated at 37°C for 24 hours as described by WHO-BS [22] to obtain a colony of the bacterial cells. The cells were maintained in Nutrient Broth Media in universal bottles labeled for each bacterium. Three tubes (1, 2, 3) each containing 9 ml of normal saline were set up for 3-fold serial dilution (1:1000) for the gram-positive bacterium, in which 1 ml of the overnight culture was transferred into the first test tube and mixed; and 1ml of this then taken into the second test tube from which another 1ml was removed into the third tube which was then deemed to be 1×10^3 colony forming unit / ml (CFU/ml).

Comment [v44]: serially diluted

The same serial dilution, but in a fourth tube containing 4.5 ml of normal saline and 4-fold dilution (1:5000) was also performed for the gram-negative bacteria to the last concentration of 1×10^5 CFU/ml. The last diluted concentrations (5 test tubes of the bacterial organisms) were incubated at 37°C for 24 hours after which, the bacterial organisms (suspension) were inoculated on prepared Molten sterile Mueller-Hinton agar plates by taking 2 ml of each of the organisms and flooding it over the agar surface by agar well diffusion method of Sharma and Aneja, [20]. Four different concentrations (10, 15, 20 and 25 mg/ml) and 5 mg/ml of the standard drug (cefuroxime axetil) were used to study the growth inhibitory effect for each of the

three extracts of *A. sativum*. Wells were bored on the agar dishes and a drop of the molten agar was used to seal the bottoms of the bored wells prior to filling them with 0.2 ml of each the various drug concentrations. The plates were then kept for 1 hour to diffuse and then incubated at 37°C for 24 hours. The experiment was performed in duplicate and the zones of growth inhibition around the two wells for each drug concentrations were measured in millimeter using a ruler. The mean of the duplicate experiments for each concentration of the extracts were then calculated and recorded as the growth inhibitory zone of the extract concentrations for each of the organisms [22, 19].

Comment [v45]: of the drug concentration.

2.8 Determination of Minimum Inhibitory Concentration (MIC) and Minimum Bactericidal Concentration (MBC)

The same extracts concentrations of 10, 15, 20 and 25 mg/ml of the three *A. sativum* were used for MIC and MBC experiments. Serial dilutions were made for each concentration as well as for the standard drug whereby ten test tubes with each containing 9 ml distilled water were set up for each concentration and 1 ml of each concentration taken into the first test tube and from which 1 ml was taken into the second test-tube and continuously like that until the last test tube after which the 1ml taken from it was discarded. Then 2 ml of each of these mixtures (varied concentrations) in test tubes were taken respectively into various petri-dishes containing 8 ml of molten agar that had been labeled for each diluted mixture and then mixed to form broth dilution. Similar dilution was carried out for each of the three extracts and in all; a hundred and fifty mixture petri-dishes were gotten (50 for JEAS, 50 for EEAS and 50 for AEAS). All of these were kept for 2 hours to solidify after which a sterile forcep was used to place 5 sterile filter paper discs of 6 mm in diameter on each solidified agar plate surface. A micropipette was used to inoculate

Comment [v46]: Serially diluted

Comment [v47]: were received

0.02 ml of each of the five bacterial isolates on each of the discs and this was done for all the four concentrations of the three extracts under study and also for the standard drug. The plates were then incubated at 37⁰C for 24hrs. The lowest diluted concentration of each extract stock concentration in which there was no visible growth of any of the bacterial organisms was considered as the Minimum Inhibitory Concentration (MIC) for the organism in question [19]. The Minimum Bactericidal Concentration (MBC) of the various extracts were performed from the lowest concentration (MIC) of the dilutions of the 4 stock concentrations and the higher concentrations above it. For instance, the 10 mg/ml has its diluted concentration of 2.5 mg/ml as its MIC and thus, three test tubes labeled as 2.5, 5.0 and 10 mg/ml and containing 2 ml nutrient broth and 3 drops of 3% v/v TW 80 were set for MBC. Then, filter paper discs of the bacterial organisms from the corresponding MIC experiment were transferred into the test tubes respectively and again incubated at 37⁰C for 24 hours and checked for growth inhibition [19, 20].

2.9 Statistical Analysis

Results obtained were recorded as mean \pm SEM and subjected to one way analysis of variance (ANOVA) and where significant differences exist, means were compared using Waller Duncan test was performed using Statistical Analysis System (SAS, software version 2002) at 0.05 significant level (P<0.05).

Comment [v48]: and was performed

3.0 RESULTS AND DISCUSSION

Results of percentage yielded of the three extracts were shown in Table 1. The results showed Ethanolic extract of *A. sativum* yielded highest percentage with light yellow powder, Aqueous extract of *A. sativum* was second yielded percentage with brown powder and Juice extract of *A. sativum* was least percentage yielded as depicted below. The results of phytochemical screening of the bulbs extracts of *A. sativum* are shown in Table 2. The results showed the presence of alkaloids, anthraquinones, carbohydrates, cardiac glycosides, fats and oils, flavonoids, saponins, steroidal ring, terpenoides and tannins. The presence of phytochemicals in the bulbs extracts (Table 2) showed that the extracts possess strong antibacterial properties. These results are agreement with similar study by Abdullahi *et al.*, [13].

Table 3, were demonstrated to possess health benefits and are used as combined tools for treating bacterial meningitis. Its zeolite herbominerals such as silver ions, zinc ions, selenium ions have bactericidal effect on bacterial meningitis pathogens through their nanopharmacologic and chelation therapy [23]. It protects entire body by stimulating the immune system, prevents infectious diseases, improves health, and supports the structures (collagens) [17].

The Table 4: shows the results of antibacterial effectiveness of *A. sativum* bulbs extracts against the test clinical bacteria isolates. Results showed that the activity of the bulbs extracts against the test bacteria decrease with decrease in the concentration ie antibacterial effect of the extracts are concentration dependent manner; with the juice extract demonstrating higher antibacterial activity (12.0-36.0 mg/ml) than the ethanolic extract. While aqueous extract lost activity due to heat applied because *A. sativum* active compounds are heat labile biodegradable. It has been observed that the more polar the solvent, the higher the yield of extract shown in the Table 1, but the

Comment [v49]: yield

Comment [v50]: delete

Comment [v51]: and stood second in its yield percentage

Comment [v52]: delete

Comment [v53]: the least in its percentage yield

Comment [v54]: in agreement with Abdullahi

Comment [v55]: italic

Comment [v56]: bulb extracts

Comment [v57]: bacterial

Comment [v58]: bulb extracts

Comment [v59]: decreased

Comment [v60]: occurred in a

Comment [v61]: the heat

Comment [v62]: are heat labile and

Comment [v63]: delete

juice and ethanolic extracts are potent due to their freeze drying process with high bioactive principles.

Figure 4 and Table 5,6,7,8,9 showed the minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) of the three extracts against the test clinical isolates respectively. These results showed the values obtained from experiment are quite higher for the juice extract than that of ethanolic extract, suggesting that extraction with ethanol could produce better active antibacterial phytochemicals with active organosulphorus volatile oil compounds which are contained in bulbs through freeze drying procedures. These results of the present study are in agreement with similar study by Sen *et al.*, [24]; Abdullahi and Ajayi, [25]. The observed antibacterial effect of the bulbs extracts on the clinical bacterial isolates though *in-vitro* demonstration of antibacterial experiment is an indication; that *A. sativum* bulbs extracts could be effective in the management of bioterrorising infections such as BM cause by these bacteria (Table 6 and 7).

Comment [v64]: bulb

Comment [v65]: bulb extracts

Comment [v66]: caused

3.1 Extraction of the Plant Material

The calculated percentage yield from the 200 g of *Allium sativum* L bulbs used for the extraction was as given in table 1 below:

Table 1: Percentage Yield of *Allium sativum* Extracts

Extract	% Yield (w/w)	Observed Colouration
JEAS	12.7	light yellow powder
EEAS	17.2	light yellow powder
AEAS	14.7	Brown powder

Comment [v67]: give the order in ascending

3.2 Phytochemical Screening of the Extract

The table below is a summary of the phytochemical components or secondary metabolites of the extract. From the obtained result, anthraquinone and tannins were not present in the extract of the bulbs.

Table 2: The Phytochemical Constituents of *Allium sativum* (AS) Bulb Extracts (JEAS, EEAS and AEAS)

Phytochemical Components	Inference		
	JEAS	EEAS	AEAS
Alkaloids	Present	Present	Present
Anthraquinone	Absent	Absent	Absent
Carbohydrates	Present	Present	Present
Cardiac glycosides	Present	Present	Present
Fats and Oils	Present	Present	Present
Flavonoids	Present	Present	Present
Saponins	Present	Present	Present
Steroidal ring	Present	Present	Present
Terpenoids	Present	Present	Present
Tannins	Absent	Absent	Absent

Note: JEAS= Juice Extract, EEAS= Ethanolic Extract and AEAS=Aqueous Extract

Comment [v68]: how come the same constituents occur in all the threecheck???

Table 3: The Herbomineral Elements and Proximate Values of Juice Extract of *Allium sativum* (JEAS)

Macro/Micro (Zeolite) Elements	Proximate Values (mg)/600mg
Sodium	19.00
Potassium	154.00
Calcium	184.00
Magnesium	26.00
Phosphorus	390.00
Sulphur	450.00
Manganese	0.03
Silver	0.01
Iodine	0.05
Zinc	1.18
Selenium	0.014

Comment [v69]: Sp.check

Comment [v70]: Sp.check

3.3 Inhibitory Effect of *Allium sativum* Extracts on Growth of Clinical Bacterial

Isolates

The result obtained showed that *A. sativum* extract inhibited the growth of clinical bacterial isolates at 30mg/ml showed significant inhibitory $P < 0.05$ in in-vitro susceptibility test compared with that of standard Drug (control) (See Table 5).

Comment [v71]: delete

Comment [v72]: at $P < 0.05$

Comment [v73]: d

Table 4: Inhibitory Effect of *A. sativum* Extracts on the Growth of Clinical Bacterial Isolates

Test Exts.	Zone of growth inhibition (mm) of the extracts at drug concentrations (mg/ml)					Bacterial orgs
	10	15	20	25	5 (cefuroxime)	
JEAS	20.5±0.50	25.5±0.50	29.5±0.50	30.5±0.50 ^a	35	<i>Streptococcus pneumoniae</i>
EEAS	-	-	-	12.5±0.50	35	
AEAS	-	-	-	-	35	
JEAS	28.0±0.00	33.5±0.50 ^a	34.0±0.00 ^a	36.0±0.50 ^a	35	<i>Neisseria meningitides</i>
EEAS	-	-	14.5±0.50	17.5±0.50	35	
AEAS	-	-	-	-	35	
JEAS	-	-	-	12.5±0.50	20	<i>Klebsiella pneumoniae</i>
EEAS	-	-	-	-	20	
AEAS	-	-	-	-	20	
JEAS	23.0±0.00	26.5±0.50	28.0±0.00	30.5±0.50 ^a	35	<i>Haemophilus influenza</i>
EEAS	-	15.5±0.50	18.5±0.50	20.0±0.50	35	
AEAS	-	-	-	-	35	
JEAS	29.0±0.00	31.5±0.50 ^a	32.0±0.00 ^a	35.5±0.50 ^a	35	<i>Escherichia coli</i>
EEAS	-	-	-	25.5±0.50 ^a	35	
AEAS	15.5±0.50	16.0±0.00	19.0±0.00	20.5±0.50	35	

- = No zone of inhibition; Values are presented as Mean ± SEM, a=P < 0.05 compared to the standard drug (control) - One way ANOVA followed by Waller Duncan Post Hoc Test, Df=4 and N=15.

Comment [v74]: A note on means with different superscripts differ significantly




Figure 4: Minimum Inhibitory Concentration of Bulb extracts of *Allium sativum* JEAS=Juice Extract of *Allium sativum*, EEAS= Ethanolic Extract of *Allium sativum*, AEAS=Aqueous Extract of *Allium sativum*, S.P= *Streptococcus pneumoniae*, N.M= *Neisseria meningitides*, K.P= *Klebsiella pneumoniae*, H.I= *Haemophilus influenzae*, E. coli= *Escherichia coli*.

Comment [v75]: In figure, abbreviation can be given in legend instead of full text

Comment [v76]: Sp.check

3.4 Determination of Minimum Inhibitory and Minimum Bactericidal Concentrations (MIC & MBC)

The result obtained shown below revealed the efficacies of *A. sativum* extracts' inhibitory effects, and the rate of killing the pathogenic bacteria isolates (Table 5). The time of killing of bacteria can easily be deduce to be the ratio of MBC/ MIC = 1 or more than 1, the rate of killing (KR) will be rapid, and is the potential weapon for eradicating or combating pathogenic bacteria and thereby reduce hemorrhagic sepsis with systemic dysfunctions [26].

Comment [v77]: Sp.check

Comment [v78]: reducing

Table 5: Minimum Inhibitory and Minimum Bactericidal Concentrations (MIC & MBC) of Standard Drug (Cefuroxime axetil) (5mg/ml)

Comment [v79]: align centre

Clinical Isolates + Cefuroxime axetil	Concentration of standard drug (cefuroxime axetil) in (mg/ml)										MIC Values	MBC Values	MBC/MIC =KR
	5	2.5	1.25	0.63	0.313	0.157	0.76	0.04	0.02	0.01			
<i>S. pneumoniae</i>	-	-	-	-	-	-	-	-	+	+	0.04	0.04	1.00
<i>N. meningitides</i>	-	-	-	-	-	-	-	-	+	+	0.04	0.76	19.00
<i>K. pneumoniae</i>	-	-	-	-	-	-	+	+	+	+	0.157	0.63	4.01
<i>H. influenzae</i>	-	-	-	-	-	-	-	-	+	+	0.04	0.157	4.00
<i>E. coli</i>	-	-	-	-	-	-	-	-	-	+	0.02	0.04	2.00

- = No viable growth, + = viable growth

Table 6: Minimum Inhibitory and Minimum Bactericidal Concentrations (MIC & MBC) of the Bulb Extracts (Juice Extract, Ethanolic Extract and Aqueous Extract) of *A. sativum* (10mg/ml)

Comment [v80]: align centre

Clinical Isolates + Extracts		Concentration of <i>A. sativum</i> extracts in (mg/ml)										MIC Values	MBC Values	MBC/MIC=KR
		10	5	2.5	1.25	0.63	0.31	0.16	0.08	0.04	0.02			
<i>S. pneumoniae</i>	JEAS	-	-	-	+	+	+	+	+	+	+	2.5	NB	
	EEAS	-	+	+	+	+	+	+	+	+	+	10	NB	
	AEAS	+	+	+	+	+	+	+	+	+	+	NA	NA	
<i>N. meningitides</i>	JEAS	-	-	-	+	+	+	+	+	+	+	2.5	NB	
	EEAS	-	+	+	+	+	+	+	+	+	+	10	NB	
	AEAS	+	+	+	+	+	+	+	+	+	+	NA	NA	
<i>K. pneumoniae</i>	JEAS											NA	NA	
	EEAS											NA	NA	
	AEAS											NA	NA	
<i>H. influenzae</i>	JEAS	-	-	+	+	+	+	+	+	+	+	5	NB	
	EEAS	+	+	+	+	+	+	+	+	+	+	NA	NA	
	AEAS	+	+	+	+	+	+	+	+	+	+	NA	NA	
<i>E. coli</i>	JEAS	-	-	-	-	-	+	+	+	+	+	0.63	NB	
	EEAS	+	+	+	+	+	+	+	+	+	+	NA	NA	
	AEAS	-	-	+	+	+	+	+	+	+	+	5	NB	

- = No viable growth, + = Viable growth, NA=No activity, NB= Non Bactericidal

Comment [v81]: don't leave the column blank

Table 7: Minimum Inhibitory and Minimum Bactericidal Concentrations (MIC & MBC) of the Bulb Extracts (Juice Extract, Ethanollic Extract and Aqueous Extract) of *A. sativum* (15mg/ml)

Clinical Isolates + Extracts	Concentration of <i>Allium sativum</i> extracts in (mg/ml)										MIC Values	MBC Values	MBC/MIC=KR	
	15	7.5	3.8	1.88	0.94	0.47	0.23	0.12	0.06	0.03				
<i>S. pneumoniae</i>	JEAS	-	-	-	-	+	+	+	+	+	+	1.88	NB	
	EEAS	-	+	+	+	+	+	+	+	+	+	15	NB	
	AEAS	+	+	+	+	+	+	+	+	+	+	NA	NA	
<i>N. meningitides</i>	JEAS	-	-	-	-	-	+	+	+	+	+	0.94	NB	
	EEAS	-	+	+	+	+	+	+	+	+	+	15	NB	
	AEAS	+	+	+	+	+	+	+	+	+	+	NA	NA	
<i>K. pneumoniae</i>	JEAS											NA	NA	
	EEAS											NA	NA	
	AEAS											NA	NA	
<i>H. influenzae</i>	JEAS	-	-	-	+	+	+	+	+	+	+	3.8	NB	
	EEAS	+	+	+	+	+	+	+	+	+	+	NA	NA	
	AEAS	+	+	+	+	+	+	+	+	+	+	NA	NA	
<i>E. coli</i>	JEAS	-	-	-	-	-	-	-	+	+	+	0.23	NB	
	EEAS	+	+	+	+	+	+	+	+	+	+	NA	NA	
	AEAS	-	-	-	+	+	+	+	+	+	+	3.8	NB	

- = No viable growth, + = Viable growth, NA=No activity, NB= Non Bactericidal

Comment [v82]: align centre

Comment [v83]: don't leave the column blank

Table 8: Minimum Inhibitory and Minimum Bactericidal Concentrations (MIC & MBC) of the Bulb Extracts (Juice Extract, Ethanolic Extract and Aqueous Extract) of *A. sativum* (20mg/ml)

Clinical Isolates + Extracts		Concentration of <i>Allium sativum</i> extracts in (Mg/ml)										MIC Values	MBC Values	MBC/MIC=KR
		20	10	5	2.5	1.25	0.63	0.313	0.16	0.08	0.04			
<i>S. pneumoniae</i>	JEAS	-	-	-	-	-	+	+	+	+	+	1.25	2.5	2.00
	EEAS	+	+	+	+	+	+	+	+	+	+	NA	NA	
	AEAS	+	+	+	+	+	+	+	+	+	+	NA	NA	
<i>N. meningitides</i>	JEAS	-	-	-	-	-	-	+	+	+	+	0.63	1.25	2.00
	EEAS	-	+	+	+	+	+	+	+	+	+	20	NB	
	AEAS	+	+	+	+	+	+	+	+	+	+	NA	NA	
<i>K. pneumoniae</i>	JEAS											NA	NA	
	EEAS											NA	NA	
	AEAS											NA	NA	
<i>H. influenzae</i>	JEAS	-	-	-	-	+	+	+	+	+	+	2.5	5.0	2.00
	EEAS	-	-	-	+	+	+	+	+	+	+	5	NB	
	AEAS	+	+	+	+	+	+	+	+	+	+	NA	NA	
<i>E. coli</i>	JEAS	-	-	-	-	-	-	-	-	+	+	0.16	0.313	2.00
	EEAS	+	+	+	+	+	+	+	+	+	+	NA	NA	
	AEAS	-	-	-	-	+	+	+	+	+	+	2.5	NB	

Comment [v84]: m

Comment [v85]: don't leave empty.mention

- = No viable growth, + = Viable growth, NA=No activity, NB= Non Bactericidal

Table 9: Minimum inhibitory and Minimum Bactericidal Concentrations (MIC & MBC) of the Bulb Extracts (Juice Extract, Ethanolic Extract and Aqueous Extract) of *A. sativum* (25mg/ml)

Clinical Isolates + Extracts		Concentration of <i>Allium sativum</i> extracts in (mg/ml)										MIC Values	MBC Values	MBC/MIC=KR
		25	12.5	6.25	3.13	1.56	0.78	0.39	0.20	0.10	0.05			
<i>S. Pneumoniae</i>	JEAS	-	-	-	-	-	-	+	+	+	+	0.78	1.56	2.00
	EEAS	-	+	+	+	+	+	+	+	+	+	25	NB	
	AEAS	+	+	+	+	+	+	+	+	+	+	NA	NA	
<i>N. meningitides</i>	JEAS	-	-	-	-	-	-	+	+	+	+	0.39	0.78	2.00
	EEAS	-	+	+	+	+	+	+	+	+	+	25	NB	
	AEAS	+	+	+	+	+	+	+	+	+	+	NA	NA	
<i>K. pneumoniae</i>	JEAS	-	-	+	+	+	+	+	+	+	+	12.5	NB	
	EEAS											NA	NA	
	AEAS											NA	NA	
<i>H. influenzae</i>	JEAS	-	-	-	-	-	+	+	+	+	+	1.56	3.13	2.00
	EEAS	-	-	-	-	+	+	+	+	+	+	3.13	NB	
	AEAS	+	+	+	+	+	+	+	+	+	+	NA	NA	
<i>E. coli</i>	JEAS	-	-	-	-	-	-	-	-	-	+	0.10	0.20	2.00
	EEAS	-	-	-	-	-	+	+	+	+	+	1.56	NB	
	AEAS	-	-	-	-	-	-	-	+	+	+	0.39	NB	

- = No viable growth, + = Viable growth, NA=No activity, NB= Non Bactericidal

CONCLUSION

The results of this study showed that bulbs extracts of *A. sativum* possessed active phytochemical with organosulphur volatile oil substances, which can be used as novel antibacterial agents. It exhibited potent antihemorrhagic and antibacterial activities which fight microbes due to its phytoconstituents, antioxidant properties, dietary medicines and coupled with zeolite herbominerals present. However, *A. sativum* extract can interact with the entire systems. It plays a vital role in human health, human nutrition as integrative medicine against bacterial meningitis pathogens. Therefore, there is need for further investigations in terms of toxicological studies and purification of bioactive components with view to using *A. sativum* in novel drug development. This study has justified and validated the folkloric usage of *A. sativum* as local anti-infectious therapeutics in any bacterial bioterrorism (as natural disaster).

Comment [v86]: italics

Comment [v87]: italics

COMPETING INTERESTS

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

REFERENCES

1. World Health Organization, (2009). Bacterial Meningitis (BM) in Sub-Saharan Africa, *Wikipedia*.
2. World Health Organization, (2007). WHO-CDS-EPR, Malnutrition Links Bacterial Meningitis, *Wikipedia*.
3. Kokori B. T., Abdullahi A. A. and Abdullahi A. S. (2019). Studies on Phytochemical, Nutraceutical Profiles and Potential Medicinal Values of *Allium sativum* Linn (Lilliaceae)

Comment [v88]: follow a uniform style

Comment [v89]:

Comment [v90]: Enter the website address

Comment [v91]: address

- on Bacterial Meningitis. *International Neuropsychiatric Disease Journal* 13(2): 1-15;
Article no.INDJ.51609, ISSN: 2321-7235, NLM ID: 101632319
4. Nikaido, H. and Pagès, J. M. (2012). Broad-Specificity Efflux Pumps and Their Role in Multidrug Resistance of Gram-Negative Bacteria. *FEMS Microbiology Reviews*, 36: 340–363.
 5. United Nations (2012). Malnutrition Linked to Bacterial Meningitis Scourge in Africa, <http://www.unitednation-malnutritionlinkmeningitis>, Health Experts.
 6. Onipede, A. O., Onayade, A. A., Elusanya, J. B. E., Obiajunwa, P. O., Ogundare, E. O. O., Olaniran, O. O. (2009). Invasive Bacterial Isolates from Children with Severe Infections in a Nigerian Hospital *Journal of Infectious Diseases*; 3(6):429-436.
 7. Okoko, B. J., Idoko, O. T., Adegbola, R. A. (2009). Prospects and Challenges with Introduction of a Monovalent Meningococcal Conjugate Vaccine in Africa. *Medical Research Council Laboratories*. 27 (14): 2023-9.
 8. IRIN Africa News (2009): *African Meningitis belt*, [Wikipedia](#).
 9. Stephens, D. S., Greenwood, B. and Brandtzaeg, P. (2007). Epidemic Meningitis, Meningococcal and *Neisseria meningitidis*. *Lancet British Journal of Infectious Disease*; 369: 2196-2210.
 10. Hackam, D. G., Wooward, M., Newby, L. K., Bhatt, D. L., Shao, M., Snuth, E. E., Donner, A., Mamdani, M, and Donketis, J.D, (2011), European publication, Pathogenesis and Pathophysiology of Hemorrhage, Bacterial Endotoxin inVital Organs and Cerebral Hemorrhagic Stroke. *European Journal of Neuroscience*124 (20):2233-42.
 11. Corey, S. J., Burkhardt, A. L., Bolen, J. B. and Geahlen, R. L. (2011). Research Laboratory, Clotting time (Coagulation). *Journal of the American Medical Association*; *JAMA* 11(2) 153-160.

Comment [v92]: comma

Comment [v93]: bold

Comment [v94]: check

Comment [v95]: . and

Comment [v96]: fullstop check

Comment [v97]: check

Comment [v98]: bold check

Comment [v99]: bold

Comment [v100]: O.T.and

Comment [v101]: Web address and line spacing

Comment [v102]: Look for the common style

Comment [v103]: check

Comment [v104]: comma

Comment [v105]: bold

Comment [v106]: check

Comment [v107]: bold

12. Tebruegge, M. and Curtis, N. (2008). Epidemiology, Etiology, Pathogenesis and Diagnosis of Recurrent Bacterial Meningitis. *Clinical Microbiology Reviews* 21(3): 519-37.
13. Abdullahi Attah A., Ayodeji, O., Donatus Teru, G. A., & Bajeh Tijani, K. (2019). Studies on the Phytochemical Compounds in the Ethanolic Leaf Extract (ELE), Ethanolic Bark Extract (EBE) and Ethanolic Root Extract (ERE) of *Bridelia ferruginea* Benth (Euphorbiaceae). *Asian Journal of Biochemistry, Genetics and Molecular Biology*, 2(4), 1-8.
14. Ncube, N., S, Afolayan, A. J and Okoh, A. J. (2008). Assessment Techniques of Antimicrobial properties of Natural Compounds of Plant Origin Current Method and Future Trends *African Journal of Biotechnology* 7:1997-1806.
15. Rader Heidi and McGuinness Julianne (2019). Growing Garlic in Alaska. University of Alaska Fairbanks. Retrieved September 24, 2019.
16. World Health Organization (2010). Traditional Medicine, *Wikipedia*.
17. Oduye, T. (2011): National Health Holistic Consultancy: Garlic Constituents and its Organosulfur are Good Interactive Herbs for Mitigating Diseases, Amelias Organics Limited; The Nation, Thursday, April 28, pg 25.
18. Fattouch, S., Imen, B., Belguitt, H. (2007). Preparation of aqueous garlic extract (AGE), Morphological and Biochemical changes of *Salmonella hadar* and fungi. *International Journal of Morphology* 44(62):231-38.
19. Monica Cheesebrough (2018). *District Laboratory Practice in Tropical Countries*. Part 2, Pg 62-71 and 116-129.

Comment [v108]: comma

Comment [v109]: bold

Comment [v110]: avoid symbols

Comment [v111]: bold

Comment [v112]: check

Comment [v113]: spacing

Comment [v114]: spacing

Comment [v115]: comma

Comment [v116]: spacing

Comment [v117]: address

Comment [v118]: check

Comment [v119]: . and

Comment [v120]: comma

Comment [v121]: bold

20. Sharma C, Aneja K. R. (2012). Agar well diffusion method, Antimicrobial potential of *Terminalia chebula* Retz fruit extracts, *World Journal of Otorhinolaryncology*, 78:42- 9.
21. Naturopathy (2010). Wikiedia. Content is available under CC BY-SA.
22. World Health Organization-Biological Standardization, (2011). WHO Expert Committee on Biological Standardization-Bacterial Suspension, *Wikipedia*.
23. Walker R. and Whittlesea C. (2008). *Clinical Pharmacy and Therapeutics*, 4th edition pg 531-33, Churchill.
24. Sen Saikat, De Biplab, Devanna N., Chakraborty Raja (2012). *Vitro* Antihelmintic (Antitrematodal) Activity in Ruminant from Plant Derived Compounds Antioxidant, Cysteine Proteases and Diallyl Disulphide. *Pubmed*, www.science.gov/garlic/vitro.
25. Abdullahi Attah Alfa and Abayomi Ajayi (2014). Bacteriological and Physiochemical Analysis of Spring Water: Case study of Osan-Ekiti, Nigeria. *Malaya Journal of Biosciences*. 1(3): 160-166, ISSN 2348-6236 print/2348-3075 online
26. Pilkhwal, S., Sharma, S., Kuhad, A., Mukesh, L. S. (2010). The direct effects of bacterial endotoxin and circulation after killing bacteria. *International Journal of Phytomedicine*; 2(2010):47-5148.

Comment [v122]: . and

Comment [v123]: spelling

Comment [v124]: address

Comment [v125]: *In vitro*

Comment [v126]: bold

Comment [v127]: . and

Comment [v128]: check