

**Grape seed proanthocyanidin alleviates toxicity, oxidative stress,  
injury and apoptosis involved in liver dysfunction of hyperthyroid  
mice**

**ABSTRACT**

**Objectives:** Thyroid hormones control the basal metabolic pace of hepatocytes, and can make oxidative harm hepatic frameworks. The reason for this investigation was to investigate how hyperthyroidism-prompted liver danger, oxidative pressure and apoptosis changes could be alleviated with Grape seed proanthocyanidin separate (GSPE). This investigation assessed some biochemical, histological and immunohistochemical changes in post pubertal hyperthyroidism and its effect on liver capacity and structures. Notwithstanding the enhancing job of Grape seed proanthocyanidin remove (GSPE) supplementation was analyzed.

**Materials and Methods:** Fifty male Swiss albino mice were randomly divided into 5 groups (G1, Control; G2, GSPE; G3, Eltroxin-induced hyperthyroid mice; G4, Post treated hyperthyroid with GSPE; G5, Self-treated hyperthyroid mice).

**Results:** Our results revealed that, a significant increase in serum T3, T4, ALT, AST, ALP, liver MDA, P53 levels, injury and P53 expression in hyperthyroid mice when compared to control and GSPE. In contrast; serum albumen, liver catalase, GSH, SOD and Bcl2 were decrease in hyperthyroid mice. Treatment of hyperthyroid mice with GSPE advantages in improving the adverse effect of hyperthyroidism and moreover the histopathological and P53 expression result approves this finding.

**Conclusions:** GSPE can used in hyperthyroidism treatment with propylthiouracil or carbimazole or methimazole replacement therapy.

*Key words: Hyperthyroidism, Grape seed proanthocyanidin, hepatic dysfunction, Oxidative stress, liver injury and PCNA.*

**1. INTRODUCTION**

Hyperthyroidism is one of the most open endocrine issues found in medicinal practice, elevated levels in serum liver and kidney chemicals are regular going with hazard elements of hyperthyroidism [1]. Hyperthyroidism is portrayed by expanded discharge of thyroid hormones tri-iodothyronin as well as thyroxin [2,3]. Thyroxine and tri-iodothyronine are thyroid hormones that basic for typical organ improvement and metabolic capacities [4-10]. Thyroid hormones modify the basal metabolic pace of hepatocytes and along these lines control hepatic capacity [11,12]. The liver is biggest organ in the body and it assumes a significant job in our digestion where it is the principle site for lipid digestion, and the thyroid hormones have a basic impact in hepatic lipid homeostasis [13,14]. Hence it isn't astounding that expanded digestion in light of hyperthyroidism can make hepatic brokenness and oxidative harm hepatic frameworks [15]. Several plant extracts have significant antioxidant activity; one of this is a grape seeds that is rich sources for proanthocyanidins [16-19]. Proanthocyanidins are consists of many polyphenolic compounds and it have become of high importance because of their biological

27 properties (anti-oxidant, anti-inflammatory and anti-cancerigen) and their protective effects by  
28 reducing mitochondria damage and inhibiting cell apoptosis [20,21]. Grape seed  
29 proanthocyanidin extract (GSPE) is believed to protect against reactive oxygen species (ROS)-  
30 mediated myocardial ischemia/reperfusion injury and apoptosis [22,23]. GSPE have free radical  
31 searching properties are more noteworthy than well known cancer prevention agents, for  
32 example, nutrients C and E, and it demonstrates a better capacity than secure cells against lipid  
33 peroxidation and DNA fracture [24]. Subsequently; the present investigation was intended to  
34 explain the conceivable enhancing impacts of GSPE in improving hepatic lethality, apoptosis,  
35 oxidative pressure, damage, and apoptosis adjustments against Eltroxin instigated  
36 hyperthyroidism in male mice.

## 37 **2.0 MATERIALS AND METHODS**

### 38 **2.1. Chemicals and drug:**

39 ELTROXIN (Thyroxine 100 mcg; 100 Tablet) was obtained from Mercury Pharma Group  
40 Limited, Capital House, London EC4N 7BL, UK.

41 Grape seed proanthocyanidin (GSPE; 200mg; Pharco Pharmaceuticals Co. Alexandria, Egypt)  
42 with 95.0–96.29% Purity of Oligomeric Proanthocyanidins.

### 43 **2.2. Animals**

44 A total of 50 male Swiss albino mice (*Mus musculus*) 6–8 weeks old, weighing  $25 \pm 2$ g, supplied  
45 from the animal house of the King Saud University, Riyadh, Saudi Arabia. Animals were  
46 provided standard mice feed and water ad libitum. The experimental protocol was approved by  
47 Local Ethics Committee and Animals Research, King Saud University, Riyadh, Saudi Arabia.

### 48 **2.3. Experimental design and mice groups**

49 The mice were equally divided into 5 groups with ten rats each.

50 G1: Control group in which mice did not received any treatment.

51 G2: GSPE; mice received GSPE (50mg/Kg/day) only for three weeks orally by a stomach tube  
52 [17].

53 G3: Hyperthyroid; mice received 100  $\mu$ g/Kg Eltroxin in drinking water for 3 weeks to induce the  
54 hyperthyroid state [2].

55 G4: Post treated hyperthyroid; mice received 100  $\mu$ g/Kg Eltroxin in drinking water for 3 weeks  
56 and then treated with GSPE for another 3 weeks (from 4<sup>th</sup> week to 6<sup>th</sup> week).

57 G5: Self-treated hyperthyroid; mice received 100  $\mu$ g/Kg Eltroxin in drinking water for 3 weeks  
58 and then mice did not received any treatment for another 3 weeks (from 4<sup>th</sup> week to 6<sup>th</sup> week)  
59 according to Tousson et al. [13].

60 At the end of the experiment period, blood samples were individually collected from the eyes by  
61 retro-orbital puncture using blood capillary tubes in non-heparinized glass tubes under mild  
62 ether anesthesia. Blood samples were centrifuged to obtain serum used for detection of thyroid  
63 hormones and liver functions.

### 64 **2.4. Determination of serum thyroid hormones:**

65 Serum was used to determine the triiodothyronine ( $T_3$ ) according to Thakur et al. [25]; thyroxine  
66 ( $T_4$ ) according to Maes et al. [26] and thyrotropin (TSH) according to Mandel et al. [27].

### 67 **2.5. Determination of serum liver enzymes**

68 Serum was analyzed to determine aspartate transaminase (AST) and alanine transaminase  
69 (ALT) activities using commercial kit (Humann, Germany) according to the method of Schumann  
70 and Klauke [28]; serum albumin levels by using commercial kit (Diamond, Egypt) according to  
71 Dumas et al. [29]; serum alkaline phosphatase (ALP) activity in serum was detected by using  
72 commercial kit (Humann, Germany) according to Moss and Henderson [30].

### 73 **2.6. Preparation of tissue homogenates**

74 Liver from different groups were removed, weighed and stored at -20°C then 10%W/V  
75 homogenate was prepared by ground 0.3 g of tissue in 3ml of saline, liver homogenates were  
76 used to estimate the oxidative stress parameters.

### 77 **2.7. Enzymatic and non-enzymatic antioxidant assays**

78 Determination of malondialdehyde (MDA) levels in homogenate was assayed according to  
79 Mesbah et al. [31], catalase (CAT) was detected after Saggu et al. [32], superoxide dismutase  
80 (SOD) activity was detected after Misra and Fridovich [33] and reduced glutathione (GSH)  
81 content was detected after Beutler et al. [34].

### 82 **2.8. Determination of P53 and Bcl2 Protein Levels**

83 P53 and Bcl2 protein levels were determine after the method of Tousson et al. [35].

### 84 **2.9. Histological preparation**

85 The changes in the liver structure were prepared according to the method of Tousson [36].

### 86 **2.10. Detection of apoptotic P53 expressions**

87 P53 immunoreactivity were detected after the method to Tousson et al. [8,9].


### 88 **2.11. Statistical Analysis**

89 Information were communicated as mean qualities  $\pm$  SE and factual examination was performed  
90 utilizing one route ANOVA to survey huge contrasts among treatment groups. The measure for  
91 factual essentialness was set at  $p < 0.05$  for the biochemical information. Every single  
92 measurable investigation were performed utilizing SPSS factual form 21 programming bundle  
93 (SPSS® Inc., USA).

## 94 **3.0 RESULTS**

### 95 **3.1. Induction of hyperthyroid mice**

96 Figures (1A-1C) showed that; the serum T<sub>3</sub> and T<sub>4</sub> levels in hyperthyroid and self-recovered  
97 hyperthyroid mice were significantly higher as compared to control; In contrast; serum TSH  
98 levels in hyperthyroid and self-recovered hyperthyroid mice were significantly lower as  
99 compared to control (Figure 1A). The treatment of hyperthyroid mice with GSPE revealed a  
100 significant decrease in T<sub>3</sub> and T<sub>4</sub> levels as compared with hyperthyroid mice (Figures 1B&1C).  
101 Also; Treatment of hyperthyroid mice with GSPE revealed a significant increase in TSH levels  
102 as compared with hyperthyroid mice group.



103 **Figure 1:** Changes in thyrotropin (TSH; µIU/ml), thyroxine (T4; ng/dl) and triiodothyronine (T3;  
 104 ng/dl) levels in different experimental groups. Data are expressed as mean ± standard error.  
 105 G1, Control; G2, GSPE; G3, Hyperthyroid; G4, post-treated hyperthyroid with GSPE; G5, Self-  
 106 treated hyperthyroid. # Significantly different from G3, \*Significantly different from G1.  
 107  
 108

109 **3.2. Changes in liver functions in different groups**

110 Table (1) showed that; a significance increase in serum ALT, AST and ALP levels in  
 111 hyperthyroid mice and self-recovered hyperthyroid mice groups as compared with control and  
 112 treated mice with GSPE. In contrast; significance decreases in serum albumin in hyperthyroid  
 113 mice and self-recovered hyperthyroid mice as compared to control and treated mice with GSPE.  
 114 On the other hand; treatment of hyperthyroid mice with GSPE revealed a significant decrease in  
 115 ALT, AST and ALP levels as compared with hyperthyroid mice group. In addition to the  
 116 treatment of hyperthyroid mice with GSPE revealed a significant increase in albumin levels as  
 117 compared with hyperthyroid group (Table 1).

118  
119  
120

**Table 1:** Changes in liver functions (ALT, AST, ALP, and Alb) parameters.

	<b>G1</b>	<b>G2</b>	<b>G3</b>	<b>G4</b>	<b>G5</b>
<b>ALT (U/l)</b>	37.5 <sup>#</sup> ± 1.81	31.1 <sup>#</sup> ± 2.15	66.3* ± 4.58	40.6 <sup>#*</sup> ± 3.19	58.4* ± 4.43
<b>AST (U/l)</b>	136.2 <sup>#</sup> ± 8.56	130.5 <sup>#</sup> ± 10.18	171.0* ± 10.45	134.52 <sup>#</sup> ± 11.22	165.1* ± 10.88
<b>ALP (U/l)</b>	161.9 <sup>#</sup> ± 11.25	147.4 <sup>#</sup> ± 8.69	193.1* ± 10.33	169.8 <sup>#*</sup> ± 11.49	180.2* ± 12.05
<b>Alb (g/dl)</b>	4.55 <sup>#</sup> ± 0.216	4.78 <sup>#</sup> ± 0.185	3.35* ± 0.164	4.60 <sup>#</sup> ± 0.228	3.40* ± 0.157

121  
122  
123  
124  
125

Data are expressed as mean ± standard error. G1, Control; G2, GSPE; G3, Hyperthyroid; G4, post-treated hyperthyroid with GSPE; G5, Self-treated hyperthyroid. <sup>#</sup> Significantly different from G3, \*Significantly different from G1.

126

### 3.3. Lipid peroxidation and reduced glutathione content

127  
128  
129  
130  
131  
132

Figure (2A) shows a significant ( $P<0.05$ ) increase in the liver MDA levels in hyperthyroid and self-treated hyperthyroid mice as compared with control mice. Our data showed a significant ( $P<0.05$ ) decrease in GSH levels in liver of hyperthyroid and self-treated hyperthyroid mice as compared with control mice (Figure 2B). On the other hand, hyperthyroid mice treated with GSPE showed significant depletion in liver MDA and alleviation in GSH levels as compared with hyperthyroid mice (Figures 2A&2B).

133

### 3.4. Antioxidant enzyme activities

134  
135  
136  
137  
138  
139

A significantly ( $P< 0.05$ ) decreased in the activities of SOD and CAT was detected in liver homogenate of hyperthyroid and self-treated hyperthyroid mice (Figures 2C&2D). Hyperthyroid mice treated with GSPE showed significant alleviation in the antioxidant enzyme activities as compared with hyperthyroid and self-treated hyperthyroid mice. Moreover, antioxidant enzyme activities were significant increase in mice treated with GSPE alone as compared to control (Figures 2C&2D).

140

### 3.5. Hyperthyroidism induced apoptosis in mice liver

141  
142  
143  
144  
145  
146

Figures (3) show the concentration of P53 and Bcl2 levels in mice liver tissues. The levels of P53 were significant ( $P< 0.05$ ) increase in hyperthyroid and self-treated hyperthyroid mice when compared with control (Figures 3A&3B).). While Bcl2 levels were significant ( $P< 0.05$ ) depletion in hyperthyroid and self-treated hyperthyroid mice when compared with control. Treatment of hyperthyroid mice with GSPE improved this alternation of P53 and Bcl2 concentrations in liver tissues (Figure 3).

147

### 3.6. Liver histopathology


148  
149  
150  
151  
152  
153  
154

Liver sections on control and treated mice with GSPE exhibits normal histological structure of hepatocytes (Figs 4A&4B). Liver sections on hyperthyroid mice showed marked vacuolated hepatocytes, marked cellular infiltrations, inflammation, moderate degeneration with necrotic area and mild congestion in central veins and portal vein, surrounded by leucocytic infiltrations (Figures 4C&4D). Liver sections in post treated hyperthyroid mice with GSPE revealed a few vacuolated hepatocytes, moderate cellular infiltrations, and cytoplasmic vacuolations observed when compared to hyperthyroid or self-treated hyperthyroid mice (Figure 4E). In contrast; self-


155 treated hyperthyroid mice showed marked vacuolated hepatocytes, marked cellular infiltrations  
156 and marked degeneration with necrotic area (Figure 4F).

### 157 3.7. Changes in P53 expression

158 Liver sections in control group and GSPE group showed negative expression of P53 (Figure  
159 5A&5B). Strong P53 expression were detected in liver sections of hyperthyroid and self-treated  
160 hyperthyroid mice (Figures 5C&5D&5F). The intensity of P53 expression in hyperthyroid group  
161 was increased when compared with control group. Mild positive expressions for P53 were  
162 observed in liver sections in post treated hyperthyroid mice with GSPE (Figure 5E).



163  
164  
165 **Figure 2:** Changes in MDA, GSH, and catalase and SOD levels in liver tissues in different  
166 experimental groups. Data are expressed as mean  $\pm$  standard error. G1, Control; G2, GSPE;  
167 G3, Hyperthyroid; G4, post-treated hyperthyroid with GSPE; G5, Self-treated hyperthyroid. #  
168 Significantly different from G3, \*Significantly different from G1.




170 **Figure 3:** The changes in P53 and Bcl2 levels in mice liver tissues in all studied groups. Data  
 171 are expressed as mean  $\pm$  standard error. G1, Control; G2, GSPE; G3, Hyperthyroid; G4, post-  
 172 treated hyperthyroid with GSPE; G5, Self-treated hyperthyroid. # Significantly different from G3,  
 173 \*Significantly different from G1.  
 174  
 175

#### 176 4. DISCUSSION

177 The current study has been represented to examine the effect of a hyperthyroidism status on  
 178 biochemical parameters, oxidative stress parameters, apoptosis, histological and P53  
 179 immunohistochemical alterations in liver tissues. Our results revealed elevation in serum  $T_3$ ,  $T_4$   
 180 and depression TSH in mice receiving Eltroxin indicating the hyperthyroid state. This finding is  
 181 compatible with other studies that used L-thyroxin as an anti-thyroid drug for induction of  
 182 hyperthyroidism [2-4]. In the current study, depletion in the level of  $T_3$  and  $T_4$  and elevation in  
 183 TSH were detected in post treated hyperthyroid mice with GSPE when compared with  
 184 hyperthyroid mice. The current outcome agrees with studies of Shibutani et al. [37]; and Beltagy  
 185 et al. [2].

186 Liver assumes a significant job in thyroid hormones digestion, and correspondingly, thyroid  
 187 hormones direct hepatic capacity and bilirubin digestion, in this manner as anyone might expect,  
 188 disorders of either organ can possibly influence capacity of the other [12, 13]. In the current  
 189 results revealed a significance increase in serum ALT, AST, ALP and significance depletion in  
 190 albumen levels in hyperthyroid mice confirmed that hyperthyroidism induced dysfunction in mice


191 liver. Giannini et al. [38]; Hull et al. [39] fined that; a significant elevation in serum direct bilirubin,  
 192 ALT, AST and ALP in hyperthyroidism. The mechanism of the elevation in serum AST, ALT and  
 193 ALP activities appears to be relative hypoxia in periventricular regions of the liver [40]. Wang et  
 194 al. [41] and Upadhyay *et al.* [42] who fined that; the elevation in the levels of T3 induces  
 195 hepatocytes apoptosis take place through the mitochondrial dependent pathway activation. In  
 196 the current results; the treatment of hyperthyroid mice with GSPE modulates liver function  
 197 parameters as compared with self-treated hyperthyroidism. Current results decide with El-Sayed  
 198 et al. [43] who indicated the liver protective effects of proanthocyanidin on acetaminophen. Also;  
 199 Kandemir et al. [44] who studies that; the protective effects of grape seed extract against  
 200 Cisplatin induced liver toxicity in rabbits.  
 201



202  
 203 **Figures 4A-4F:** Photomicrographs of mice liver sections in the different experimental groups  
 204 stained with Haematoxylin & Eosin. **A&B:** Normal structure of hepatocytes (Hp) and central vein  
 205 (cv) in control and GSPE groups revealed **C&D:** Marked vacuolated hepatocytes, marked  
 206 necrosis (star), marked cellular infiltrations (White arrows), and moderate degeneration with  
 207 necrotic area (Black arrows) in hyperthyroid mice revealed. **E:** Mild vacuolated hepatocytes



208 (Hp), diffuse Kupffer cells proliferation in between the some hepatocytes and moderate cellular  
 209 infiltrations (White arrows) in post treated hyperthyroid mice with GSPE. **F:** Marked vacuolated  
 210 hepatocytes, marked cellular infiltrations (White arrows) and marked degeneration with necrotic  
 211 area (star) in self-treated hyperthyroid mice.  
 212



213  
 214 **Figures 5A-5F:** Photomicrographs of mice liver sections marked with apoptotic P53 protein.  
 215 **A&B:** Negative or faint positive reactions (arrows) in hepatocyte nuclei in hepatocytes in control  
 216 and GSPE. **C&D:** Strong positive reactions (arrows) in hyperthyroid. **E:** Moderate to strong  
 217 positive reactions (arrows) in self-treated hyperthyroid mice. **F:** Mild positive reactions (arrows)  
 218 in post treated hyperthyroid mice with GSPE.

219

220 Changes in thyroid hormone levels are known to tweak elements of numerous tissues by  
221 adjusting their cancer prevention agent resistances [45]. Hyperthyroidism is distinguished to  
222 improve ROS age and it instigates oxidative worry in liver tissue through upsetting the  
223 endogenous star oxidant/against oxidant balance [46]. In the present study, treatment of mice  
224 with Eltroxin induced hyperthyroidism that induced oxidative stress which play an important role  
225 in the damage of DNA [47]. In the current study; elevation in liver MDA and depletion in liver  
226 GSH, catalase and SOD levels in hyperthyroid mice. On the other hand, hyperthyroid mice  
227 treated with GSPE showed significant depletion in liver MDA and alleviation in GSH, catalase  
228 and SOD levels as compared with hyperthyroid and self-treated hyperthyroid mice. Asayama  
229 and Kato [48] reported that; the damage in MDA was increase in some organs in rats.  
230 Antioxidants contained in the red grape seed extract are able to inactivate superoxide anions  
231 and prevent lipid peroxidation [49]. Our results confirm GSPE as a potent antioxidant that can  
232 reduce lipid peroxidation in liver tissues. Current results agree with Davies [50] who fined that  
233 hyperthyroidism tends to increase catalase activity [51].

234 Reduced glutathione (GSH) as an oxidative stress marker showed a significant decrease in  
235 hyperthyroid rats. This decrease was improved by treatment with GSPE. Administration of  
236 GSPE exhibit an increase in antioxidant enzyme activities in hyperthyroid which might be due to  
237 its ability to reduce the accumulation of free radical generation [52]. Long et al. [52] reported  
238 that; GSPE showed an increase in GSH, catalase and SOD levels that clearly suggest their  
239 antioxidant property. GSPE has scavenging character and it can effectively inhibit liver toxicity  
240 induced by hyperthyroidism and this is may be attributed to its antioxidant properties.

241 Apoptosis is a urgent cell action in the conduct of mammalian cells in a wide scope of  
242 pathophysiological conditions. Apoptosis of individual cells may introduce a defensive system  
243 against neoplastic improvement in the life form by wiping out hereditarily harmed cells [41]. In  
244 the present examination; a critical increment in P53 and lessening in Bcl2 levels in hyperthyroid  
245 mice and the treatment of hyperthyroid mice with GSPE decline in P53 and increment in Bcl2  
246 levels. Our immunohistochemical results for P53 in liver additionally affirmed these outcomes, in  
247 this way, our outcomes uncover the plausibility of the apoptosis event after Eltroxin  
248 organization. Our outcomes concur with Kumar et al. [47] who announced that; hyperthyroidism  
249 incites apoptosis in rodent liver through enactment of death receptor-interceded pathways.  
250 Likewise; Tousson et al. [8,9] accomplishment to discover backwards connection somewhere in  
251 the range of's P53 and Bcl2. Our outcomes not concur with Diebold et al [53] who neglected to  
252 uncover any connection somewhere in the range of' P53 and Bcl2.

253 In the current study; many signs of pathological alterations were observed in liver sections in  
254 hyperthyroid mice as marked vacuolated hepatocytes, marked necrosis, marked cellular  
255 infiltrations, and moderate degeneration with necrotic area, this outcomes perhaps were a result  
256 of oxidative pressure brought about by Eltroxin that expansion digestion and free radicals with  
257 the goal that the body's barriers of cell reinforcements can't avoid free radicals that harm liver  
258 tissue. Additionally, in our investigation on hyperthyroidism actuated in mice showed that  
259 Eltroxin hormone animates the quality answerable for customized passing in liver cells and it  
260 was seen in the histological examination a distinction in cell shading and degeneration in the  
261 cytoplasm nature just as hyperthyroidism causing ischemia in the liver [24,47,54]. In the present  
262 investigation GSPE supplementation improvement of liver harm initiated by Eltroxin, this may  
263 come back to the job of viability grape seed concentrate mixes against oxidative pressure,  
264 irritation and customized cell passing opposition [43].

265 In conclusion, current work confirmed that; hyperthyroidism induced liver toxicity, oxidative  
266 stress, and apoptosis also the treatment with GSPE improved these alterations in liver tissues.

267  
268  
269  
270  
271  
272  
273  
274  
275  
  
276  
277  
  
278  
279  
280  
  
281  
282  
283  
  
284  
285  
286  
  
287  
288  
289  
  
290  
291  
292  
  
293  
294  
  
295  
296  
297  
  
298  
299  
300  
  
301  
302  
  
303  
304  
305  
  
306  
307  
308  
  
309  
310  
311  
  
312  
313

## CONSENT

It is not applicable.

## ETHICAL APPROVAL

It is not applicable.

## REFERENCES

1. Joeimon JL, Mohanraj K, Karthikeyan R, et al. Thyroid dysfunction in patients with liver cirrhosis. *Journal of Dental and Medical Sciences* 2017; 16(4):18-22.
2. Beltagy D, Mohamed T, El Said A, Tousson E. The cardioprotective effects of L-carnitine on rat cardiac injury, apoptosis, and oxidative stress caused by amethopterin. *Environ Sci Pollut Res* 2016; 23: 20600–20608.
3. Salama A, Tousson E, Ebrahim W, Elwan M. Ameliorating Role of Folic Acid in Eltroxine Induced Hyperthyroid and Oxidative Stress in Rat Cortex, Hypothalamus and Hippocampus. *Alexandria Journal of Veterinary Sciences* 2017; 55 (1): 210-216.
4. Tousson E, Ali EM, Ibrahim W, Mansour MA. Proliferating Cell Nuclear Antigen as a Molecular Biomarker for Spermatogenesis in PTU-Induced Hypothyroidism of Rats. *Reprod Sci* 2011; 18: 679-686.
5. Tousson E, Ali EM, Ibrahim W, Mansour MA. Treatment with folic acid ameliorated the histopathological alterations caused by propylthiouracil-induced hypothyroid rat testes. *Toxicology and Industrial Health* 2012a; 28: 566–576
6. Tousson E, Ali EM, Ibrahim W, Ashraf RM. Histopathological and immunohistochemical alterations in rat heart after thyroidectomy and the role of hemin and ketoconazole in treatment. *Biomedicine & Pharmacotherapy* 2012b; 66: 627–632.
7. Tousson E, Hafez E, Massoud A, Sweef O, Atta N. Protective role of folic acid in thyroxine-induced cardiac hypertrophy in hyperthyroid rat. *Biomedicine & Aging Pathology* 2013; 3: 89–95.
8. Tousson E, Hafez E, Zaki S, Gad A. P53, Bcl-2 and CD68 expression in response to amethopterin-induced lung injury and ameliorating role of l-carnitine. *Biomedicine & Pharmacotherapy* 2014; 8(5): 631–639.
9. Tousson E, Hafez E, Zaki S, Gad A. The cardioprotective effects of L-carnitine on rat cardiac injury, apoptosis, and oxidative stress caused by amethopterin. *Environ Sci Pollut Res* 2016; 23:20600–20608
10. Kim, D. et al. Subclinical hypothyroidism and low-normal thyroid function are associated with nonalcoholic steatohepatitis and fibrosis. *Clin. Gastroenterol. Hepatol.*, 2018; 16: 123–131.
11. Aydemir S, Bayraktaroglu T, Demircan N, Sert M, Açikgoz S, Tekin IO, et al . Effect of hyperthyroidism and propylthiouracil treatment on liver biochemical tests. *Int J Clin Pract* 2005; 59(11): 1304-1308.
12. Mansourian AR. A review of literatures on the adverse effects of thyroid abnormalities and liver disorders: An overview on liver dysfunction and hyperthyroidism. *P J Biological Science* 2013; 16: 1641-1652.
13. Bolkiny Y, Tousson E, El-Atrsh A, Akela M, Farg E. Costus Root Extract Alleviates Blood Biochemical Derangements of Experimentally-Induced Hypo-and Hyperthyroidism in Mice. *Annual Research & Review in Biology* 2019; 31(5):1-0.
14. Sinha RA, Yen PM. Thyroid hormone-mediated autophagy and mitochondrial turnover in NAFLD. *Cell Biosci* 2016; 6: 46.

- 314 15. Khemichian S, Ling Fong T. Hepatic Dysfunction in Hyperthyroidism. *Gastro enterol Hepatol*  
315 2011;7(5): 337–339.
- 316 16. Bagchi D, Swaroop A, Preuss HG, Bagchi M. Free radical scavenging, antioxidant and cancer  
317 chemoprevention by grape seed proanthocyanidin: an overview. *Mutat Res* 2014;768: 69-73.
- 318 17. Li SG, Ding YS, Niu Q, Xu SZ, Pang LJ, Ma RL, Jing MX, Feng GL, Liu JM, Guo SX. Grape seed  
319 proanthocyanidin extract alleviates arsenic-induced oxidative reproductive toxicity in male mice.  
320 *Biomed Environ Sci* 2015; 28: 272–280.
- 321 18. Schelz Z, Ocsovszki I, Bózsity N, Hohmann J, Zupko I. Antiproliferative Effects of Various  
322 Furanoacridones Isolated from *Ruta graveolens* on Human Breast Cancer Cell Lines. *Anticancer*  
323 *Res* 2016; 36: 2751–2758.
- 324 19. Sarshar S, Brandt S, Asadi Karam MR, Habibib M, Bouzari S, Lechtenberg M, Dobrindt U, Qin X,  
325 Goycoolea FM, Hensel A. Aqueous extract from *Orthosiphon stamineus* leaves prevents bladder  
326 and kidney infection in mice. *Phytomedicine* 2017; 28: 1–9
- 327 20. Zhou K, Raffoul JJ. Potential anticancer properties of grape antioxidants. *J Oncol* 2015; 80:3294.
- 328 21. Al-Rasheed NM, El-Masry TA, Tousson E, Hassan H, Al-Ghadeer A. Protective Potential of Grape  
329 Seed Proanthocyanidins Extract against Glivec (Imatinib Mesylate) Induced Liver Toxicity and  
330 Oxidative Stress in Male Rats. *Annual Research & Review in Biology* 2017; 20(6): 1-9.
- 331 22. Zhu FM, Du B, Li J. Effect of ultrafine grinding on physicochemical and antioxidant properties of  
332 dietary fiber from wine grape pomace. *Food Sci Technol Int* 2014; 20: 55–62.
- 333 23. El-Atrash A, Zaki S, Tousson E, Shoir M. Protective potential of grape seed extract against  
334 monosodium glutamate induced liver toxicity and oxidative stress in young rats. *Journal of*  
335 *Advanced Trends in Basic and Applied Science* 2017;1(3) :257-262.
- 336 24. Al-Naely A, Shattnan D. Effect of grape seed extract on biochemical factor and histological changes in  
337 liver and the kidney in albino rat infected hypo- hyperthyroidism induced laboratory by  
338 carbimazole and l-thyroxine. *Journal of Global Pharma Technology* 2017; 9(9):174-181
- 339 25. Thakur C, Saikia TC, Yadav RN. Total serum levels of triiodothyronine (T3) thyroxine (T4) and  
340 thyrotropine (TSH) in school going children of Dibrugarh district: an endemic goitre region of  
341 Assam. *Indian J Physiol Pharmacol* 1997; 41(2):167-70.
- 342 26. Maes M, Mommen K, Hendrickx D, Peeters D, D'Hondt P, Ranjan R, De Meyer F, Scharpe S.  
343 Components of biological variation, including seasonality, in blood concentrations of TSH, TT3,  
344 FT4, PRL, cortisol and testosterone in healthy volunteers. *Clin Endocrinol (Oxford)* 1997; 46 (5),  
345 587–598.
- 346 27. Mandel SJ, Brent GA, Larsen PR. Levothyroxine therapy in patients with thyroid disease. *Ann Intern*  
347 *Med* 1993; 119: 492–502.
- 348 28. Schumann G, Klauke R. New IFCC reference procedures for the determination of catalytic activity  
349 concentrations of five enzymes in serum: preliminary upper reference limits obtained in  
350 hospitalized subjects. *Clin Chim Acta* 2013; 327: 69-79.
- 351 29. Dumas BT, Watson WA, Biggs HG. Albumin standards and the measurement of serum albumin with  
352 bromocresol green. *Clin Chim Acta* 1977; 31: 87–96.
- 353 30. Moss DW, Henderson AR. *Clinical enzymology* C.A. Burtis, E.R. Ashwood (Eds.). Tietz Textbook of  
354 *Clinical Chemistry* (3<sup>rd</sup> Ed) 1999. W.B Saunders Company, Philadelphia, 1999; pp: 617- 721.
- 355 31. Mesbah L, Soraya B, Narimane S, Jean PF. Protective effect of flavonoides against the toxicity  
356 of vinblastine cyclophosphamide and paracetamol by inhibition of lipid- peroxidation and  
357 increase of liver glutathione. *Haema* 2004; 7: 59-67.

- 358 32. Saggi S, Sakeran M, Zidan N, Tousson E, Mohan A, Rehman H. Ameliorating effect of chicory  
359 (Chichorium intybus L.) fruit extract against 4-tert-octylphenol induced liver injury and oxidative  
360 stress in male rats. *Food and Chemical Toxicology*, 2014; 72: 138–146.
- 361 33. Misra HP, Fridovich I. The role of superoxide anion in the auto-oxidation of epinephrine and a simple assay  
362 for superoxidedismutase. *Journal of Biological Chemistry* 1972; 247: 3170 - 3175.
- 363 34. Beutler E, Duron O, Kelly BM. Improved method for the determination of blood glutathione. *J. Lab.*  
364 *Clin. Med.*, 1963; 61: 882–888.
- 365 35. Tousson E, Rafat BM, Hessien M, El Barbary AA, Sami A. P53 and Bcl2 apoptosis proteins in meso-  
366 2, 3-dimercaptosuccinic acid treated lead-intoxicated rabbits. *Toxicology and industrial health*.  
367 2011 Apr;27(3):271-8.
- 368 36. Tousson E. Histopathological alterations after a growth promoter boldenone injection in rabbits.  
369 *Toxicology and Industrial Health* 2016 Feb;32(2):299-305.
- 370 37. Shibutani, M, Woo GH, Fujimoto H, Saegusa Y, Takahashi M, Inoue K Hirose M, Nishikawa A.  
371 Assessment of developmental effects of hypothyroidism in rats from in utero and lactation  
372 exposure to anti-thyroid agents. *Reprod Toxicol* 2009; 28: 297–307
- 373 38. Giannini EG, Testa R and Savarino V. Liver enzyme alteration:A guide to clinicians. *MAJ* 2005;  
374 172(3): 367-79.
- 375 39. Hull K, Horenstein R, Naglieri R, Munir K, Ghany M, Celi FS. Two cases of thyroid storm associated  
376 cholestatic jaundice. *EndocrPract*.2007; 13: 476-480.
- 377 40. Biscoveanu M, Hasinski S. Abnormal results of liver function test in patients with Grave's disease.  
378 *Endocr Pract* 2000; 6: 367-369.
- 379 41. Wang R, Tan J, Zhang G, Zheng W, Li C. Risk factors of hepatic dysfunction in patients with Graves'  
380 hyperthyroidism and the efficacy of 131iodine treatment. *Medicine* 2017; 96: 5(6035).
- 381 42. Upadhyay G, Singh R, Kumar A, Kumar S, Kapoor A, God bole MM. Severe hyperthyroidism induces  
382 mitochondria mediated apoptosis in rat liver. *Hepatology* 2004; 39(4): 1120-1130.
- 383 43. Tousson E, Elgharabawy RM, Elmasry TA. Grape seed proanthocyanidin ameliorates cardiac toxicity  
384 induced by boldenone undecylenate through inhibition of nadph oxidase and reduction in the  
385 expression of NOX2 and NOX4. *Oxidative Medicine and Cellular Longevity* vol. 2018, ID  
386 9434385, 12 pages, <https://doi.org/10.1155/2018/9434385>, 2018.
- 387 44. Kandemir FM, Benzer F, Ozkaraca M, Ceribasi S, Yildirim NC, Ozdemir N. Protective antioxidant  
388 effects of grape seed extract in a cisplatin-induced hepatotoxicity model in rabbits. *Revue Méd*  
389 2012; 163(11): 539-545.
- 390 45. Sahoo D, Roy A, Bhanja S, Chainy G. Hypothyroidism impairs antioxidant defence system and  
391 testicular physiology during development and maturation. *General Comp Endocrinol* 2008; 156:  
392 63–70.
- 393 46. Das K, Chainy GN. Modulation of rat liver mitochondrial antioxidant defence system by thyroid  
394 hormone. *Biochim Biophys Acta* 2001; 1537: 1-13.
- 395 47. Kumar A, Sinha RA, Tiwari M , Singh R, Koji T, Manhas N, Rastogi L, Pal L, Shrivastava A, Sahu RP,  
396 Godbole MM (2007) Hyperthyroidism induces apoptosis in rat liver through activation of death  
397 receptor-mediated pathwaysq. *Journal of Hepatology* 46: 888–898.
- 398 48. Asayama K, Kato K. Oxidative muscular injury and its relevance to hyperthyroidism. *Free Radic Biol*  
399 *Med* 1990; 8: 293-303.
- 400 49. Bagchi D, Bagchi M, Stohs SJ, Das DK, Ray SD, Kuszynski CA, Joshi SS, Pruess HG: Free radicals  
401 and grape seed proanthocyanidin extract: importance in human health and disease prevention.  
402 *Toxicology* 2000;148: 187–197.

- 403 50. Davies KJ. Oxidative stress, antioxidant defenses, and damage removal, repair, and replacement  
404 systems. *IUBMB Life* 2000; 50: 279-289.
- 405 51. Chattopadhyay S, Sahoo DK, Subudhi U, Chainy GB. Differential expression profiles of antioxidant  
406 enzymes and glutathione redox status in hyperthyroid rats: a temporal analysis. *Comp. Biochem.*  
407 *Physiol C Pharmacol Toxicol Endocrinol* 2007; 146: 383-391.
- 408 52. Long M, Yang S, Han J, Li P, Zhang Y, Dong S. The Protective Effect of Grape-Seed  
409 Proanthocyanidin Extract on Oxidative Damage Induced by Zearalenone in Kunming Mice Liver.  
410 *Int J Mol Sci* 2016; 17(6): 808.
- 411 53. Diebold J, Baretton G, Felchner M, Meier W, Dopfer K, Schmidt M, Lohrs U. Bcl-2 expression, p53  
412 accumulation, and apoptosis in ovaria carcinomas. *Am J Clin Pathol* 1996; 105: 341–334.
- 413 54. Tousson E, Hafez E, Massoud A, Sweef O, Atta N. Protective role of folic acid in thyroxine-induced  
414 cardiac hypertrophy in hyperthyroid rat. *Biomedicine & Aging Pathology* 2013; 3: 89–95.  
415  
416

UNDER PEER REVIEW