

Determination of Acetylcholinesterase and Butylcholinesterase inhibitory activities of *G. asiatica* and *G. tiliaefolia* (family Tiliaceae) leaves..

ABSTRACT

Aims: Our study was carried out to appraise the acetylcholinesterase (AChE) and butyrylcholinesterase (BChE) inhibitory activities of *Grewia asiatica* and *Grewia tiliaefolia* leaves extracts.

Comment [A1]: aim

Comment [A2]: omit

Study Design: For the purpose of these experiments the extracts were subjected to an in-vivo study.

Comment [A3]: no mice groups are included in methodology

Place and Duration of Study: The study was carried out between June 2016 to December 2016 in the Department of Pharmacy, Southeast University, Dhaka, Bangladesh.

Comment [A4]: design should be explained

Comment [A5]: 3 years ago?

Methodology: In this study, acetylcholinesterase (AChE) and butyrylcholinesterase (BChE) effect of different fraction of crude ethanolic extract of both plants were examined using swiss albino mice at 300 mg/kg b.w. dose using slightly modified Elman coupled enzyme assay.

Comment [A6]: effects

Comment [A7]: fractions

Comment [A8]: rewrite as shown

Results: The highest inhibition of bovine brain acetylcholinesterase and human blood butyrylcholinesterase were exhibited by PEF and CLF of *G.asiatica* with the IC₅₀ values were found to be 55.88µg/ml and 26.14 µg/ml respectively whereas the highest inhibition of bovine brain acetylcholinesterase and human blood butyrylcholinesterasewere exhibited by CLF of *G.tiliaefolia*.

Comment [A9]: abb?

Comment [A10]: These enzymes

Comment [A11]: Sentence is incomplete

Conclusion: The result of the present study of various fractions of this plants have considerable amount of anti-acetylcholinesterase and anti-butyrylcholinesterase activity which suggest their effectiveness against various neurodegenerative disorders.

Comment [A12]: on

Comment [A13]: these

Comment [A14]: omit

Comment [A15]: activities

Keywords: Free radicals, Acetylcholinesterase, Butyrylcholinesterase, *G.asiatica*, *G.tiliaefolia*.

ABBREVIATIONS

IC₅₀: Concentration of an Inhibitor

Comment [A16]: not complete

PEFGA: Petroleum ether fraction of *G. asiatica*

CEEGA: Crude ethanolic extract of *G. asiatica*.

CLFGA=Chloroform fraction of *G.asiatica*

EAFGA=Ethyl acetate fraction of *G. asiatica*,

AEFGA=Aqueous Ethanolic fraction of *G. asiatica*.

AD: Alzheimer's Disease

AChE: Anti-acetylcholinesterase

Comment [A17]: not anti

BChE: Anti-butrylcholinesterase

Comment [A18]: C

Comment [A19]: not anti

1. INTRODUCTION

Plants serve various purposes and their usefulness to man is not limited to their role as sources of raw materials for industries; they are also consumed as food and sometimes used as medication. For ages, plants have provided man with diverse means of healing. In fact, many parts of plants such as fruits, seeds, barks, roots, and flowers have been used as medication to provide alternative therapies for various diseases that affect man and animals [1]. Medicinal plants contain potentially useful chemicals that are currently used for the manufacturing of modern therapeutic agents [2]. The evaluation of medicinal plants, used traditionally in treating Alzheimer's disease (AD) is of growing interest. Alzheimer's disease (AD) is a silent killer, one of the major leading causes of mortality after heart disease, cancer and stroke. AD associated with memory impairment that progressively declines in cognitive abilities and behaviors, which lead to the complete functional dependency that defines the dementia phase of the illness [3]. Therefore, inhibition of acetylcholinesterase (AChE), butyrylcholinesterase (BChE) enzyme and oxidation are the two promising strategies in the development of drug for neurological diseases like Alzheimer's and as well as in the treatment of Parkinson's disease, ataxia and dementia [4]. *Grewia asiatica* and *Grewia tiliaefolia* belonging to the family Tiliaceae [5] is a tree and has been used in traditional medicine to relief of various health problems such as cold, hepatitis, diarrhoea, heat stroke, dyspepsia, tuberculosis, sexual debility troubles, rheumatism and also important to promote intellect and enhancing memory, thus supporting its possible anti-Alzheimer's properties [6-7]. Several literature reviews demonstrated that the plant *G. asiatica* and *G. tiliaefolia* possess analgesic, anti-inflammatory, antioxidant, antimalarial, antidiabetic, antiemetic, antipyretic, antifungal, antiviral, antiplatelet, anticancer and immune-modulatory activities [8-16]. Thus, our main intention of the present study is to investigate the neuroprotective effect of different extracts of two medicinal plants *i.e.* *G. asiatica* and *G. tiliaefolia* (Family-Tiliaceae) available in Bangladesh on animal model.

Comment [A20]: AD is

Comment [A21]: enzymes

Comment [A22]: for

Comment [A23]: plants

2. MATERIALS AND METHODS

2.1 Collection of the plant materials and preparation of extracts:

For this present investigation whole plant of *Grewia asiatica* & *Grewia tiliaefolia* were collected from Moulovibazar, Bangladesh, in July 2016. After collection the plant was thoroughly washed with water and dried. The plants were identified by expert of Bangladesh National Herbarium, Mirpur, and DACB Accession number 73883 for *Grewia asiatica* & DACB Accession number 73882 for *Grewia tiliaefolia*. For whole plant part of the dried and powdered materials (100 g for) were soaked in 500 ml of 90% ethanol

Comment [A24]: omit

Comment [A25]: parts

Comment [A26]: what about other extracts (chloroform and ethyl acetate? where is the fractionation

for about 14 days at room temperature with occasional shaking. After 14 days the solution was filtered using cotton filter and Whitman's filter paper.



Fig 1: Accession number -DACB 73883



Fig 2: Accession number -DACB 73882 for *G. tiliaefolia*

for *G. asiatica*

2.1.1 Drugs and chemicals

5, 5'-dithio-bis-(2-nitro) benzoic acid (DTNB) (Sigma-Aldrich, Japan), Acetylcholine iodide (Sigma-Aldrich, Japan), Brain homogenate (Crude enzyme), Tris-Hcl buffer (Merck, Germany), Triton X-100 (Sigma chemical company, USA), BCA kit (bicinchoninic acid; Sigma Co., USA), Bovine serum albumin (Merck, India), Donepezil (Sigma-Aldrich, Japan),

Comment [A27]: ?

2.1.2 Acetyl cholinesterase inhibitory activity assay

The anti-acetylcholinesterase (AChE) assay was performed according to the colourimetric method of Ellman et al. [17], using acetylthiocholine iodide as a substrate. For the enzyme source, the rat brains were homogenised in a homogeniser with 5 volumes of a homogenisation buffer 910 mM Tris-HCl (pH

Comment [A28]: colorimetric

Comment [A29]: mice or rat

7.2), which contained 1M NaCl, 50mM MgCl₂ and 1% Triton X-1000 [18], centrifuged at 10,000 rpm for 15 min. The resulting supernatant was used as an enzyme source. All of the extraction steps were carried out at 4°C. Protein concentration was determined using the BCA kit (bicinchoninic acid) with bovine serum albumin (BSA) as a protein standard. The rates of hydrolysis by acetylcholinesterase were monitored spectrophotometrically. Each extract or standard (500µl) was mixed with an enzyme solution (500µl) and incubated at 37°C for 15 min. Absorbance at 405 nm was read immediately after adding an Ellman's reaction mixture [3.5 ml; 0.5 mM acetylthiocholine, 1 mM 5, 5'-dithio-bis (2-nitro benzoic acid)] in a 50 mM sodium phosphate buffer (pH 8.0) to the above reaction mixture. Reading was repeated for 10 min at 2 min intervals to verify that the reaction occurred linearly. The blank reaction was measured by substituting saline for the enzyme.

Acetylcholine → Thiocholine + Acetate; Thiocholine + dithiobisnitro → Benzoate yellow color

2.1.3 Anti-butrylcholinesterase activity assay

The anti-butrylcholinesterase (BchE) assay was performed according to the colourimetric method of Doctor et al. [19], using butrylthiocholine iodide as a substrate. For the enzyme source, the human blood was homogenised in a homogeniser with 5 volumes of a homogenisation buffer [10 mM Tris-HCl (pH 7.2), which contained 1M NaCl, 50mM MgCl₂ and 1% Triton X-100] [17], centrifuged at 10,000 rpm for 15 min. The resulting supernatant was used as an enzyme source. All of the extraction steps were carried out at 4°C. Protein concentration was determined using the BCA kit (bicinchoninic acid) with bovine serum albumin (BSA) as a protein standard. The rates of hydrolysis by butrylcholinesterase were monitored spectrophotometrically. Each extract or standard (500 µl) was mixed with an enzyme solution (500 µl) and incubated at 37°C for 15 min. Absorbance at 405 nm was read immediately after adding an Ellman's reaction mixture [3.5 ml; 0.5mM acetylthiocholine, 1mM 5, 5'-dithio-bis (2nitro benzoic acid)] in a 50 mM sodium phosphate buffer (pH 8.0) to the above reaction mixture. Reading was repeated for 10 min at 2 min intervals to verify that the reaction occurred linearly. The blank reaction was measured by substituting saline for the enzyme.

Comment [A30]:

Comment [A31]: italic

Comment [A32]: ????

Butrylcholine → Thiocholine + Acetate; Thiocholine + dithiobisnitro → Benzoate yellow Color

3. RESULTS

3.1 Acetyl cholinesterase inhibitory activity assay of *G. asiatica*

The acetylcholinesterase inhibitory activity of different extractives was determined by Ellman's method. This method estimates acetylcholinesterase (AChE) using acetylcholine iodide (substrate) and dithiobisnitro benzoic acid (DTNB). The enzymatic activity was measured by the yellow color compound produced by thiocholine when it reacts with dithiobisnitro benzoate ion. The results for anticholinesterase activity of crude ethanolic extracts different fraction of *G. asiatica* are given in Table 1.

Table 1: Acetylcholinesterase inhibitory activity of the crude ethanol extract of *G. asiatica* and Donepezil (standard) at different concentration.

Comment [A33]: the table is not only the crude ethanolic extract

Comment [A34]: concentrations

Name of Sample	Conc. (µg/ml)	% of Scavenging			% of Scavenging Mean ± STD	IC ₅₀ (µg/ml)±STD
		a	b	C		
CEEGA	31.25	22.22	22.79	21.65	22.22±0.465	286.78±1.77
	62.5	24.50	25.64	25.07	25.07±0.465	
	125	30.77	30.20	84.64	30.77±0.465	
	250	43.87	43.30	43.87	43.48±0.269	
PEFGA	31.25	57.55	58.12	56.98	57.55±0.465	55.88±0.889
	62.5	56.41	54.70	56.70	55.94±0.881	
	125	69.52	68.95	70.09	69.52±0.465	
	250	71.22	70.66	70.94	70.94±0.232	
CLFGA	31.25	55.27	55.84	54.70	55.27±0.465	59.62±0.529
	62.5	51.85	52.42	52.99	52.42±0.465	
	125	62.39	62.68	61.54	62.20±0.484	
	250	63.82	64.39	63.53	63.91±0.355	
EAFGA	31.25	27.92	28.49	28.77	23.08±0.355	250.29±3.58
	62.5	34.19	34.76	33.90	34.28±0.355	
	125	43.30	42.17	43.87	43.11±0.711	
	250	49.00	50.71	50.14	49.95±0.711	
AEFGA	31.25	23.36	22.79	23.07	23.08±0.232	268.66±0.3.37
	62.5	37.04	37.61	36.47	37.07±0.465	
	125	40.46	42.17	41.60	41.41±0.711	
	250	45.87	46.44	47.29	46.53±0.585	
Donepezil	31.25	54.13	54.70	55.84	54.89±0.711	28.47±0.150
	62.5	71.23	71.23	71.79	71.41±0.269	
	125	73.50	73.79	74.07	73.79±0.232	
	250	75.21	75.78	76.07	75.68±0.355	

Comment [A36]: Scavenging what?

Comment [A35]: concentrations tested in vivo or in vitro?

Further, the anticholinesterase activity of all the fractions of crude ethanol extract such as CEEGA, PEFGA, CLFGA, EAFGA and AEFGA have been investigated at 250 µg/ml concentration. Among the fractions the highest activity was found in PEFGA (55.88 % inhibition).

Comment [A37]: Not highest inhibition

Fig 3: Acetylcholinesterase inhibitory activity of different fractions of crude ethanol extract different fraction of *G. asiatica* and donepezil at different concentration.

Comment [A38]: concentrations





Fig 4: IC₅₀ (µg/ml) values of different extractives of *G. asiatica* for Anti-acetylcholinesterase activity Assay.



3.2 Acetyl cholinesterase inhibitory activity assay of *G. tiliaefolia*

The acetylcholinesterase inhibitory activity of different extractives was determined by Ellman's method. This method estimates acetylcholinesterase (AChE) using acetylcholine iodide (substrate) and dithiobisnitro benzoic acid (DTNB). The enzymatic activity was measured by the yellow color compound produced by thiocholine when it reacts with dithiobisnitro benzoate ion. The results for anticholinesterase activity of crude ethanolic extracts different fractions of *G. tiliaefolia* are given in Table 2.

Table 2: Acetylcholinesterase inhibitory activity of the crude ethanol extract of *G.tiliaefolia* and Donepezil (standard) at different concentration.

Name of Sample	Conc. (µg/ml)	% of Scavenging			% of Scavenging Mean ± STD	IC ₅₀ (µg/ml)±STD
		a	b	C		
CEEGT	31.25	31.91	32.48	33.05	32.48±0.465	272.53±2.73
	62.5	38.18	38.46	37.61	38.08±0.355	
	125	39.32	38.75	39.89	39.31±0.465	
	250	45.58	46.15	46.72	46.15±0.465	
PEFGT	31.25	56.98	57.83	58.12	57.64±0.484	59.61±0.529
	62.5	52.42	51.85	52.99	52.42±0.465	
	125	64.10	63.25	63.82	63.72±0.355	
	250	68.95	68.38	69.23	68.85±0.355	
CLFGT	31.25	46.15	46.72	48.43	47.10±0.968	57.13±0.485
	62.5	54.13	55.27	54.70	54.70±0.465	
	125	56.69	57.83	58.40	57.64±0.711	
	250	60.11	58.12	59.54	59.26±0.839	
EAFGT	31.25	29.06	29.63	28.77	29.15±0.355	277.20±5.01
	62.5	36.75	37.03	34.19	35.99±1.28	
	125	40.17	40.46	42.16	40.93±0.880	
	250	44.16	46.15	45.01	45.10±0.816	
AEFGT	31.25	26.78	26.21	26.78	26.59±0.268	268.66±3.37
	62.5	31.91	32.19	32.48	32.19±0.232	
	125	39.60	39.03	40.46	39.69±0.585	
	250	45.87	46.44	47.29	46.53±0.585	
Donepezil	31.25	54.13	54.70	55.84	54.89±0.711	28.47±0.150
	62.5	71.23	71.23	71.79	71.41±0.269	
	125	73.50	73.79	74.07	73.79±0.232	
	250	75.21	75.78	76.07	75.68±0.355	

Further, the anticholinesterase activity of all the fractions of crude ethanol extract such as CEEGT, PEFGT, CLFGT, EAFGT and AEFGT have been investigated at 250 µg/ml concentration. Among the fractions the highest activity was found in CLFGT (57.17 % inhibition).

Comment [A39]: not found

Fig. 5: Acetylcholinesterase inhibitory activity of different fractions of crude methanol extract different fraction of *G. tiliaefolia* and donepezil at different concentration.

Comment [A40]:





Fig.6: Acetylcholinesterase inhibitory activity of different fractions of crude methanol extract different fraction of *G. tiliaefolia* and *donepezil* at different concentration.



3.3 Butyrylcholinesterase inhibitory (BchE) activity assay of *G. asiatica*

The Butyrylcholinesterase inhibitory activity of extractives was determined by Ellman's method. This method estimates Butyrylcholinesterase (BchE) activity using butyrylthiocholine iodide (substrate) and dithiobisnitro benzoic acid (DTNB). The enzymatic activity was measured by the yellow color compound produced by thiocholine when it reacts with dithiobisnitro benzoate ion. The results for anticholinesterase activity of crude methanol extracts different fraction of *G. asiatica* are given in Table 3.

Table 3: Anti-butylcholinesterase activity assay of the crude ethanol extracts different fraction of *G. asiatica* and Donepezil (standard) at different concentrations.

Name of Sample	Conc. (µg/ml)	% of Scavenging			% of Scavenging Mean ± STD	IC ₅₀ (µg/ml) ±STD
		a	b	C		
CEEGA	31.25	50.33	51.00	52.67	51.33±0.981	30.84±0.577
	62.5	53.00	53.67	54.33	53.67±0.544	
	125	55.67	57.00	57.67	56.77±0.831	
	250	58.33	59.67	60.33	59.44±0.831	
PEFGA	31.25	52.33	53.33	53.00	52.89±0.416	29.54±0.233
	62.5	55.33	56.33	57.00	56.22±0.685	
	125	60.33	61.00	61.67	61.00±0.544	
	250	64.33	63.67	64.00	64.00±0.272	
CLFGA	31.25	59.67	60.00	59.67	59.78±0.157	26.14±0.069
	62.5	61.33	62.33	61.67	61.78±0.416	
	125	66.67	69.00	69.67	68.44±1.29	
	250	75.67	75.67	76.33	75.89±0.314	
EAFGA	31.25	56.33	57.00	57.33	56.89±0.416	27.47±0.201
	62.5	58.33	59.00	60.00	59.11±0.685	
	125	61.33	62.33	63.00	62.22±0.685	
	250	65.00	64.67	65.67	65.11±0.415	
AEFGA	31.25	34.33	33.67	34.33	34.11±0.314	129.33±1.47
	62.5	42.67	43.00	43.67	43.11±0.415	
	125	47.67	48.33	49.00	48.33±0.544	
	250	56.33	56.67	57.00	56.67±0.272	
Donepezil	31.25	55.00	55.67	56.00	55.56±0.416	28.13±0.170
	62.5	60.00	60.33	61.00	60.44±0.416	
	125	70.33	71.00	71.67	71.00±0.544	
	250	84.33	85.33	85.00	84.89±0.416	

Among the fractions of crude ethanol extract, CLFGA and EAFGA showed the most potent activity with IC₅₀ value of 26.14µg/ml and 27.47 µg/ml which is higher than that of DON (Standard) With IC₅₀ value of 28.13 µg/ml. On the other hand, (PEFGA), (CEEGA) and (AEFGA) [fraction] showed free radical scavenging activity with IC₅₀ value of 29.54 µg/ml, 30.84µg/ml and 129.33 µg/ml respectively. Our results clearly demonstrate that the extractives of *G.asiatica* possess antiradical activity.

Comment [A41]: fractions

Comment [A42]: clarify the scavenging % and difference between IC50

Fig.7: Anti-Butyrylcholinesterase activity of different fractions of crude ethanol extract of *G. asiatica* and donepezil at different concentrations.





Fig.8: IC₅₀ (µg/ml) values of different extractives of *G. asiatica* for Anti-Butyrylcholinesterase activity Assay.



3.4 Butyrylcholinesterase inhibitory (BchE) activity assay of *G. tiliacifolia*

The Butyrylcholinesterase inhibitory activity of extractives was determined by Ellman's method. This method estimates Butyrylcholinesterase (BchE) activity using butyrylthiocholine iodide (substrate) and dithiobisnitro benzoic acid (DTNB). The enzymatic activity was measured by the yellow color compound produced by thiocholine when it reacts with dithiobisnitro benzoate ion. The results for anticholinesterase activity of crude methanol extracts different fraction of *G. tiliacifolia* are given in Table 4.

Table 4: Anti-butylcholinesterase activity assay of the crude ethanol extracts different fraction of *G. tiliaefolia* and Donepezil (standard) at different concentrations.

Name of Sample	Conc. (µg/ml)	% of Scavenging			% of Scavenging Mean ± STD	IC ₅₀ (µg/ml)±STD
		a	b	C		
CEEGT	31.25	47.67	49.00	49.67	48.78±0.831	32.33±0.550
	62.5	52.33	53.00	53.67	53.00±0.544	
	125	57.33	57.67	58.67	57.89±0.567	
	250	61.00	61.67	59.67	60.78±0.831	
PEFGT	31.25	54.67	56.00	56.33	55.67±0.720	28.07±0.366
	62.5	60.33	61.67	62.00	61.33±0.720	
	125	62.33	63.00	63.33	62.89±0.415	
	250	67.00	67.67	68.33	67.67±0.544	
CLFGT	31.25	57.67	59.00	59.67	58.78±0.831	26.59±0.378
	62.5	63.33	64.33	64.67	64.11±0.567	
	125	66.33	67.67	68.33	67.44±0.831	
	250	73.00	71.67	73.33	72.67±0.720	
EAFGT	31.25	53.00	54.00	54.33	53.78±0.567	29.06±0.308
	62.5	61.67	61.00	60.00	60.89±0.685	
	125	63.67	63.33	64.00	63.67±0.272	
	250	68.00	67.00	65.67	66.89±0.956	
AEFGT	31.25	32.00	33.00	31.67	32.22±0.567	134.25±0.92
	62.5	40.33	39.67	41.00	40.33±0.544	
	125	46.33	47.00	46.33	46.56±0.314	
	250	58.67	57.67	57.00	57.78±0.685	
Donepezil	31.25	55.00	55.67	56.00	55.56±0.416	28.13±0.170
	62.5	60.00	60.33	61.00	60.44±0.416	
	125	70.33	71.00	71.67	71.00±0.544	
	250	84.33	85.33	85.00	84.89±0.416	

Fig.9: Anti-Butyrylcholinesterase activity of different fractions of crude ethanol extract of *G.tiliaefolia* and donepezil at different concentrations.





Fig.10: IC₅₀ (µg/ml) values of different extracts of *G.tiliaefolia* for Anti-Butyrylcholinesterase activity Assay.



4. DISCUSSIONS

Alzheimer's disease (AD) is a progressive neurodegenerative disorder characterised by loss of memory and cognition. The AD has revealed the development of new therapeutic drugs for the AD is underway. Currently, only five medications are approved by the Food and Drug Administration to treat AD. Four of them are acetylcholinesterase inhibitors such as donepezil, galantamine, rivastigmine, tacrine and the fifth is the N-methyl-d-aspartate antagonist memantine [19]. The history of drug discovery showed that plants are highly rich sources of bioactive compounds and lead to the development of drugs for the treatment of neurological diseases including Alzheimer's disease [20]. In traditional practices of medicine

Comment [A43]: omit

plants have been used to enhance cognitive function and to alleviate other symptoms associated with the AD. Inhibition of acetylcholinesterase enhances cholinergic transmission by reducing enzymatic degradation of acetylcholine is a promising strategy for the development of AD-drug. Acetylcholinesterase inhibitors are the only source of the compound that is currently approved for the treatment of Alzheimer's disease.

However, our results revealed moderate anti-acetylcholinesterase effect of *G.asiatica* and *G. tiliaefolia*. Also, the activity, the different fractions of the crude extract such as petroleum ether, crude ethanol extract, chloroform, ethyl acetate and aqueous fractions were examined similarly at a concentration of 250 µg/ml. Among the fractions, the CLFGA and CLFGT had the highest % of inhibition on Butyrylcholinesterase at a concentration of 250 µg/ml. However, our results revealed significant anti-butyrylcholinesterase effect of *G. asiatica* and *G. tiliaefolia*.

5 CONCLUSIONS

The present study was undertaken to investigate the *in vivo* anticholinesterase and anti-butyrylcholinesterase effects of *G. asiatica* and *G. tiliaefolia*. Inhibition of acetylcholinesterase, which enhances cholinergic transmission by reducing the enzymatic degradation of acetylcholine, is a promising strategy for the development of neurodegenerative disorders like Alzheimer's diseases drug. *In vivo* effectiveness of *Grewia nervosa* and its components remains to be investigated. The results indicate that *G. asiatica* and *G. tiliaefolia* may be of value for an effective treatment for Alzheimer's disease.

Comment [A44]: omit

CONSENT

It is not applicable

ETHICAL APPROVAL

The protocol of the experiment was approved by the animal ethics committee of the Department of Pharmacy, Southeast University, Dhaka, Bangladesh. The animals care and health were maintained according to the guidelines of the National Institutes of Health.

REFERENCES

1. Phyllistin AB, James FB. Tips for preventing food poisoning "Herbs" American No. 1 Guide Natural health. Publ. Averige. 2000; 9:383-6.
2. Okigbo RN, Anuagasi CL, Amadi JE. Advances in selected medicinal and aromatic plants indigenous to Africa. Journal of Medicinal Plants Research. 2009 Feb 28;3(2):086-95.

3. McKhann G, Drachman D, Folstein M, Katzman R, Price D, Stadlan EM. Clinical diagnosis of Alzheimer's disease: Report of the NINCDS-ADRDA Work Group* under the auspices of Department of Health and Human Services Task Force on Alzheimer's Disease. *Neurology*. 1984 Jul 1;34(7):939-.
4. Ahmad W, Ahmad B, Ahmad M, Iqbal Z, Nisar M, Ahmad M. In vitro inhibition of acetylcholinesterase, butyrylcholinesterase and lipoxygenase by crude extract of *Myricaria elegans* Royle. *J Biol Sci*. 2003; 11:1046-9.
5. Heywood VH, Brummitt RK, Culham A, Seberg O. Flowering plant families of the world. Ontario: Firefly Books; 2007.
6. Zia-Ul-Haq M, Stanković MS, Rizwan K, Feo VD. *Grewia asiatica* L., a food plant with multiple uses. *Molecules*. 2013 Mar;18(3):2663-82.
7. Singariya P, Mourya KK, Kumar P. In vitro studies of antimicrobial activity of crude extracts of the Indian grasses *Dhaman* (*Cenchrus ciliaris*) and *Kala-Dhaman* (*Cenchrus setigerus*). *Indian journal of pharmaceutical sciences*. 2012 May;74(3):261.
8. Gupta MK, Sharma PK, Ansari SH, Lagarkha RE. Pharmacognostical evaluation of *Grewia asiatica* fruits. *Int J Plant Sci*. 2006;1(2):249-51.
9. Gupta MK, Lagarkha R, Sharma DK, Sharma PK, Singh R, Ansari HS. Antioxidant activity of the successive extracts of *Grewia asiatica* leaves. *Asian Journal of Chemistry*. 2007 Jul 1;19(5):3417.
10. Sharma KV, Sisodia R. Hepatoprotective efficacy of *Grewia asiatica* fruit against oxidative stress in swiss albino mice. *International Journal of Radiation Research*. 2010 Sep 1;8(2):175.
11. Paviaya US, Kumar P, Wanjari MM, Thenmozhi S, Balakrishnan BR. Analgesic and anti-inflammatory activity of root bark of *Grewia asiatica* Linn. in rodents. *Ancient science of life*. 2013 Jan;32(3):150.
12. Khattab HA, El-Shitany NA, Abdallah IZ, Yousef FM, Alkreathy HM. Antihyperglycemic potential of *Grewia asiatica* fruit extract against streptozotocin-induced hyperglycemia in rats: anti-inflammatory and antioxidant mechanisms. *Oxidative medicine and cellular longevity*. 2015;2015.
13. Khan RS, Asghar W, Khalid N, Nazir W, Farooq M, Ahmed I, Syed QA. Phalsa (*Grewia asiatica* L) fruit berry a promising functional food ingredient: A comprehensive review. *Journal of Berry Research*. 2019(Preprint):1-5.
14. Ahamed MB, Krishna V, Dandin CJ. In vitro antioxidant and in vivo prophylactic effects of two γ -lactones isolated from *Grewia tiliaefolia* against hepatotoxicity in carbon tetrachloride intoxicated rats. *European journal of pharmacology*. 2010 Apr 10;631(1-3):42-52.
15. Thamizh SN, Vengatakrishnan V, Murugesan S, Damodar KS. Antioxidant and antiproliferative activity of methanolic extract of *Grewia tiliaefolia* (Vahl) bark in different cancer cell lines. *International Journal of Pharmacy and Life Sciences (IJPLS)*. 2010;1(2):54-60.
16. Sakat SS, Juvekar AR. Analgesic and antipyretic activity of aqueous extract of *Grewia tiliaefolia* Vahl leaves. *Journal of Pharmacy Research*. 2009;2(9):1475-8.
17. Ellman GL, Courtney KD, Andres Jr V, Featherstone RM. A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochemical pharmacology*. 1961;7(2):88-95.

18. Rieger F, Shelanski ML, Greene LA. The effects of nerve growth factor on acetylcholinesterase and its multiple forms in cultures of rat PC12 pheochromocytoma cells: increased total specific activity and appearance of the 16 S molecular form. *Developmental Biology*. 1980;76(1):23843.
19. McShane R, Areosa AS, Minakaran N. Memantine for dementia. 2006; CD003154-CD003154.
20. Tomlins SA, Mehra R, Rhodes DR, Cao X, Wang L, Dhanasekaran SM, KalyanaSundaram S, Wei JT, Rubin MA, Pienta KJ, Shah RB. Integrative molecular concept modeling of prostate cancer progression. *Nature Genetics*. 2007; 39(1):41.

UNDER PEER REVIEW