

Original Research Article

Pedogenic and contemporal issues of fluvisol characterization and utilisation in the Wetlands of Bamenda Municipality, Cameroon

ABSTRACT

Fluvisols in urban wetlands in Bamenda Municipality play a vital role in vegetable production but under immense pressure. Seven representative soil profiles and 21 surface soil samples were morphologically and physico-chemically characterized to classify the soils, evaluate their agro-utilization constraints, and to provide adequate data for planning sustainable land management. The soil samples were analyzed using standard procedures. Critical levels established for tropical crops and vegetables was used to declare deficiency of soil nutrients. The coefficient of variation was used as an index of soil variability, while sources of soil variation and subsequent grouping into management units were identified using principal component analysis. The soils classified as Humi-umbric fluvisols are developed from young alluvio-colluvial material of granitic origin. Like other physico-chemical properties, organic matter varied irregularly down the profile. With the exception of pH which was slightly ($CV < 15\%$) variable, most soil properties were moderately ($CV = 15-35\%$) to highly ($CV > 35\%$) variable. Some correlation coefficients between the soil parameters were highly significant ($p < 0.01$) ranging - 0.95 to 0.99, but most of them have correlation values less than 0.5. Six principal components (PCs) grouping soils in management units explained 96.2% of the variations observed in the soil properties. The PCs were: base status, organic matter, weathering and moisture retention, acidity, dispersal and N-mineralization, and mineral neo-synthesis factors. We recommend that a detailed mapping of soil properties be carried out for the establishment of a soil fertility map; and individual soil management practices defined for identified units instead of a common management for all units in the municipality.

Key words: Fluvisols, soil classification, soil characteristics, soil utilization, soil management

Introduction

The World Reference Base for Soil Resources (WRB) has 32 soil groups (FAO, 2006) some of which occur in West and Central Africa. Deckers (1993) and Bationo *et al.* (2006) outlined the typical distribution in terms of agroecology. Fluvisols, an important soil group, and found in wetlands covers an estimated area of over 350 million hectares worldwide (FAO, 2006). In Cameroon, they occupy 2.6% of the total land area (Ngachie, 1992). Found on all continents under all kinds of climatic conditions they occur mainly on flood plains, fans and deltas of rivers (Driessen and Dudal, 1991). In the upper part of the drainage basin, they are normally confined to narrow strips along the river. The fluvisols of floodplain wetlands are characterized by seasonal hydrological dynamics, which strongly influence vegetation stand (Haase and Neumeister, 2001). The later equally noted that in the last century, hydrological

system of seasonal flooding of flood plains has been influenced intensively by man in the form of riversides regulation, and utilisation of parts of the floodplain wetlands for agricultural purposes thus drying up wetland areas and soils. Most importantly, these soils perform an ecological function of accumulating and fixing airborne and fluvially transported nutrients and contaminants. Since soils are good contamination indicators of floodplain ecosystems, they have been the object of research on the soil characteristics and dynamics. The mobilization of organic and inorganic contaminants, results to considerable damages of sensitive soil edaphic factors and the vegetation as well as migration of contaminants into the water saturated zone (Haase and Neumeister, 2001). Increases in the soil acidity will fasten these migration processes.

Comment [RA1]: insert reference

In fluvisols, soil forming processes, other than the formation of a surface horizon through accumulation of organic matter, have not left their marks (Yerima and Van Ranst, 2005a; FAO, 2006). These juvenile soils show few or no evidence of weathering and soil formation below 25 cm depth except possible gleying. Permanent or seasonal saturation with water, because of permanent or recurrent anaerobic conditions and low biological activity, tends to preserve the original stratified nature of the original deposits. But in the course of time, when the effects of pedogenetic agents such as soil animals, roots, repeated wetting and drying proceed downward in the profile, a cambic horizon may be formed. This implies the transformation of the Fluvisol into a Cambisol or Gleysol, depending on the drainage conditions. Flooding of some parts of the wetlands with fresh water, will change the geochemistry of the soils as the basis for flora and fauna.

Comment [RA2]: insert reference

Comment [RA3]: reference

Today, the nutrients contents of the flooded and non-flooded areas differ considerably. Fluvisols harbor a wide range of crops or are under grassland. They are usually fertile but may need flood control. Many Fluvisols in the humid tropics, Cameroon inclusive, are under vegetable production especially during a dry period where microbial activity is stimulated which promotes mineralization of organic matter and hence release of plant nutrients. Other factors that influence the potential productivity of wetland soils are their topography, minimal soil erosion, natural fertility, and location in climatic regions of adequate rainfall for most crops.

Comment [RA4]: reference

Fertility levels have a direct bearing on the potential for development of wetland soils, especially in urban areas. A vast majority of wetland benefits accrue to the local population which makes it important to conserve these valuable natural resources. Today, their exploitation particularly for economic purpose is alarming (Kometa 2013, Asongwe *et al.*, 2014). The urban wetlands of Bamenda Municipality, which is part of the Western Highlands of Cameroon and the focus of this study, are coming under increased pressure for the production of food to meet the needs of the increasing population. The municipality and the wetlands are characterized with a heavy demographic, technogenic and municipal load. The problem with the wastewaters and solid wastes acquires increasing importance. Here, these wetlands ecological niches have been drained for vegetable cultivation. These valuable areas are not only exploited for market gardening involved in the cultivation of vegetables, carrots, tomatoes, green beans, Irish potatoes but also for the planting of eucalyptus which help to drain the water in these wetlands (Kometa, 2013; Asongwe *et al.*, 2014). The result is that at the present rate of exploitation the survival of this resource is at stake. Enhanced understanding of the morphological and physico-chemical properties of the fluvisols would avail pertinent information for assessing the potentials and constraints of the soils for different uses and management options. It is on this backdrop that the objectives of this study were to a) characterize the soils based on the morphological field description and physico-chemical properties; and to classify the soils using the World Reference Base (WRB) for Soil

Comment [RA5]: *et al.*, or *et al.*, uniform please

Resources, b) Identify the current utilization constraints and, c) to provide data for use by stakeholders in planning sustainable land management in the Bamenda urban Municipality.

Comment [RA6]: none of these objectives or a joint proposal appears in the summary

Materials and Methods

Description of the study area

The area covered by this study includes urban and peri-urban wetlands in the Bamenda City Council of the North West Region of Cameroon (Figure 1). It is part of the Bamenda escarpment and located between latitudes 5° 55' N and 6° 30' N and longitudes 10° 25' E and 10° 67' E. The town shows an altitudinal range of 1200 - 1700 m, and is divided into two parts by escarpments; a low lying gently undulating part with altitudes ranging from 1200 to 1400 m, with many flat areas that are usually inundated for most parts of the year, and an elevated part at 1400 to 1700 m altitude that forms the crest from which creeks and streams supplying the low lying parts take their rise. This area has two seasons; a long rainy season, which runs from mid-March to mid- October and a short dry season that spans from mid-October to mid-March. The area lies within the thermic and hyperthermic temperature regimes. Mean annual temperatures stand at 19.9 °C. January and February are the hottest months with mean monthly temperatures of 29.1 °C and 29.7 °C, respectively. Yerima and Van Ranst (2005a) reported that the area is dominated by the Ustic and Udic moisture regimes with the Udic extending to the south. Annual rainfall ranges from 1300-3000 mm (Ndenecho, 2005). The area has a rich hydrographical network with intense human activities and a dense population along different water courses in the watershed. The main human activity in and around this area is agriculture, which according to GP-DERUDEP (2006) involves over 70% of the population and make use of rudimentary tools. The area equally harbors the commercial center that has factories ranging from soap production, and garages to metallurgy, which may be potential sources of pollutants. An important vegetation type in this area is the raffia palm bush, which is largely limited to the wetlands (Valleys and depressions). The raffia palm (*Raffia farinifera*) provides raffia wine, a vital economic resource to the indigenes who are fighting against the cultivation of these wetlands by vegetable farmers.

Comment [RA7]: masl (meters above sea level), or refers to a slope

Comment [RA8]: they only differ by 60 days, please correct the wording

Comment [RA9]: After 14 years, this proportion is still maintained, it is necessary to update the reference and the value condition, if it changed. The migration factor has not modified this figure?

Sampling strategy

In the study (Figure 1), seven representative soil profiles were described. These profiles represented different variability in landscapes and were pedologically and morphologically described in the field as outlined by the Soil Survey Staff (2003). The depth of soil profiles and thickness of horizons were determined with the help of a measuring tape. Soil colours were determined with the help of a Munsell color chart (USDA, 1975). The soils of the study area are composed of poorly drained fluvisols derived from aluvi-colluvial parent materials, on a less than 1% slope. Samples were collected by genetic horizons from pits excavated to maximum 1.5 m deep depending on the water table. Twenty bulk samples were collected from the various horizons. Similarly, 21 top soil samples (0 - 25 cm) were randomly collected with the use of a hand trowel into black plastic bags in the wetlands. The soil samples were air-dried and ground using a ceramic mortar. The fine earth fractions were screened through a 2-mm sieve. They were analysed for routine parameters in the Environmental and Analytical Chemistry Laboratory of the University of Dschang Cameroon. Particle size distribution, cation exchange capacity (CEC), exchangeable bases, EC, and pH were determined on the fine earth fraction by standard procedures (Pauwells *et al*, 1992). pH was measured both in water and KCl (1:2.5 soil/water mixture) using a glass electrode pH meter. Part of the fine soil was ball-milled for organic carbon (OC) and Kjeldahl-N analysis (Pauwells *et al.*, 1992). Available P was determined by method of Bray II, exchangeable cations were determined by extracting with 1N ammonium acetate at pH 7, potassium and Na in the extract were

Comment [RA10]: Greater precision is needed in the description of sampling of wetlands, which methodology was mainly followed for this. How representative was the sample based on the area collected? How many wetlands did they locate to be representative of the locality?

Comment [RA11]: Based on which standard they applied to the handling of the samples, how was the transfer to the laboratory? Was it direct from the field to the laboratory or was it previously refrigerated? No questions to be answered since they are elements that influence the possible variation of the chemical composition of the samples.

determined using flame photometer and Mg and Ca determined by complex metric titration. Exchangeable acidity was extracted with 1M KCl followed by quantification of Al and H by titration (Pauwels *et al.*, 1992). Effective cation exchange capacity (ECEC) was determined as sum of bases and exchanged acidity. Apparent CEC (CEC7) was determined directly as outlined by Pauwels *et al.* (1992). Based on critical values of nutrients established for vegetables, nutrients were declared sufficient or deficient.




Figure 1: Map of Bamenda Municipality showing fluvisols sampling areas.

Analytical Approach

Data obtained were subjected to statistical analyses (univariate statistics: mean, range, maximum and minimums and multivariate statistics, correlation and principal component analyses: factorial analyses) using Microsoft Excel 2007 and SPSS statistical package 20.0. Soil properties were assessed from their variability using coefficient of variation (CV %) and compared with variability classes (Table 1 and 2, respectively).

In order to identify the factors causing variation in the soil properties, principal component analysis (PCA) were applied. Varimax rotation was applied to do away with the problem of autocorrelation and to reduce the contributing soil factors to orthogonal principal components (Phil-Eze, 2010). The coefficients of the principal components are the eigen values. Analysis of correlation and coefficient of variations also assisted to identify soil factors that correlate significantly and differ, respectively.

$$\%CV = \frac{Sd \times 100}{X}$$

Where:

Comment [RA12]: It is important that they declare the experimental design they followed for the described procedure, or if it was a non-experimental design. As well as omitting the declaration of the criteria of normality and homoscedasticity for the development of the correlation, it is essential to incorporate these aspects.

Sd = standard deviation

X = arithmetic mean of soil properties

Table 1: Grouping coefficient of variation into variability classes.

CV %	Variability grouping (class)
< 15 %	Slightly variable
15-35 %	Moderately variable
➤ 35 %	Highly variable

Table 2: Critical Values of nutrients and soil properties

Properties	Critical values				
	Very low	Low	Medium	High	Very high
OM %	<1	1-2	2-4.2	4.2-6	>6
Total N g/Kg	< 0.5	0.5-1.25	1.25-2.25	2.25-3.0	>3.0
C/N	<10 = good, 10-14 = medium and >14 = poor				
Ca cmol/kg	<2	2-5	5-10	10-20	>20
Mg cmol/kg	<0.5	0.5-1.5	1.5-3	3-8	>8
K cmol/kg	<0.1	0.1-0.3	0.3-0.6	0.6-1.2	>1.2
Na cmol/kg	<0.1	0.1-0.3	0.3-0.7	0.7-2.0	>2
Bray 2-P mg/kg	<7	< 7-16	16-46	>46	
pH	5.3= 6.0 moderately acid, 6.0-7.0 = slightly acid, 7.0-8.5 moderately alkaline				
ESP %		<2	2-8	8-15	15-27
CEC ₇ cmol/kg	0-20	21-40	41-60	61-80	81-100

Adapted from Beernaert and Bitondo, (1992)

In order to identify the factors causing variation in the soil properties, principal component analyses (PCA) was applied.

RESULTS AND DISCUSSIONS

Soil profile Characterization and Classification

Pedological and Morphological characterization of the soils showed that, they were fluvisols with parent materials of alluvi-colluvial deposits, of granitic origins. The soils also indicated varying degree of oxidation and reduction as indicated by red and blue colours of (ferric and ferrous of Fe) mottles. Munsell color values of the various horizons of the soils did not show a definite pattern throughout the profiles and generally ranged around 2.5 to 10. These marked differences observed could be ascribed to variation in drainage, topographic and/or anthropogenic influences. Similar variations in hydric soils have been noticed in many parts of the world (Van Diepen, 1986, Eswaran and Cook, 1986, and Egbuchua and Ojeifo, 2007). Numerous mottles predominantly of reddish coloration of varying percentages were noticed within the various horizons in the different profiles. According to Wilding and Rehage (1986), major pedogenetic processes in hydric soils apart from the mobilization and immobilization of iron and manganese concretions, includes mottling (FAO/UNESCO, 1999). The texture of soils under investigation were dominated by sandy loams, with a few sandy clay loams, based on particle size distribution. At the surface horizons, the structure of the soils were dominantly granular parting to weak sub-angular blocky shapes. Lower horizons showed varied structures with numerous relics of man-used materials (cloths, plastics, pieces of metals, charcoal, etc.) within most of the profiles (Figure 2). Consistence,

when moist, generally varied from friable through slightly sticky to sticky and was plastic when wet. Few to many fine roots dominated the surface horizons. Clear smooth and gradual boundaries were some prominent morphological properties. Soil moisture regime was ustic, while the temperature was isothermic. The water table was very close to the surface with a maximum depth of 120 cm obtained at the profile Fuwambi – Ntahsen. Profiles at lower river courses were less stony indicating the river had lost a greater part of its energy and could thus transport and deposit only fine particles.

Comment [RA13]: Paragraph too long




Figure 2: Artifacts in (a) soil profile (b) in river banks of the wetland of Bamenda Municipality Cameroon

Classification

Based on the USDA soil Taxonomy and with approximate correlation of FAO/UNESCO soil legend, the near absence of diagnostic horizons qualified the soils for classification in the Entisol order of soil Taxonomy. The ustic moisture regime of the profiles coupled with gleyzation processes qualified the soils to be fitted into the suborder Fluvent, Great group Ustivluvet and subgroup Typic Ustifluent. FAO/ISRIC/ISSS (1998) allocated these alluvial soils to the group of azonal soils. Other correlations made include: alluvial soils, Regosols, Sols tropicaux recents; Sols mineraux bruts d'apport alluvial ou colluvial, Sols peu evolues non climatiques d'apport alluvial ou colluvial.

The FAO/UNESCO equivalent is Humi-umbric Fluvisol.

Physical and chemical properties of soil profiles

Studies of the seven soil profiles revealed that the physico-chemical properties varied considerably within and between the profiles (Table 3). Similar variations in hydric soils have been noticed in many parts of the world (Van Diepen, 1986, Eswaran and Cook, 1986, and Egbuchua and Ojeifo, 2007). The sand fractions were the dominant soil separates and irregularly distributed within the profiles with mean values of 69.5%. Silt contents were moderately distributed. Moderate values of silt could be allied to constant fluvial deposits and sedimentation processes. Clay contents of the soils were generally low and ranged from

10% in the profiles at Ntenefor (Below Foncha) and that at Mulang toward Army rescue to 20% in the Cg horizon of the Fuwambi – Ntahsen profile. The low clay content could be attributed to some extent to the processes of selective destruction of clay in the surface soil horizons and to ferrollysis. The irregular distribution of, and stratification of sand in these profile horizons suggest different periods of deposition of sediments rather than the deep weathered nature of parent materials. Similar findings in wetlands have been reported by Ojannuga (1991). In most of the profiles, the A horizons were dark in colors attributed to higher organic matter contents (Table 3). The organic matter content did not show regular decreases with depth often witnessed in most undisturbed soil profiles around the world (Yerima and Van Ranst, 2005b, Yerima *et al.*, 2009). This inexistence of definite trends of organic matter in the profiles could be ascribed to the continuous excavation activities and/or seasonal deposition of run-off sediments and urban wastes as evident in the profiles and river banks (Figure 2), respectively). These results are in concordance with reports by FAO-ISRIC-ISSS (1998) and Yerima and Van Ranst (2005b) who noted that fluvisol profiles have irregular distribution of organic matter. Total nitrogen showed similar pattern as organic matter. The pH (1:2.5 soil: water) of these soils ranged from very acidic (4.5) in the Ap horizon of the profile at Fuwambi – Ntahsen to moderately acidic (6.8) in the CAg horizon of the same profile. However, the profiles showed slight decreases of pH with depths which were not significant. Such variation is uncommon in stable ferralitic soils where pH decreases are always significant with depth increases (Yerima and Van Ranst, 2005b, Yerima *et al.*, 2009). Lower pH values observed in some surface horizons might be as a result of deposition of acidic substances through routine farm management practices such as the application of fertilizers, a common phenomenon of the area and/or from municipal swept offs and this was also reported by Asongwe *et al.* (2014). The very acidic nature of some of the samples might result to Al and Mn toxicity. This would affect the availability of available P. Beernaert and Bitondo (1992) reported that in soils with such acidic nature, Ca, Mg and Mb may also be deficient. To address the situation of acidity, ammonium sulphate and triple superphosphate fertilizers should be avoided. The EC values were high and ranged from 59 to 289 uS/cm with an average value of 203.3 uS/cm. The concentrations of Ca and Mg which dominated the exchange complex of the soils ranged from 0.20 to 5.13 and 0.12 to 3.80 cmol/kg, respectively. A potential local source of calcium might be relatively unweathered gneissic rocks rich in oligoclase (soda-lime plagioclase feldspar), hornblende (an amphibole containing calcium, magnesium, sodium, iron and aluminium) or augite (a pyroxene containing calcium, magnesium, iron and aluminium). Of particular importance, crushed or ground animal bones (bone meal) can be used as a slow-acting source of both calcium and phosphorus (Dagerskog *et al.*, 2010). All vegetable and bone waste can be re-used in various ways as fertilizers and, using processes that have proved successful in other African countries (Dagerskog *et al.*, 2010), this could be extended to include human faeces and urine. They respectively showed highly significant ($P < 0.01$) positive correlation coefficients of 0.776 and 0.780 with base saturation and CEC (Table 5). The sand/silt ratios were fairly constant but evidenced stratigraphic breaks in some profiles. According to Yerima and Van Ranst (2005b), in environments characterized by slowly flowing or stagnant water, stratification of deposits may be difficult to detect, and as such successive deposits will seem to be homogenous. This would have been the case of the fluvisols of the wetland gardens in Bamenda Municipality. Sombrek and Zonneveid (1971) reported that, silt/clay ratios also predicts ages of parent materials, young or old ones. Where the ratio is more than 0.15, the parent material is a young one and vice versa. In this case, the ratios exceeded 0.15, an indication that the parent materials of the Fluvisols in urban wetland gardens are young. The CEC of the soils ranged from low to high with an average value of 37.73 cmol/kg. The ECEC

showed a weak positive relationship ($r = 0.044$) with exchange acidity at the 5% probability level. Though CEC clay was not determined directly but calculated from CEC soil, the values ranged from 3.87 in the Apg horizon of the Fuwambi – Ntansen profile to 178.38 cmol/Kg in the Cg2 horizon of the profile at Mulang council junction. This predicts suiting mineralogy of the soils to be dominated by 1:1 minerals such as Kaolinites and sesquioxides with some limited interstratification with 2:1 minerals. The clay fraction of soil showed an insignificant ($p > 0.05$) positive correlation ($r = 0.302$) with organic matter at the 5% probability level (Table 5). The insignificant positive relationship is an indication that the distribution of organic matter does not depend on the clay content but on external factors possibly varying degrees of sedimentation.

Comment [RA14]: Review this value, perhaps they tried to mention 95%, which is the confidence level, because 5% is not equal to 0.05 and in case they refer it as a confidence level it is also incorrect

Comment [RA15]: Excessively long paragraph

UNDER PEER REVIEW

Table 3: Physico-chemical properties of soil profiles within wetland gardens of Bamenda Municipality

Comment [RA16]: include caption in footer

Horizon	Depth cm	Sand %	Silt %	Clay %	Sand/Silt	pH (1:2.5) H ₂ O KCl	EC uS/cm	OC %	OM %	Tot. N	C/N	Av. P mg/kg	Ca ²⁺ cmol/kg	Mg ²⁺ cmol/kg	K ⁺	Na ⁺	Σ Bases	Acidity meq/100g	Al ³⁺	H ⁺	CECsoil cmol/kg	CECclay cmol/kg	ECEC	B.S. CEC %	B.S EC %
Profile 1: Fuwambi-Ntahsen Quarter																									
Ap1	0 - 15	68	17	15	4.0	4.5 4.2	233	9.49	16.37	0.6	15.82	17	0.49	0.15	0.10	0.13	0.87	0.045	0	0.04	39.76	46.80	39.81	2.19	2.19
Apg	15 - 27	68	17	15	4.0	6.3 4.5	248	8.53	14.71	0.4	21.33	17	0.84	3.10	0.40	0.12	4.46	0.010	-	-	30.00	3.87	30.01	14.87	14.86
Cag	27 - 34	66	16	18	4.1	6.8 4.5	273	9.70	16.72	0.5	19.40	14	0.30	0.35	0.04	0.12	0.81	0.010	-	-	36.80	18.62	36.81	2.19	2.19
2CAg	34 - 59	69	15	16	4.6	6.4 4.4	244	10.25	17.67	0.5	20.50	19	0.34	0.17	0.05	0.10	0.66	0.010	-	-	41.20	36.63	41.21	1.60	1.60
2CAg2	59 - 83	70	16	14	4.4	6 4.3	262	7.89	13.61	0.5	15.78	60	0.43	0.20	0.05	0.10	0.78	0.010	-	-	50.16	164.57	50.17	1.56	1.55
Cg	83 - 120	72	8	20	9.0	5.6 4.2	219	7.44	12.82	0.2	37.18	7	0.44	0.36	0.03	0.14	0.97	0.070	-	-	33.93	41.43	34.00	2.85	2.85
Profile 6: Below Cow Slaughter																									
Apg	0 - 12	70	16	14	4.4	5.5 4.7	147	9.57	16.49	0.6	15.94	8	0.42	0.24	0.04	0.12	0.83	0.090	-	-	33.72	5.30	33.81	2.45	2.44
Cg	12 - 44	68	18	14	3.8	6.3 4.8	158	9.24	15.93	0.6	15.40	9	0.37	0.26	0.04	0.12	0.79	0.010	-	-	34.03	15.46	34.04	2.32	2.32
Profile 7: Below Slap																									
Apg	0 - 20	67	17	16	3.9	5.5 4.3	243	9.35	16.12	0.8	11.69	12	0.42	0.16	0.04	0.12	0.75	0.090	-	-	33.52	8.01	33.61	2.23	2.22
Cg	20 - 41	70	16	14	4.4	6.3 5.5	246	9.13	15.75	1.2	7.61	15	0.20	0.16	0.04	0.12	0.52	0.010	-	-	35.67	29.82	35.68	1.45	1.45
Profile 2: Ntenefor (Below Foncha)																									
Apg	0 - 20	80	10	10	8.0	6 4.9	134	9.00	15.51	0.6	15.00	14	0.54	0.26	0.04	0.12	0.96	0.010	-	-	34.67	36.36	34.68	2.77	2.77
ACg	20 - 30	82	8	10	10.3	6 4.9	137	8.73	15.05	0.4	21.83	13	0.20	0.35	0.04	0.12	0.71	0.010	-	-	33.72	36.22	33.73	2.10	2.10
Cg	30 - 38	81	9	10	9.0	5.7 4.6	132	8.00	13.80	0.3	26.68	8	5.13	1.12	0.04	0.12	6.41	0.090	-	-	35.57	79.74	35.66	18.02	17.97
Profile 3: Mulang -Ntahkah																									
Apg	0 - 41	63	18	19	3.5	5.7 4.5	59	9.13	15.75	0.6	15.22	13	0.54	0.26	0.04	0.09	0.93	0.060	-	-	34.13	13.89	34.19	2.71	2.71
Cg	41 - 70	62	20	18	3.1	6.2 4.6	68	9.54	16.44	0.5	19.08	10	0.20	0.16	0.04	0.11	0.51	0.010	-	-	35.77	16.02	35.78	1.41	1.41
Profile 4: Mulang Council junction																									
Ap	0 - 27	70	16	14	4.4	6.1 5.1	289	9.60	16.56	0.6	16.00	10	0.24	1.46	0.05	0.10	1.85	0.010	-	-	41.20	57.71	41.21	4.49	4.49
Cg1	27-38	62	28	10	2.2	5.9 5.0	256	8.00	13.79	0.6	13.33	27	1.34	3.80	0.05	0.10	5.29	0.090	-	-	36.96	93.80	37.05	14.31	14.28
Cg2	38 - 59	63	21	16	3.0	5.7 4.7	227	10.99	18.97	0.7	15.70	12	0.56	1.90	0.20	0.13	2.79	0.060	-	-	66.48	178.38	66.54	4.20	4.19
Profile 5: Mulang towards Army rescue																									
Ap	0 - 27	62	28	10	2.2	6.3 5.5	248	8.81	15.19	0.4	22.03	24	0.42	0.12	0.04	0.12	0.71	0.010	-	-	34.13	37.53	34.14	2.07	2.07
ACg1	27 - 37	64	26	10	2.5	6.2 4.4	214	8.92	15.38	0.5	17.84	11	0.77	0.26	0.04	0.12	1.19	0.010	-	-	35.57	48.16	35.58	3.35	3.35
ACg2	37 - 46	82	8	10	10.3	6 5.2	233	8.22	14.17	0.8	10.27	13	0.51	0.16	0.04	0.12	0.83	0.010	-	-	35.44	71.08	35.45	2.35	2.35

Table 4: Descriptive statistics for some physico-chemical properties of soil profiles in Bamenda Municipality

	N	Range	Minimum	Maximum	Mean	Std. Deviation	Variance
Sand	21	20.00	62.00	82.00	69.4762	6.60014	43.562
Silt	21	20.00	8.00	28.00	16.5714	5.96298	35.557
Clay	21	10.00	10.00	20.00	13.9524	3.30872	10.948
Sand /silt	21	8.10	2.20	10.30	5.0048	2.60796	6.801
pH water	21	2.30	4.50	6.80	5.9524	.46864	0.220
pH KCl	21	1.30	4.20	5.50	4.7048	.38791	0.150
EC	21	230.00	59.00	289.00	203.3333	66.63207	4439.833
OC	21	3.55	7.44	10.99	9.0252	.83786	0.702
OM	21	6.15	12.82	18.97	15.5619	1.44691	2.094
N	21	1.00	0.20	1.20	0.5667	.20575	0.042
CN	21	29.57	7.61	37.18	17.7919	6.20378	38.487
Av.P	21	53.00	7.00	60.00	15.8571	11.30171	127.729
Ca	21	4.93	0.20	5.13	0.7000	1.04752	1.097
Mg	21	3.68	0.12	3.80	0.7143	1.03059	1.062
K	21	.37	0.03	0.40	0.0690	0.08414	0.007
Na	21	.05	0.09	0.14	0.1162	0.01203	0.000
SBases	21	5.90	0.51	6.41	1.6010	1.68893	2.852
Acidity	21	0.08	0.01	0.09	0.0345	.03376	0.001
Al	1	0.00	0.00	0.00	0.0040	.	.
H	1	0.00	0.04	0.04	0.0410	.	.
CECsoil	21	36.48	30.00	66.48	37.7348	7.76672	60.322
CECclay	21	174.51	3.87	178.38	49.4952	47.24357	2231.955
ECEC	21	36.53	30.01	66.54	37.7695	7.76819	60.345
BSCEC	21	16.61	1.41	18.02	4.3567	4.86537	23.672
BSECEC	21	16.56	1.41	17.97	4.3505	4.85492	23.570

Comment [RA17]: declare acronyms in footer this table does not appear in results and discussion, delete or develop in results and discussion

UNDER

Table 5: Pearson correlation coefficients for some physico-chemical parameters of soil profiles in wetlands of Bamenda Municipality

	Sand	Silt	Clay	Sand/silt	pH H ₂ O	pH KCl	EC	OC	OM	N	C/N	Av. P	Ca	Mg	K	Na	ΣBases	Acidity	CECsoil	CECclay	ECEC	BSCEC	BSECEC	
Sand	1																							
Silt	-0.866**	1																						
Clay	-0.434*	-0.075	1																					
Sand/silt	0.937**	-0.903**	-0.242	1																				
pH H ₂ O	-0.083	0.134	-0.076	-0.107	1																			
pH KCl	0.138	0.144	-0.534*	0.045	0.370	1																		
EC	-0.148	0.205	-0.075	-0.207	0.147	0.101	1																	
OC	-0.373	0.245	0.302	-0.444*	0.044	-0.011	-0.008	1																
OM	-0.373	0.245	0.302	-0.445*	0.043	-0.010	-0.007	1.000**	1															
N	-0.069	0.102	-0.047	-0.204	-0.007	0.041*	0.175	0.345	0.346	1														
C/N	0.124	-0.267	0.234	0.332	-0.011	-0.382	-0.114	-0.352	-0.352	-0.869**	1													
Av. P	-0.132	0.222	-0.135	-0.218	0.080	-0.067	0.359	-0.297	-0.296	-0.024	-0.172	1												
Ca	0.323	-0.169	-0.341	0.275	-0.152	-0.079	-0.194	-0.352	-0.351	-0.315	0.300	-0.108	1											
Mg	-0.212	0.318	-0.152	-0.226	0.047	0.088	0.278	-0.128	-0.127	-0.108	0.004	0.080	0.271	1										
K	-0.158	0.106	0.124	-0.188	0.009	-0.153	0.213	0.125	0.127	-0.097	0.055	0.025	0.008	0.603**	1									
Na	0.238	-0.261	-0.005	0.322	-0.273	-0.082	0.111	-0.042	-0.042	-0.094	0.350	-0.409	0.040	-0.150	0.174	1								
ΣBases	0.065	0.093	-0.298	0.025	-0.068	-0.003	0.061	-0.291	-0.289	-0.267	0.194	-0.020	0.786**	0.807**	0.424	-0.050	1							
Acidity	-0.085	0.023	0.130	-0.007	-0.613**	-0.290	-0.153	-0.123	-0.123	-0.071	0.138	-0.197	0.456*	0.277	-0.082	0.112	0.449*	1						
CECsoil	-0.213	0.160	0.137	-0.219	-0.126	-0.103	0.261	0.0461*	0.463*	0.141	-0.146	0.323	-0.065	0.152	0.165	-0.007	0.060	0.039	1					
CECclay	0.012	0.095	-0.195	-0.001	-0.137	-0.010	0.300	-0.038	-0.035	0.009	-0.069	0.542*	0.200	0.282	0.071	-0.052	0.298	0.100	0.060	1				
ECEC	-0.213	0.160	0.137	-0.219	-0.129	-0.105	0.260	0.460*	0.463*	0.140	-0.145	0.322	-0.063	0.153	0.165	-0.006	0.062	0.044	1.000**	0.849**	1			
BSCEC	0.103	0.053	-0.302	0.058	-0.029	-0.015	0.035	-0.382	-0.381	-0.305	0.226	-0.035	0.776**	0.780**	0.451*	-0.056	0.979**	0.399	-0.123	0.141	-0.121	1		
BSECEC	0.103	0.054	-0.302	0.058	-0.028	-0.015	0.035	-0.382	-0.381	-0.305	0.226	-0.036	0.776**	0.780**	0.451*	-0.056	0.979**	0.398	-0.123	0.140	-0.121	1.000**	1	

*Correlation significant at 5%

**Correlation significant at 1

Comment [RA18]: In this regard, it would be convenient to include the ANOVA table of the correlation analysis and thus be able to know with certainty the possible statistical differences between variables, regardless of the coefficient of determination reached. This statistical procedure (ANOVA) is available in R (free software)

Physico-chemical properties of surface soil samples collected from the wetland gardens of Bamenda Municipality

Descriptions of adequacy of nutrient levels were done using critical values for nutrients and soil properties (Table 2).

The results of the physico-chemical properties (Table 6) of soils in the Urban and Peri-urban wetlands of the Bamenda Municipality under smallholder agricultural farms vary considerably. Ninety percent of the soils in the wetlands have a sandy loam texture while 10% are of the sandy clay-loam textural class. The clay contents of the soils ranged from 10% to 21% with an average of 16.7%. (Table 7). According to Mengel and Kirkby (1985), sites with high percentage of clay and silt are recommended for agricultural practices as they are capable of providing good aeration and retention and therefore supply of nutrients and water. However, these soils were poor in such parameters, predicting agronomic lapses. The soils of the area vary from acidic, through moderately acidic to slightly acidic. Average soil pH (H₂O) was 5.3 and 4.6 for pH (KCl). Generally, pH KCl range from 4.0 to 5.4 and are lower than those of pH water which ranged from 4.3 to 6.3. The variation of ΔpH ($\text{pH}_{\text{KCl}} - \text{pH}_{\text{H}_2\text{O}}$) was negative throughout. This indicates that the net charge on the exchange complex is negative, and thus exhibit cation exchange capacity. However, according to Yerima and Van Ranst (2005a), some tropical soils due to intensive rainfalls and weathering are dominated by positive charges with anion exchange capacity predominantly. Percent organic carbon ranged from 3.21% (Mile 4 market area) to 13.63% (Mulang 4 near houses) with an average value of 8.19 % in the entire area. The organic matter according to critical values by Beernaert and Bitondo (1992) (Table 2), varied from high to very high values, with a range from 5.67% to 23.50%. It had a weak positive correlation with the clay fraction of soil ($r = 0.218$) at the 5% probability level. This is an indication that the distribution of organic matter in the soil is not influenced by clay. This variation might be attributed to the constant addition of organic matter from varying anthropogenic activities (application of poultry manure, municipal wastes (Table 6-etc) and varying levels of stratification due to seasonal flooding. In the tropics, soil organic carbon is central to sustaining soil fertility on smallholder farms (Swift and Woomer 1993, Woomer *et al.*, 1994). In low-input agricultural systems in the tropics, it helps retain mineral nutrients (N, S, micronutrients) in the soil and make them available to plants in small amounts over many years as it is mineralized. In addition, soil organic carbon increases soil flora and fauna (associated with soil aggregation, improved infiltration of water and reduced soil erosion), complexes toxic Al and manganese (Mn) ions (leading to better rooting), increases the buffering capacity on low-activity clay soils, and increases water- holding capacity (Woomer *et al.* 1994). Continuous cropping, with its associated tillage practices, provokes an initial rapid decline in soil organic matter, which then stabilizes at a low level (Woomer *et al.*, 1994).

Total N ranged from 0.3 to 0.8% (Table 7). Landon (1991) reported that for tropical soils, total nitrogen content of 0.13% is sufficient. Nitrogen is highly mobile and easily lost and vegetables need high quantities. This necessitates high application of nitrogen fertilizers to maintain the production of vegetable on the wetland which are already vulnerable given that they are dominated by the sand fraction. The organic carbon was perfectly correlated to organic matter ($r = 1.00$) (Table 8). The C/N ratio varying from 5.58 to 36.03 indicated that, the soils range from good to poor. Despite of the fact that the soil are rich in organic carbon, the very high C/N ratio witnessed in some areas predicts difficulties in mineralization which could be ascribed to water stagnation. Likewise, farm specific practices might have also influenced the inconsistent pattern in mineralisation. Areas characterized by rapid mineralization would result to high nitrogen losses which necessitate high nitrogen fertilization, a constraint to peasant farmers caused by poverty. Generally, a majority of the

Comment [RA19]: It is very important for the understanding of the study to locate the graphs and tables immediately after the first call or mention, this aspect is not fulfilled in the present proposal, please correct

soils had C/N ratios that were less than 25. According to Mengel and Kirkby (1987), soils with such C/N ratios (less than 25) are ameliorating and the litter are of the mull type.

Available P is associated to organic carbon. It ranged from 8 mg/Kg to 76 mg/Kg. Available P concentrations lower than 16 mg/Kg in soils are considered low to ensure adequate phosphate supply to most plants (Landon, 1991). The low P in some of the soils could be related to the rapid loss due to fast mineralization rate of organic matter given by low values of C/N values. The availability of phosphorus might also be limited due to the nature of parent material that are generally basaltic in this area, and probably high sorption due to surface mineralogy (Clulombe *et al.*, 1996).

Ca and Mg dominates the exchange complex but their concentrations were low ranging from 0.20 to 0.54 cmol/Kg for Ca (with a variance of 0.015) and 0.37 to 0.68 cmol/Kg for Mg (with a variance of 0.011). According to Landon (1991) deficiencies of Ca are normal in soils with $\text{pH} \leq 5.5$ which have been witness in some sites of this study. Continuous cultivation without returning residues to soil depletes these nutrients and thus low future yields. Major sources of magnesium in soils include amphiboles, olivine, pyroxene, dolomites and clay minerals (Todd, 1980). According to the later, dolomites have a high concentration of magnesium while clay minerals have low concentration. Ca and Mg showed a highly significant positive correlation ($r = 0.869$ and $r = 0.780$) with exchangeable bases at the 1% probability level (Table 8). The Al concentration ranged from 0.02 to 0.05 meq/100g. Within the exchanged acidity, it concentrations were significantly ($r = 0.710$) lower than that of H, at the 5% probability level. This is an indication that the sources of charges on the exchange colloids displaced by neutral salts are pH dependent.

The CEC of the soil according to the critical values of soil nutrients varied from low to medium, ranging from 29.96 to 48.99 cmol/Kg of soil with a standard deviation of 4.52347. These soils could thus have few weatherable minerals desiring nutrient application when extensively cultivated. Nkona (1998) reported that ECEC values of 4 meq/100g of soil marks the minimum limits. The soils in this regard have moderate potential to hold nutrient and avoid nutrients lost. It is thus imperative to raise the pH of the soil in areas of low pH in order to harness this potential.

Base saturation CEC for the soils were low ranging from 1.79% at Ngomegham to 4.02% at Foncha right of the road. This parameter is a good indicator of soil fertility in terms of the availability of nutrient elements. An ideal soil should have the exchange complex saturated to 76, 18, 6 by the elements Ca/Mg/K, respectively (Yerima and Van Ranst, 2005a). In this study, Ca, Mg, and K occupied respectively 38.33%, 56.98%, and 4.68% on the absorbed complex, necessitating amendment by Ca 1.7894 cmol/kg Mg 0 cmol/kg values for K. CEC clay was not directly determined but was calculated from CEC soil. It ranged from 3.2 to 7.5 cmol/Kg. This indicates that the mineralogy of the soil is dominated by low activity clays of the 1:1 type and sesquioxides. The Na content was fairly constant throughout the area with 1.3 value.

Table 6: Physico-chemical properties of surface soil samples, poultry manure and Municipal waste within the wetlands of the Bamenda Municipality

No	Site	Sand %	Silt %	Clay %	pH (1:2.5) H ₂ O KCl	EC uS/cm	OC	OM %	Tot. N	C/N	Av. P mg/Kg	Ca ²⁺	Mg ²⁺	K ⁺ cmol/Kg	Na ⁺	ΣBases	Exch. Acid, meq/100g	Al ³⁺	H ⁺	CECsoil	CECclay	ECEC cmol/Kg	B.S. CEC %	B.S. ECEC	
1	Fuwambi Near Ntasen	56	24	20	4.6	4.3	235	5.88	10.13	0.8	7.34	18	0.31	0.56	0.03	0.12	1.02	0.45	0.04	0.41	32.30	60.21	32.75	3.16	3.11
2	Fuwambi near GTTC	58	24	18	5.6	4.7	212	7.52	12.96	0.6	12.53	8	0.20	0.37	0.04	0.13	0.73	0.09	-	-	33.93	44.44	34.02	2.16	2.15
3	Slap 1	58	21	21	4.5	4.3	230	10.58	18.23	0.6	17.63	14	0.31	0.46	0.05	0.12	0.94	0.4	-	-	37.62	5.49	38.02	2.51	2.48
4	Slap 2	57	22	21	6.3	5.2	197	9.40	16.21	0.7	13.43	12	0.20	0.56	0.04	0.13	0.92	0.01	-	-	35.98	16.98	35.99	2.57	2.56
5	Slap 3	58	20	20	5.3	4.0	237	10.81	18.64	0.3	36.03	22	0.20	0.52	0.04	0.13	0.88	0.4	-	-	38.04	3.81	38.44	2.32	2.30
6	Slap 4	55	26	19	4.5	4.1	247	9.4	16.21	0.6	15.67	33	0.31	0.37	0.06	0.15	0.89	0.47	0.05	0.42	34.34	10.14	34.81	2.59	2.56
7	Slap 5	56	24	20	4.3	4.0	248	7.52	12.96	0.3	25.07	17	0.54	0.6	0.04	0.14	1.32	0.49	0.05	0.44	35.77	49.21	36.26	3.69	3.64
8	Mile 4 market	57	23	20	5.2	4.7	233	3.29	5.67	0.5	6.58	19	0.31	0.56	0.04	0.13	1.04	0.1	0.04	0.06	32.60	106.26	32.70	3.19	3.18
9	Foncha right of road	63	25	12	6.3	4.7	113	8.23	14.18	0.3	27.42	22	0.54	0.68	0.04	0.13	1.39	0.01	-	-	34.44	50.67	34.45	4.02	4.02
10	Foncha left of road	64	26	10	5.7	4.5	105	10.11	17.42	0.4	25.26	17	0.20	0.37	0.03	0.13	0.72	0.06	-	-	35.46	6.22	35.52	2.03	2.03
11	Ndamukong	63	26	11	6.0	4.8	167	4.70	8.10	0.7	6.71	13	0.31	0.56	0.04	0.13	1.04	0.09	-	-	33.93	161.11	34.02	3.06	3.05
12	Ntahkah inn	63	18	19	5.7	4.5	78	8.23	14.18	0.6	13.71	13	0.54	0.46	0.04	0.13	1.17	0.06	-	-	34.13	30.38	34.19	3.41	3.41
13	Ntahkah out	61	19	20	5.6	4.8	127	4.47	7.70	0.7	6.38	11	0.31	0.37	0.04	0.13	0.85	0.09	-	-	34.34	94.71	34.43	2.46	2.46
14	Ntahkah before bridge	62	18	20	5.1	4.5	130	10.58	18.23	0.3	35.25	20	0.54	0.68	0.04	0.13	1.39	0.3	0.03	0.27	37.74	6.38	38.04	3.67	3.64
15	Mulan council junction	79	10	11	5.4	5.0	458	6.58	11.34	0.4	16.45	76	0.31	0.44	0.04	0.13	0.92	0.09	0.02	0.07	34.54	107.77	34.63	2.65	2.64
16	Mulang left of road	68	17	15	5.8	5.4	279	9.40	16.21	0.6	15.67	25	0.20	0.56	0.02	0.13	0.90	0.05	-	-	34.24	12.16	34.29	2.63	2.63
17	Mulang middle	80	10	10	4.9	4.3	437	4.47	7.70	0.8	5.58	17	0.54	0.46	0.04	0.13	1.17	0.3	0.03	0.27	32.39	169.96	32.69	3.60	3.56
18	Mulang 4 near houses	62	20	18	4.6	4.1	273	13.63	23.50	0.6	22.72	16	0.31	0.66	0.04	0.13	1.14	0.4	0.04	0.36	48.99	11.05	49.39	2.32	2.30
19	Army Rescue	62	18	20	5.4	4.9	269	9.17	15.80	0.6	15.28	8	0.31	0.56	0.05	0.12	1.04	0.10	0.03	0.07	39.06	10.61	39.16	2.66	2.66
20	Ngomegham	80	10	10	6.0	5.1	140	6.82	11.75	0.5	13.63	20	0.31	0.37	0.04	0.12	0.84	0.01	-	-	47.00	116.59	47.01	1.79	1.79
21	Mbelewa	67	18	15	4.5	4.2	57	11.28	19.45	0.7	16.11	12	0.31	0.4	0.07	0.12	0.87	0.40	0.04	0.36	29.96	1.72	30.36	2.90	2.87
22	Poultry manure	-	-	-	7.4	7.1	1500	23.50	40.51	29.00	0.81	130	24960.00	8043.3	13442.00	1329.00	47774.3	0.002	-	-	50.32	-	50.32	94940.98	94937.20
23	Municipal solid waste	-	-	-	7.6	5.5	5201	47.71	82.24	17.20	2.77	112	21961.00	7698.0	13711.00	14213.00	57583.0	0.004	-	-	44.43	-	44.43	129603.87	129592.20

Table 7: Descriptive statistics of surface soil samples from the Wetland gardens of Bamenda Municipality

Descriptive Statistics								
	N	Range	Minimum	Maximum	Mean		Std. Deviation	Variance
	Statistic	Statistic	Statistic	Statistic	Statistic	Std. Error	Statistic	Statistic
Sand	21	25.00	55.00	80.00	63.2857	1.68345	7.71455	59.514
Silt	21	16.00	10.00	26.00	19.9524	1.11372	5.10369	26.048
Clay	21	11.00	10.00	21.00	16.6667	0.91894	4.21110	17.733
pHwater	21	2.00	4.30	6.30	5.3000	0.13680	0.62690	0.393
pHKCl	21	1.40	4.00	5.40	4.5952	0.08521	0.39048	0.152
EC	21	401.00	57.00	458.00	212.9524	22.44756	102.86762	10581.748
OC	21	10.34	3.29	13.63	8.1938	0.57950	2.65558	7.052
OM	21	17.83	5.67	23.50	14.1224	0.99936	4.57965	20.973
N	21	0.50	0.30	0.80	0.5524	0.03560	0.16315	0.027
CN	21	30.45	5.58	36.03	16.8786	1.94166	8.89779	79.171
AV.P	21	68.00	8.00	76.00	19.6667	3.09172	14.16804	200.733
Ca	21	0.34	0.20	0.54	0.3386	0.02707	0.12407	0.015
Mg	21	0.31	0.37	0.68	0.5033	0.02315	0.10608	0.011
K	21	0.05	0.02	0.07	0.0414	0.00221	0.01014	0.000
Na	21	0.03	0.12	0.15	0.1290	0.00153	0.00700	0.000
SBases	21	0.67	0.72	1.39	1.0086	0.04245	0.19453	0.038
Acidity	21	0.48	0.01	0.49	0.2081	0.03893	0.17840	0.032
Al	10	0.03	0.02	0.05	0.0370	0.00300	0.00949	0.000
H	10	0.38	0.06	0.44	0.2730	0.04844	0.15319	0.023
CECsol	21	19.03	29.96	48.99	36.0381	0.98710	4.52347	20.462
CECclay	21	168.24	1.72	169.96	51.2319	11.71137	53.66824	2880.280
ECEC	21	19.03	30.36	49.39	36.2462	0.98819	4.52846	20.507
BSCEC	21	2.23	1.79	4.02	2.8281	0.13132	0.60180	0.362
BSECEC	21	2.23	1.79	4.02	2.8114	0.12999	0.59570	0.355
Valid N (listwise)	10							

Table 8: Pearson correlation coefficient of surface soil samples from wetland gardens in Bamenda Municipality significant at the 5% probability

Comment [RA20]: Idem to the indication regarding the ANOVA of the correlation

Sand	Silt	Clay	pH _{H₂O}	pH _{KCl}	EC	OC	OM	N	CN	AV.P	Ca	Mg	K	Na	ΣBases	Acidity	Al	H	CEC _{sol}	CEC _{clay}	ECEC	BSCEC	BSECEC
------	------	------	------------------------------	-------------------	----	----	----	---	----	------	----	----	---	----	--------	---------	----	---	--------------------	---------------------	------	-------	--------

UNDER PEER REVIEW

*correlation significant at the 95% probability level, **correlation significant at the 1% probability level

Variability of surface soil properties in the wetlands of Bamenda Municipality

Table 9 show the mean physico-chemical properties of surface soil properties in the three major laps assessed. The coefficient of variations (CV %) was used to evaluate the variability of the soils. Similar groupings have been used by Tabi and Ogunkunle (2007). According to Ogunkunle, (1993), Tabi and Ogunkunle (2007), Tabi *et al.* (2012), CV values ranging 0-15% are considered slightly variable, 15-35% moderately variable, while > 35% they are considered highly variable (Table 10). Soil pH (H₂O or KCl), percent sand, CEC soil, and Na were consistently slightly variable. Tabi and Ogunkunle (2007) similarly reported slight variability of soil pH for Alfisols in Southern Nigeria. In a like manner for vertisols under rice cultivation in the Logone flood plain of Northern Cameroon slight variability was also obtained (Tabi *et al.*, 2012). Soil pH is one of the most very important parameter that influences the availability of nutrients and minor changes in pH units have significant effects on nutrient availability. The variability of Silt, clays, EC, ECEC, and Al³⁺, were slight to moderate (dominated by the moderate) class. This could be attributed to variation in levels of alluvial materials received. Also, variation in Chrono sequences of materials that have been subjected to different intensities of weathering could have a significant effect on these physical parameters. All these factors have significant implications on water and nutrient management of the wetlands of Bamenda Municipality. The exchangeable bases were moderately variable in a greater portions of the farms, probably associated to variation of soil amendment activities. The bases moderate variation implies that a single policy such as the application of organic or inorganic fertilizer will not be sufficient to properly manage the farms of the area. Organic carbon, organic matter, available P, exchange acidity, CEC clay, BSCEC, BSECEC, H⁺, Tot. N, were moderately to highly variable. Most of these parameter owe a lot to organic materials. Natural denudational activities inclusive, urban swept off, farming activities such as residue management, crop species and land preparation/management practices could have influence the nature and management of soil properties of the area. The available phosphorus and base saturation which is equally slightly variable reflects pH variability.

Comment [RA21]: percentage

Comment [RA22]: paragraph too long, correct

Sources of variation of soil properties.

The correlation coefficients were used to group various soil properties in each side. The correlation coefficients of 21 soil properties examined are shown in Table 8. The coefficients ranged between -0.950 to 0.999. Even though some of the correlations were highly significant ($p < 0.01$), most of them have correlation values less than 0.5. The highest positive correlation coefficients were observed between BSECEC and BSCEC (1.00), ECEC and CECsoil (0.999), exch. Acidity and H⁺ (0.999), exch. Acidity and Al (0.739), BSCEC and exchangeable bases (0.866), Ca and BSCEC (0.833), Ca and exchangeable bases (0.869), Ca and BSCEC (0.834), Mg and exchangeable bases (0.585), OC and CN (0.687), OM and CN (0.687) all at the one percent confidence interval. The high correlation value between OC and CN, could be an indication that they mainly contributed to the soils of this area from a single source making management of this parameter easier when identified. The highest negative correlation coefficients were obtained between pH H₂O and H⁺ (-0.950), pH H₂O and Al (-0.827), and pH H₂O and acidity (-0.919) which was significant at the 1 % confident interval. Similarly at the 1% probability level, sand equally had a negative correlation with Al content of the soils ($r = -0.768$). The negative correlation could be attributed to the facts that as the soil becomes sandier, very little colloids are left in place for retention of Al which is toxic to the plants. The data of 24 variables considered in this study were also subjected to R-mode analysis using the six-factor model, which accounted for 96.240% of the total data variance. The resulting varimax is summarized in Table 11. The computations were performed by means of the SPSS computer software package version 20.0. Only variables with loadings > 0.5 were considered significant. Contributions of the various principal components (PCI) were 20.896% (PC1), 18.655 % (PC2), 17.125% (PC3), 16.562% (PC4), 14.9837% (PC5), and 8.020% (PC6).

Table 9: Physico-chemical properties of surface soil properties at the Mile Four, Foncha and Mulang Laps of the Bamenda Municipality

Site	Sand Silt Clay %			pH(1:2.5) H2O KCl		EC uS/cm	OC	OM	Tot. N	C/N	Av. P mg/Kg	Ca ²⁺	Mg ²⁺	K ⁺	Na ⁺	ΣBases	Exch. Acid.	Al ³⁺	H ⁺	CECsoil	CECclay	ECEC	B.S. CEC	B.S. ECEC
								%									meq/100g	meq/100g		cmol/Kg		%		
Mile Four Lap	57	23	20	5.04	4.41	230	8.05	13.88	0.55	16.8	18	0.30	0.50	0.04	0.1305	0.97	0.3013	0.045	0.333	35.07	37.07	35.37	2.77	2.75
Foncha Lap	63	22	15	5.73	4.63	120	7.72	13.30	0.50	19.1	16	0.41	0.52	0.04	0.1265	1.09	0.1017	0.030	0.270	35.01	58.25	35.11	3.11	3.10
Mulang Lap	72	14	14	5.35	4.87	309	8.34	14.38	0.58	14.9	27	0.33	0.51	0.04	0.1239	1.00	0.1583	0.030	0.193	39.37	71.36	39.53	2.61	2.60

Table 10 Variability classes of soil properties for the different sites

Site	Least variable (CV<15%)	Moderately Variable (15 < CV ≤ 35%)	Highly Variable CV> 35%)
Mile Four area	Sand, silt, clay pH _{H2O} , pH _{KCl} , EC, Na ⁺ , Al ³⁺ , CECsoil, ECEC	OC, OM, Tot.N, Ca ²⁺ , Mg ²⁺ , K ⁺ , ΣBases, BSCEC, BSECEC.	C/N, Av. P, Exch.acidity, H ⁺ , CECclay.
Foncha area	Sand, pH _{H2O} , pH _{KCl} , K ⁺ , Na ⁺ , CECsoil, ECEC.	Silt, clay, EC, OC, OM, Av.P, Mg ²⁺ , ΣBases, BSCEC, BSECEC.	Tot. N, C/N, Ca ²⁺ , Exch. Acidity, ECECclay
Mulang	Sand, pH _{H2O} , pH _{KCl} , Na ⁺	Silt, Clay, Tot. N, Ca ²⁺ , Mg ²⁺ , K ⁺ , Na ⁺ , ΣBases, Al ³⁺ , CEC sol, BSCEC	EC, OC, OM, C/N, Av.P, Exch. Acidity, H ⁺ , CECclay, BSCEC

The 5 PCs identified constitute the minimum dataset required to group fluvisols for vegetable cultivation in the Wetlands of Bamenda Municipality. The factors (PCI) extracted from Table 11 are as follows (Table 12):

Factor 1 was made up of pH H₂O, pH KCl, Exch. acidity, Al, and H. This factor is an acidity factor because of the high negative loading (- 0.949) for pH water and (- 0.949) pH KCl and high positive loading of exch. Acidity (0.945), H⁺ (0.942) and Al³⁺ (0.738). Acidity of this environment is highly influenced by acid dependent charges as opposed to isomorphic substitution. The source of this acidity would be anthropogenic from the urban area.

Factor 2 constituted BSCEC, BSECEC, ΣBases, Ca, Mg. The principal component was named the base status factor because it had a high positive loading on base saturation CEC (0.930), base saturation ECEC (0.933), sum of bases (0.871), exchangeable Ca (0.916), and moderate loading of Mg (0.508). Same naming was given to soils of the Logone and Chari plain in the northern region of Cameroon under rice cultivation with similar constitution (Tabi *et al.*, 2012).

Factor 3 was composed of sand, silt, clay, Al, and EC. This component was termed weathering and associated moisture retention factor because of a high positive loading clay (0.950) greater than silt (0.901), moderate positive loading of Al (0.616) and a moderate negative loading of EC (-0.579).

Factor 4 grouped EC, OC, OM, C/N, K, and CECclay: the factor was ascribed an organic matter factor because of a high (same) positive loading for organic matter and organic carbon content (0.862) and organic matter quality (0.732) quality (C/N ratio), moderate positive loading from K (0.626); but a high negative loading from CECclay (-0.811). The major source of this component would be the breakdown of natural vegetation in the area.

Factor 5 converged Mg, CECsoil, ECEC, K. The factor had high positive loading from CECsoil (0.937), ECEC (0.937), moderate loading from Mg (0.719) and a moderate negative loading from K (-0.540). This factor was described as a mineral neo-synthesis related component derived mainly from the deposition of Mg eroded from upland in the wetland which gradually replaces K in interlayers of micaceous minerals in the wetlands.

Factor 6 grouped N, available P, and Na. this is a dispersal-mineralization factor influenced by anthropogenic activities of the International Soap Factory (high sodium released) and the Cow slaughter house (high organic matter containing nitrogen which is rapidly mineralized due to dispersion caused Na). The factor had high positive loading from Na (0.818), moderate loading from available phosphorus (0.690) but a negative yet moderate loading from total N (-0.690). Potentially mineralizable nitrogen is an important measure of N supplying capacity in wetland soils, which is most often not calculated. The findings of this component are in conformity with the reports of Mengel and Kirkby (1985), and Tabi *et al.* (2012) that nitrogen is one of the most important factors controlling potentially mineralizable N in wetland soils. Generally, the results reported here agree with those of other authors. Salami *et al.* (2011) identified potential fertility, available phosphorus, organic matter, acidity and sand-silt as factors responsible for soil fertility variation in the northern guinea savanna agro-ecological zone of Nigeria. Kosaki and Juo (1989) identified inherent fertility status (represented by Mg, Ca, K, sand, silt and clay) and available phosphorus as major factors causing soil variation in wetland fields in southwestern Nigeria. In fact, no sampling site was completely homogeneous in relation to soil management units

From this study, it is concluded that the fluvisols of the Wetland of Bamenda Municipality have been highly influenced by human activities. The soils are developed from young alluvio-colluvial material of granitic origin. The organic matter just as other physico-chemical properties varied irregularly down the profile. The surface soils of the wetland

gardens were slightly acidic to neutral, of low organic matter and nitrogen contents, and moderate to high exchangeable bases. Except pH that was slightly variable, soil properties were moderately variable to highly variable. Six principal components (PCs) grouping soils in management units explained 96.240% of the variations observed in the soil properties. The PCs were: base status, organic matter, weathering and moisture retention, acidity, dispersal and N-mineralization, and mineral neo-synthesis factors. The PCs constitute a minimum dataset required to group soils in the Bamenda wetland gardens. We recommend that a detailed mapping of soil properties be carried out for the establishment of a soil fertility map for Bamenda Municipality; fertilizer recommendations should be established for identified soil management units; and farming systems which favors buildup of soil organic matter and use of animal manure should be encouraged. Soil management practices defined for identified units instead of a common management for all units will boost and sustain vegetable yield. Soil characterization and the implementation of some of the findings of these studies are believed to result in increased food production per unit area.

Table 12: Total Variance explained from Physico-chemical properties of wetland soils in the wetland gardens of Bamenda Municipality

Component	Initial Eigenvalues			Extraction Sums of Squared Loadings			Rotation Sums of Squared Loadings		
	Total	% of Variance	Cumulative %	Total	% of Variance	Cumulative %	Total	% of Variance	Cumulative %
1	7.501	31.254	31.254	7.501	31.254	31.254	5.015	20.896	20.896
2	4.933	20.553	51.807	4.933	20.553	51.807	4.477	18.655	39.550
3	4.437	18.486	70.293	4.437	18.486	70.293	4.110	17.125	56.675
4	2.536	10.567	80.860	2.536	10.567	80.860	3.975	16.562	73.237
5	1.921	8.004	88.864	1.921	8.004	88.864	3.596	14.983	88.220
6	1.770	7.377	96.240	1.770	7.377	96.240	1.925	8.020	96.240
7	0.623	2.595	98.836						
8	0.198	0.827	99.663						
9	0.081	0.337	100.000						
10	6.649E-016	2.770E-015	100.000						
11	4.939E-016	2.058E-015	100.000						
12	3.993E-016	1.664E-015	100.000						
13	3.097E-016	1.290E-015	100.000						
14	2.069E-016	8.621E-016	100.000						
15	1.530E-016	6.376E-016	100.000						
16	5.699E-017	2.375E-016	100.000						
17	4.010E-017	1.671E-016	100.000						
18	-5.979E-017	-2.491E-016	100.000						
19	-1.267E-016	-5.281E-016	100.000						
20	-1.565E-016	-6.519E-016	100.000						
21	-1.663E-016	-6.930E-016	100.000						
22	-2.446E-016	-1.019E-015	100.000						
23	-3.771E-016	-1.571E-015	100.000						
24	-6.116E-016	-2.548E-015	100.000						

Extraction Method: Principal Component Analysis

Comment [RA23]: You do not necessarily have to extend the ACP up to six, there is a consensus that explanation of the variance <70% is sufficient, so you could try leaving it up to four components

Comment [RA24]: Likewise, you included without discrimination or cleaning of data any composite variables within the CPA, which is not correct, you mention that you developed the procedure in the SPSS software, this software has a highly valued indicator in the CPA analysis that measures the incorrectness of the variables based on 1, that is, the closer it is to one, the greater will be the correlation of the variables and the more robust will be the statistical procedure, this indicator is called KMO, so I urge you to do a new procedure, eliminating the composite variables and considering the KMO value, so your results will be reliable

Comment [RA25]: I consider that it is not necessary to show this table as a result, the following table is more important than it is and actually shows the resolution of the operation so it should be removed

Formatted: Highlight

References

- Asongwe GA, Yerima BPK, and Tening AS. Vegetable production and the livelihood of farmers in Bamenda Municipality, Cameroon. *International Journal of Current Microbiology and Applied Sciences*. 2014; 3:682 – 700.
- Bationo AJ, Kihara B, Vanlauwe B, Waswa P and Kimetu J. Soil organic carbon dynamics, functions and management in West African agro- ecosystems. *Agricultural Systems*. 2006.
- Bernaert F, and Bitondo D. A simple and Practical Method to Evaluate Analytical Data of Soil Profiles. CUDs, Soil Science Department. Belgia Cooperation Dschang Cameroon, 1992; 65PP.
- Coulombe CE, Dixon B, Wilding L P. Mineralogy and Chemistry of Vertisols. *Developments in Soil Science*. 1996; 24:115-200.
- Dagerskog L, Coulibaly C, and Ouandaogo I. The emerging market of treated human excreta in Ouagadougou. *Urban Agriculture*, 23. Resource Centres on Urban Agriculture and Food Security (www.ruaf.org). 2010
- Deckers J, and Spaargaren O. The State of the Art regarding the World Reference Base for Soil Resources: WRB towards Montpellier 1998. *Proceedings of the Int. WRB Workshop*, Pretoria. 1996.
- Driessen PM, and Dudal R. The major soils of the world. *Lecture notes on their geography, formation, properties and use*. Agricultural University Wageningen, the Netherlands, and Katholieke Universiteit Leuven, Belgium. 1991; 310 p.
- Duchaufour PH. *Pedologie*. Masson ed. 1988; 224 p.
- Dudal R. Dark clay soils of tropical and subtropical regions. *FAO Agric. Development Paper* 83. Rome, Italy. 1965; 161 p.
- Egbuchua CN and Ojeifo IM. Pedogenetic characterization of some wetland soils in Delta State. Ph.D. Thesis Delta State University, Abraka, Nigeria, 2007.
- Eswaran H and Cook T. Classification of wetland soils in Soil Taxonomy. *Wetland soils: Characterization, classification and utilization*. IRRI, Los Banos, Philippines. 1986.
- FAO/ISRIC/ISSS. *World Reference Base for Soil Resources*. FAO, Rome, 1998.
- FAO-WRB, *World reference base for soil resource. A framework for international for soil classification, correlation and communication*. *World Soil Resources Reports*, 2006; 103: 145-159.
- GP-DERUDEP. *Grassfield Participatory: Decentralized and Rural Development Project*. 2006. Baseline study of the North West Province. SIRDEP Bamenda Cameroon. 298pp.

Comment [RA29]: The bibliography is outdated, the following is represented in periods of ten years, which is the representativeness of the bibliographic elements:

17.9% (2010-2020)
28.2% (1999-2009)
33% (1988-1998)
12.8% (1977-1987)
7.69% (1976-xx)

- Haase D, and Neumeister H. Anthropogenic impacts on fluvisols in German Floodplains: Ecological processes in soils and methods of investigation. *International Agrophysics* 2001; 15:19-26.
- Kometa SS. Wetlands Exploitation along the Bafoussam – Bamenda road Axis of the Western Highlands of Cameroon. *Journal of Human Ecology*, 2013; 41(1): 25-32
- Kosaki T, and Juo ASR. Multivariate approach to grouping soils in small fields. Extraction of factors causing soil variation by PCA. *Soil Sci. Plant Nutr.* 1989; 38(3):469-477.
- Landon J. Booker Tropical Soil Manual: A handbook for soil survey and agricultural land evaluation in the tropics and subtropics, N. York, 1991; 450p
- Mengel K and Kirkby EA. Principles of plant nutrition. International Potash Institute. Worblaufen-Ben Switzerland. 1987; 687pp.
- Moloua EL, and Lambi CM. Climate, Hydrology, and Water Resources in Cameroon. CEEPA, Pretoria, 2006, 37pp.
- Ndenecho EN. Biological Resource Exploitation in Cameroon: From crises to sustainable management. Unique Printers Bamenda Cameroon: 2005, 181pp.
- Ngachie N. A general assessment of soil resources and soil fertility constraint in Cameroon on the basis of FAO-UNESCO soil map analysis. *Tropicultura*, 1992; 10(10):61-63.
- Ogunkule AO. Variation of some soil properties along two topo sequence on quartzite schist and banded gneiss in South-western Nigeria. *Geo-journal*, 1993; 30 (40): 399-402.
- Pauwels JM, Van Ranst E, Verloo, M and Mvondo Ze A. Methodes d'analyses de sols et de plantes, equipements, gestion de stocks de verrerie et de produits chimiques. A. G. Bulding. Bruxelles, 1992.
- Phil-Eze PO. Variability of soil properties related to vegetation cover in a tropical rainforest landscape. *Journal of Geography and Regional Planning*, 2010; 3(7): 177-184.
- Ramsar. Wetlands and Agriculture, Partners for Change. Ramsar publication for the World Wetland Day 2014, Gland Switzerland, 2014; 16pp.
- Salami BT, Okogun JA and Sanginga N. Delineation of management zones by classification of soil physico-chemical properties in the northern savanna of Nigeria. *African Journal of Agricultural Research*, 2011, 6(6): 1572-1579.
- Sombrok, W.G and I. S. Zoineveld. 1971. Ancient dune fields and fluvial deposit in Rima-Sokoto River basin.
- Swift M J, and Woome P. Organic matter and the sustainability of agricultural systems: Definition and measurement. In K. Mulongoy and R. Merckx (eds.), *Soil Organic Matter Dynamics and Sustainability of Tropical Agriculture*. Chichester, U.K.: Wiley-Sayce. 1993, 3-18
- Tabi FO, and Ogunkunle AO. Spatial variation of some soil physico-chemical properties of an Alfisol in Southwestern Nigeria. *Nigeria Journal of Soil and Environmental Resources*. 2007; 7:82-91.
- Tabi, FO, M. Omoko, A. Boukong, A. D. Mvondo Ze, D. Bitondo, and Fuh-Che, C. 2012. Evaluation of Lowland rice (*Oryza sativa*) production system and management

recommendations for Logone and Chari flood plain-Republic of Cameroon. *Agricultural Science Research Journals*, 2(5): 261-273.

Todd DK. *Groundwater Hydrology*. John Wiley and Sons. New York, 1980; 535pp.

USDA. United States Department of Agriculture.. Munsell soil colour chart. Macbeth Division of Kollmorgen Corporation 1441 North Calvert Street Baltimore, Maryland 1975; 218.

Van – Diepen CA. Wetland soils of the World, their characteristics and distribution in the FAO/UNESCO approach. Proceedings of an international workshop on wetland soils. 25th March-5th April, 1984 Los Banos, Philippines. 1986

Wilding LP and Rehage JA. Pedogenesis of soils with aquic moisture regimes. Proceedings of International workshop on wetlands: Characterization, classification and utilization IRRI- Los Banos, Philippines. 1986

Woomer PL, Martin A, Albrecht A, Resck DVS, and Scharpenseel HW.. The importance and management of soil organic matter in the tropics. In P.L. Woomer and M.J. Swift (eds.), *The Biological Management of Tropical Soil Fertility*. Chichester, U.K.: Wiley-Sayce. 1994; 47- 80.

Yerima BPK and Van Ranst E. Major soil classification systems used in the Tropics: Soils of Cameroon. TRAFFORD Publishers, Victoria Canada. 2005b; 295pp.

Yerima BPK and Van Ranst E. Introduction to Soil Science, soil of the Tropics. TRAFFORD Publishers, Victoria, 2005a; 397pp.

Yerima BPK, Van Ranst E, and Verdoot A. Use of Correlation Relationships to Enhance Understanding of Pedogenic Processes and use potential of Vertisols and Vertic Inceptisols of the Bale Mountain Area of Ethiopia. *Tropicultura*, 2009, 27(4):223-232.