

Over-liming in the construction of the fertility of red yellow latosol with different phosphorus sources

0

ABSTRACT—The availability of P in Cerrado soils is a limiting factor for the satisfactory development of crops. It is known that the efficiency of phosphate fertilization is low and depends on several factors, such as solubility of fertilizers, soil texture, and soil acidity. Thus, the objective of this work was to evaluate the availability of phosphorus as a function of different phosphate sources under the influence of pH, as well as to determine its fertilization efficiency in a red latosol. The experiment was carried out in a greenhouse at Tocantins federal university, in DIC, with five replications. A red-yellow Latosol was used and three sources of phosphate fertilization were evaluated: Mono-ammonic phosphate - MAP, single superphosphate - SS and Natural phosphate - FN, plus one treatment without fertilization, in five evaluation periods (0, 7, 14, 21 and 28 days). Therefore, the pH in CaCl₂ of the soil stabilized between (5.5 – 6.5) in approximately 83 days of incubation with the acidity concealer. The SS and MAP phosphate fertilizers promoted the highest available P contents, with 6.32 and 6.23 mg.d m⁻³, respectively. The use of FN showed low P levels during the evaluated incubation period, mainly due to its lower solubility.

Keywords—fertilization efficiency, adsorption, soil incubation

INTRODUCTION

The Cerrado is the second largest Brazilian biome, covering about 25% of the total area of Brazil [1]. Through the intensive use of knowledge and technology, the agricultural frontier is expanding and, in this scenery, the Cerrado gains prominence in the conjuncture of the national agribusiness, especially the MATOPIBA region, the most recent Brazilian agricultural frontier [2]. Approximately 80 million hectares (40% of the total Cerrado area) have been cultivated with different types of land use, such as cultivated pastures and agricultural crops, in which soybean presents itself as the main culture responsible for the expansion of agribusiness in this region [3].

According to the last survey of Conab (2018) [2], soybean production in the Cerrado represents 78% of the amount produced in Brazil. The region comprised of Maranhão, Tocantins, Piauí, and Bahia (MATOPIBA) is responsible for 12% of Brazilian production, with the expected increase in the coming years.

The soil classes with the highest representatives of the region are the Latosols, Neossols, Argissols, Plinthosols, lytic soil and Cambisols [4]. Of these, the Latosols class occupies the largest territorial extension reaching 46% of the total Cerrado area, with great potential for agriculture [5, 6].

The Latosols comprise soils that present an advanced stage of tempering, consisting of mineral material, with a horizon diagnosis of the Latosol B immediately below any of the types of horizons, except the hystic. Generally, acidic soils, low clay content, low organic matter content, present colloidal material with low CTC and low natural fertility, emphasizing phosphorus (P) and potassium (K) [4].

The amount of available nutrients found in the soils in the form available to plants is one of the main

Comment [D1]: State names of authors and other relevant information.

Comment [D2]: Arrange in alphabetical order.

Comment [D3]: State parts of the region.

study objectives when it comes to evaluating soil fertility. However, the construction of soil fertility is not easy, because there is a great complexity of the processes that involve the dynamics of nutrients in the soil-solution-plant system [7].

Among the factors influencing soil fertility, acidity is a limiting factor in plant growth. Whenever the pH is low (acidity is high), one or more harmful effects may affect crop growth. For practical purposes, the pH considered appropriate for the majority of plants cultivated in Brazil is in the range between 6.0 and 6.5. Therefore, as the mineralogy of Brazilian soils are generally acidic, the use of liming stands out, which in turn allows the majority of essential nutrients to be readily available [8].

Liming and phosphate fertilization are practices used in the management of fertility, where the yield of crops is raised, especially in the regions where acidic and deficient soils in phosphorus are excelled^[9]. The correction made with limestone elevates the contents of Ca and Mg, decreasing or even eliminating the exchangeable Al, and with this practice increases the negative loads in these soils. These chemical alterations that occur in the soil may, however, influence some physical properties of the soil, by altering the electrochemical behavior of the colloids [10].

Phosphorus is one of the macro-nutrients less demanded by plants, but its low availability in the soil limits production, mainly in annual crops. More than 90% of the soil analyses in Brazil show a deficiency of this nutrient, with emphasis on the Cerrado soils, where the contents of this element are even smaller, reaching the worrying values of 1 mg dm^{-3} [11]. The P dynamics in Cerrado soils is quite complex due to its chemical characteristic, favoring great interaction with the soil, such as P adsorption to Fe and Al Oxy-hydroxides, which are predominant in the Latosols, which is the most representative class of soil in tropic regions [12]. This process can reduce the efficiency of phosphate fertilization by 70% and thus the use of high fertilizer dosages [5].

Currently, there are numerous fertilizers available on the market, which have differentiated solubility, as well as agronomic efficiency [13]. In addition, the efficient use of fertilizer can, in addition to providing greater productivity, reduce production costs and reflect on a positive margin at the end of the harvest. Therefore, different studies are being developed to improve the quality and efficiency of fertilizers, in order to decrease losses in the field [14].

According to Sousa & Lobato (2004) [15] the quality of the fertilizer used, soil type, an epoch of application, a form of application, uniformity of fertilizer application, are factors that when associated with soil moisture and management of the field interfere in fertilization efficiency.

Therefore, knowledge of phosphorus availability in Cerrado soils is of great importance for the measurement of fertility, seeking the best sources for the phosphate fertilization required to supply the needs of the crops, taking into account the interaction of nutrients with the soil. Therefore, the objective was to evaluate the availability of phosphorus as a function of different phosphate sources under pH

influences, as well as to determine its efficiency in a red-yellow latosol.

MATERIAL AND METHODS

The experiment was conducted in a greenhouse at the University Federal of Tocantins (UFT), campus Gurupi, located in the southern region of the state of Tocantins, with the coordinates: 11° 44' 44" S and 49° 03' 04" W. The soil used for the conduction of this assay was a red-yellow Latosol^[4], which was collected in the depth of 0 – 20 cm in the experimental farm of the UFT. The physical and chemical attributes were determined according to [16] in the Soil Laboratory - UFT (table 1).

Table 1. Chemical and Physical (0 – 20 cm) initial attributes of the red-yellow latosol. Gurupi – TO, 2018.

Ca ²⁺	Mg ²⁺	Al ³⁺	H+Al	CTC(T)	SB	CTC(t)	K	P		
	cmol _c dm ⁻³mg dm ⁻³			
0,66	0,43	0,12	2,78	3,92	1,14	1,26	18	1,04		
V	M	Organic matter.	pH CaCl ₂	Sand	Silt	Clay	Cu	Zn	Fe	Mn
	(%).....	Texture (%).....		mg dm ⁻³			
29,1	9,52	2,57	4,73	49,02	13,6	37,39	0,4	0,54	42,31	5,72

A completely randomized experimental design (IHD) was used, in a 3 x 5 + 1 scheme, where three sources of phosphate fertilization were evaluated: Mono-ammonium-phosphate - MAP (N - 11%, P₂O₅- 45%), single superphosphate — SS (P₂O₅ - 18%, Ca — 18%, S — 11%) and natural phosphate - FN (P₂O₅ - 7.5%, CaO — 4%, MgO — 0.5%, Si — 4%), with five replications and in five evaluation periods (0, 7, 14, 21 and 28 days), in addition to a treatment without fertilization (control). The dosage of P₂O₅ and FTE BR12 (9% Zn — 1.8% B — 0.8% Cu — 2% Mn — 3.5% Fe — 0.1% Mo) were fixed in 120 kg ha⁻¹ e 100 kg ha⁻¹ for all sources, respectively.

The experiment was carried out in plastic bags with 1 kg of soil previously sifted into a 2 mm sieve. Subsequently, dolomitic limestone (30% CaO, 18% — MgO — PRNT = 97.55%) was applied, and gypsum (30.70% CaO and 19.40% S). The lime and gypsum dosages used were 6 t ha⁻¹ and 1.8 t ha⁻¹, respectively, according to calibrated dosage in work performed by [17]. The soil was incubated for 55 days until the complete reaction of limestone and plaster, during this period the moisture was maintained at optimum levels, through the addition of distilled water whenever necessary. Subsequently, phosphate fertilization (MAP, SS, and FN) was added in each plot and soon after a new incubation cycle was performed for another 30 days.

The evaluated characteristics were the pH values (CaCl₂) and the P content in the soil. Soil sampling for the evaluations was performed before fertilization (0 days), at 7, 21 and 28 days after fertilization (DAA), which represents the days 55, 62, 69, 76 and 83 days after the beginning of the experiment. The soil was forwarded to the soil laboratory — LABSOIL of the UFT and prepared as TFSA for the chemical analysis of P (Mehlich⁻¹) and also the pH values (CaCl₂). The P and pH (CaCl₂) analyses were performed according to the methodology proposed by Embrapa (1997)^[16].

Comment [D4]: MATERIALS

Comment [D5]: Use capital T

Comment [D6]: Not clear. Are you referring to phosphate rock?

Comment [D7]: Passed through a 2 mm sieve.

Comment [D8]: Delete.

Comment [D9]: Total P?

The data obtained were subjected to analysis of variance and tests of the mean (Tukey), using the T-test and adopting 1 and 5% probability. Data analysis was determined using the statistical software Sisvar[®] version 6.5 [18] and the graphs were plotted using the software SigmaPlot[®] version 10 [19].

Comment [D10]: Give reference.

Comment [D11]: Reference?

The fertilization efficiency with the sources was calculated based on Eq. 1.

$$\text{Eq.1} \quad \text{Efic. (\%)} = \frac{(\text{P initial} - \text{P added})}{\text{P sample}} * 100 \quad \text{Eq. 1}$$

Where: P_{sample} P content available after treatments (mg/dm⁻³),
P_{initial} P content available in the initial condition (mg/dm⁻³) and
P_{added} P dosage of applied p (26.20 mg/dm⁻³).

RESULTS AND DISCUSSION

The analysis of variance and significance level, as well as the general mean and coefficient of variation for the characteristics P content and pH values, is found in table 2. For P contents, there was a significant effect ($p \leq 0.01$) for sources of variation, sources, epoch and interaction, and between source and epoch. The same result was observed for the pH values, except for the source. There is a low coefficient of variation for P and pH values in this study, which, according to Silva et al., (2010) [20], demonstrates homogeneous results for both traits.

Comment [D12]: 1. State your findings. 2. Give reasons for the findings. 3. Support the reasons with references.

Comment [D13]:

Comment [D14]: Presented

Comment [D15]: Capital T.

Comment [D16]: Italics.

Table 2. Analysis of variance, level of significance, general mean and coefficient of variation related to phosphorus content (p), and PH as a function of phosphate fertilizer sources (MAP, SS and FN, and 120 kg there⁻¹ of P₂O₅), Gurupi-TO, 2018.

Features	Source of variation				General average	C.V. (%)
	Sources	Epoch	Source x Epoch	Waste		
	Degree of freedom					
	3	4	12	80		
P	162,54**	62,7**	10,88**	9,82	4,07	8,6
Ph	0,009 ^{ns}	2,92**	0,03**	0,007	6,75	1,25

C.V.: Coefficient of variation. ^{*}: Significant at level 1% probability ($P \leq 0.01$); ^{*}: Significant at 5% probability level ($0.01 \leq p < 0.05$); ^{ns}: Non-significant ($p \geq 0.05$) by F test.

The Soil PH (CaCl₂), regardless of the source of P applied, obtained a quadratic response due to the days of evaluation after the liming with 6 t ha⁻¹ (Figure 1). The phosphate sources did not influence the soil pH, because the values in these treatments are equal to the values found for the treatment without application of P. The pH (CaCl₂) of the soil increased on average up to 72 days after the liming (6 t ha⁻¹), and thus obtained an increase of 29.8% in relation to the beginning of the experiment.

Comment [D17]: Small letter.

Comment [D18]: pH

Soil pH is of fundamental importance because it determines the availability of nutrients contained in the soil or added to it and also the assimilation of nutrients by the plants. Therefore, liming is a decisive factor in the efficiency of fertilization, since most Brazilian soils have a considerable degree of acidity [21]. The solubility of phosphate fertilizers is also influenced by soil pH, that is, the degree of acidity determines

the flow of P release to the soil solution. For acidulated phosphate fertilizers, such as MAP and SS, the pH range close to neutrality and considered ideal for plants (5.5 – 6.5) is that they present maximum solubility efficiency of P. Already in acidic condition, the nutrient available by these sources are quickly adsorbed by the colloids of the soil, and may not make them available to the plants. Low solubility fertilizers, such as reactive natural phosphates, require the action of H⁺ in the soil so that their orthophosphates are released to the solution. Therefore, for the natural phosphates, the increase of the pH to close to neutrality decreases its power of solubilization of P for the solution of the soil [22].




Figure. 1: PH values (CaCl) in the different sources of phosphate fertilization (MAP, SS, and FN) as a function of incubation time.

The treatments with the highest P content available in the soil were SS and MAP, with 6.32 and 6.23 mg dm⁻³ of P, respectively, which represents a superiority of 68.35 and 67.89% in relation to the treatment with FN (2.0 mg.dm⁻³). Moreover, the SS and MAP treatments when compared to the treatment without fertilization presented superiority of 72.62 and 72.23%. In relation to the FN, this presented P content available 13.5% higher than the treatment without the addition of P. Regarding P levels in relation to different epochs, the MAP treatment responded significantly. At 7 days after fertilization, P levels were higher than 85.71% in relation to 0 days. However, the nutrient release peak occurred between 14 and 21 days. After 28 days the contents showed a decrease being close to the values recorded on the 7th day.

For the SS, the epoch that presented the highest P contents was at 14, 21 and 28 days, statically equal

to each other, these times resulted in an increase of 86.83%, 86.73%, and 86.65% higher when compared to the treatment represented for day zero. Similarly, to the SS, the treatment represented by the FN had a maximum release of P at 14, 21 and 28 days, equal statically, with results 65.20, 64.28 and 63.60% higher than the control, represented by the time zero. It is also worth noting that at 7 days, the P content in the soil did not change, being statistically equal to the control treatment.




Figure. 2: For the phosphorus contents, there was a significant difference between the fertilization sources and the evaluation time.

However, the treatment without fertilization (control), also had positive results at 14, 21 and 28 days, with results 68.28, 71.87, and 65.05% higher when compared to the time 0 days. In relation to the control, there is an increase in P levels as the incubation time increases, the highest levels were observed at 14, 21 and 28 days. There is a small reduction in P levels at 28 days, but this reduction was not statistically significant.

According to Monteiro (2008) [23], the best result presented by the SS in this work may have been due to its high solubility in water and for being a relatively fast action fertilizer. For Reis (2009) [24], this does not happen with FN because it presents low solubility of P in the soil as a function of epoch.

In this study, it was observed that the MAP had a faster phosphorus solubilization behavior with 14 days and the nutrient release peak, followed by a decline in the next 21 and 28 days periods. The SS obtained a higher release with 14 days and maintained the peak release in the period from 21 to 28 days.

Therefore, for the construction of P fertility in this study for the average level, it is necessary to 120 kg

of P_2O_5 [25]. For the efficiency of P (Figure 3), there was also a significant difference between the fertilizer sources and the evaluation time.




Figure. 3: The efficiency of different sources of phosphate fertilizers as a function of sources and epochs.

The treatments that had higher fertilization efficiencies were SS and MAP, with 28.10 and 27.67%, respectively. These treatments had results 75.18 and 74.67% higher compared with the treatment of absence of P, which presented 7.05% efficiency. In acidic soils, FN is more efficient in solubilizing P, and with this, it presented lower results when compared to other fertilization, being slightly higher than the treatment that had an absence of P, but without statistical difference.

When it comes to efficiency in relation to different eras, the evaluation performed at 7 days after fertilization obtained an efficiency of the treatments MAP, SS, FN that corresponded to 26.73, 25.93 and 3.51%, respectively, being better to the treatment with absence of fertilization, which also presented a slight increase in P content over time. The increase in P availability in the treatment with the absence of fertilization probably occurred due to the neutralization of soil acidity by means of liming, promoting the best solubilization of the nutrient as pH reached the ideal levels [13].

At 14 days, the MAP, SS, and FN had 30.15, 28.97 and 10.01% efficiency, respectively, being relevant when compared to the treatment with an absence of fertilization that obtained 8.36% efficiency. With 21 days, the MAP, SS, and FN obtained 28.83, 28.78 and 9.76% efficiency, respectively, this efficiency is

considered better when compared to the treatment with an absence of fertilization that had an efficiency of 9.40%.

At the end of the incubation period, with 28 days the MAP treatment showed a decline and efficiency when compared to the previous times (24.83%). According to Yamada & Abdalla (2004) [13], phosphate fertilizers with higher efficiency solubility are monocalcium phosphates (SS) and monoammonium phosphate (MAP), but in this work, the MAP presented results of the release of unbelieve P as a function of epochs, which did not happen with SS. The SS and FN in the same period maintained the level of efficiency in the release of P for soil solution.

CONCLUSIONS

The MAP, SS and FN sources promoted an increase in phosphorus content in the soil. The best fertilization efficiencies were observed with the use of MAP and SS.

The applied dosage (120 kg ha⁻¹ of P₂O₅) increased the P levels at the average level of soil availability. The application of 6 t ha⁻¹ of dolomitic limestone promoted an increase in pH up to 71 days after incubation.

REFERENCES

- [1] Durigan, G. and Ratter, J. A. (2016). The need for a consistent fire policy for Cerrado conservation. *Journal of Applied Ecology*. 53(1):11–15. Available from: <https://doi.org/10.1111/1365-2664.12559>.
- [2] Conab, National Supply Company. Grain crop follow-up - Second November survey; (2018). Available from: <https://www.conab.gov.br/info-agro/safras/graos/boletim-da-safra-de-graos>.
- [3] Guareschi FR, Pereira MG, Perin A. (2012). Deposition of plant residues, light organic matter, carbon and nitrogen stocks and remaining phosphorus under different management systems in the Cerrado Goiano. *Brazilian Journal of Soil Science*. 36(3). Available from: <http://dx.doi.org/10.1590/S0100-06832012000300021>.
- [4] Embrapa; (2018). Embrapa solos. In: and others, editor. *Brazilian soil classification system*. p. 523–523. Brasileira de Pesquisa Agropecuria.
- [5] Coelho MR, Fontana A, Santos HG, Perez DV; (2014). Brazilian Society of Soil Science. In: *Soil and agricultural sustainability in Brazil, a pedological approach*. Viosa - MG. p. 30–30.
- [6] Leite RDC, Silva JSC, Freitas GA, Casali ME, Silva RR. (2017). Phosphate fertilization in soybeans during three consecutive harvests in the new Brazilian agricultural frontier. *Scientia Agraria*. 18(4):28–35. v.18, n.4. Available from: <http://dx.doi.org/10.5380/rsa.v18i4.50310>.

Comment [D19]: As corrected. Please effect the correction in all the references.

- [7] Cantarutti RB, Barros NF, Martinez HEP, Novais RF. (2007). Evaluation of soil fertility and recommendation of fertilizers. Brazilian Society of Soil Science. p. 769–850.
- [8] Lopes AS; (1995). International Soil Fertility Manual.
- [9] Ernani PR, Nascimento AL, Campos ML. (2000). Influence of phosphorus and limestone combination on corn yield. Brazilian Journal of Soil Science. 24(3). Available from: <http://dx.doi.org/10.1590/S0100-06832000000300007>.
- [10] Haynes RJ, Naidu R. (1998). Influence of lime, fertilizer and manure applications on soil organic matter content and soil physical conditions: A review Nutrient Cycling Agroecosystems. 51:123–137. Available from: <http://dx.doi.org/10.1023/A:1009738307837>.
- [11] Faquin V, Rodas CL (2015). Nutrient functions in plants: study guide. Lavras - MG: Federal University of Lavras. p.135.
- [12] Novais RF, Alvarez VH, Barros NF, Fontes RL, Cantarutti RB, et al.; (2007). Brazilian Society of Soil Science. In: and others, editor. Soil Fertility Manual. p. 117–117.
- [13] Yamada T, Abdalla SRS; (2004). Brazilian Association for Potassa and Fosfique research. In: and others, editor. Phosphorus in Brazilian Agriculture. p. 99–99.
- [14] Anda, Embrapa; (2005). National Association for the diffusion of fertilizers. In: and others, editor. Statistical yearbook: Fertilizer sector. p. 165–165.
- [15] Sousa DMG, Lobato E, Embrapa; (2004). Cerrado: soil correction and fertilization. p. 416–416.
- [16] Embrapa, Embrapa solos; (1997). Brazilian Agricultural Research Company. In: and others, editor. Brazilian soil classification system. p. 49–49.
- [17] Machado AF, Federal University of Tocantins. Dissolution, limestone doses, methods and nutrient availability in five Tocantins soils. Gurupi - TO; (2017). Tese Mestrado.
- [18] Ferreira DF, Federal University of Lavras. (2011). Sisvar: a computer statistical analysis system. Cincia e Agrotecnologia. 35(6):1039–1039.
- [19] Systat S; (2018). User manual of the Sigma Plot 10. San Jose - USA: Systat.
- [20] Silva SA, Lima JSS, Cândido A. (2010). Spatial variability of chemical attributes of a red-yellow hymic latosol grown with coffee. Brazilian Journal of Soil Science. 34(1). Available from: <http://dx.doi.org/10.1590/S0100-06832010000100002>.
- [21] Malavolta E; (2006). Agronômica Ceres. In: Plant mineral nutrition manual. So Paulo -SP; Agronmica Ceres. p. 59–59.

- [22] Prochnow LI, Escola superior de Agricultura Luiz de Queiroz; (2001). Fertilizer Phosphates: Some beliefs and some scientific facts.
- [23] Monteiro MF, Universidad Federal de Bahía; (2008). Life cycle assessment of the superphosphate simple fertilizer. Salvador-BA.
- [24] Reis THP. Evaluation of the life cycle of simple fertilizer. Lavras - MG; (2009).
- [25] Alvarez V, Novaes R, Barros N, Cantarutti R, Lopes A, et al.; (1999). Recommendation for the use of correctives and fertilizers in Minas Gerais. vol. 359.

UNDER PEER REVIEW