

Review Article

Future Therapeutic Strategies in the Glaucoma Management

ABSTRACT

Glaucoma is a group of eye diseases in which progressive damage to the ocular nerves may cause retinal ganglion cell (RGC) death. ~~Glaucoma is still the main reason for poor eyesight and the inability to see things around the world.~~ Intraocular pressure (IOP) is considering to be the main identified cause of danger so far, and lowering intraocular pressure is the only recognized technique for inhibiting disease progression. Furthermore, blood vessels and genetic components of glaucoma are considered additional risk factors. In order to realize the potential progress of glaucoma treatment, new treatment strategies and ambitious goals are constantly being developing. These treatments will provide specific tissue goals to reduce IOP and ensure neuroprotective effects on RGCs. Consequently, physicians can shortly have an expanded range of medical choices to choose from, coupled with therapies that are more successful. Therefore, this study has reviewed the recent studies that were conducted on cellular mechanisms of glaucoma treatment.

KEYWORDS: Glaucoma; IOP; ~~mechanism of action~~

ORCID: 0000-0002-8851-5783

Introduction

Glaucoma refers to a series of eye diseases that can cause progressive damage to the optic nerve and may cause RGC ~~deaths~~. According to a 2010 report by the World Health Organization, glaucoma accounts for 2% of visual disability and 8% of total blindness. Due to the growth of population, the total number of glaucoma patients is expecting to be increased^[1]. The classification of glaucoma depends on the condition of the drainage route. The

Formatted: Strikethrough

Comment [kk1]: Replace with:
Worldwide Glaucoma is a leading cause of avoidable blindness.

Comment [kk2]: Replace with:
Pharmacokinetics

Formatted: Strikethrough

Formatted: Strikethrough

Comment [kk3]: death (it is not plural)

Formatted: Strikethrough

Formatted: Strikethrough

Comment [kk4]: Please add the Reference:
Global estimates of visual impairment: 2010, S.P.Mariotti, D. Pascolini, Br J Ophthalmol. 2012 May;96(5):614-8.

drainage pathway of open-angle glaucoma looks normal, while the drainage pathway of closed-angle glaucoma is blocked. Glaucoma was moreover categorized to the primary and secondary glaucoma. The most common subtype of glaucoma is primary open-angle glaucoma. Therefore, glaucoma is associated with increased intraocular pressure, and currently ~~the drugs that lower intraocular pressure are still the only clinically proven treatment for~~ glaucoma[2]. The focus of the current review is to briefly summarize and discuss the new goals for reducing IOP.

Comment [kk5]: (Erase article)

Formatted: Strikethrough

Formatted: Strikethrough

Current Treatment Strategies for Glaucoma

Glaucoma is a disease that is not well understood, but the main goal of treatment is to reduce intraocular pressure [2][3]. A 40% reduction in intraocular pressure will reduce the progress of the ocular loss by about half. The first drug for glaucoma was released in 1875, and now there are many IOP eye drops for reducing glaucoma. Compromise options for ophthalmic classification: beta blockers, prostaglandin analogues, carbonic anhydrase inhibitors, and α 2-sympathomimetic receptor agonists [4], as shown in table 1.

Future Therapy Plans for Glaucoma

The pathogenesis of glaucoma includes many mechanisms. However, no one seems to fully describe the disease, and different etiologies have carried out key trials in the progress of new anti-glaucoma strategies. In this part, treatment plans can emphasize on IOP lowering approach, therefore, an effort has been done to discover novel, curative goal to lower synthesis of aqueous humor (AqH) (Fig.1 and 2) or to promote the uveoscleral outflow into the anterior chamber angle using assorted subcellular pathways from those already existing[5].

Nitric Oxide

The endogenous signaling molecule is called nitric oxide (NO), which is a new goal to reduce IOP [6]. In many tissues of the eye, both the anterior and posterior endogenously produce NO, and it has a stimulator of the intracellular receptor soluble guanylate cyclase (GC). The latest evidence links NO to GC through the IOP director and the enzyme cyclic guanosine monophosphate (cGMP) pathway in glaucoma's retinal pathophysiology. There are cumulative signs that the direct guide to intraocular pressure is NO, and the failure of the NO-sensitive soluble GC-1 (NO-GC1) pathway is associated with glaucoma. Ciliary body (CB), the trabecular meshwork (TM), and the Schlemm's canal (SC) are tissues able to generate NO in normal eyes [7]. In some cases, the impact of NO on IOP is related to the activities of the downstream second messenger cGMP, mainly in the usual outflow route. Taking of NO-generating compounds to encourage the NO-GC 1-cGMP pathway can reduce IOP through TM relaxation, TM

volume changes, and increased SC cell permeability [8]–[11]. "Many studies conducted in rabbits have shown that NO donors and cGMP analogues can significantly reduce IOP and lead to increase outflow facilities [12]. In a dose-dependent manner, the intravitreal or intraluminal injection of a cGMP analog named 8-Br-cGMP may increase the AqH outflow facility [13]. The 8-Br-cGMP is an activator of cGMP enzyme, it can easily penetrate cells, and its hydrolysis by phosphodiesterases is less sensitive than that of native cGMP [14]. When used in low doses, 8-Br-cGMP will show a decrease in AqH flow, but the outflow facility will not be affected. On the other hand, 8-Br-cGMP displays an improvement in outflow facility at high doses [13]. One possible mechanism for promoting the inhibitory effect of NO on AqH synthesis includes the inhibition of Na, K-ATPase (NKA) pumps. In the ocular tissues, NKA is the main active transporter involved in the formation of ion gradients that control the formation of AqH[15]. Ouabain lowers the secretion of AqH by about 62% by inhibiting the NKA pump (17). In addition, the use of sodium nitroprusside (SNP) on bovine and porcine eyes cut the second messenger cGMP and PKG, thereby inhibiting NKA pump thereby inhibiting the secretion of AqH, and thus reducing IOP(16).

Phosphodiesterases-5 Inhibitor

A family of enzymes that control intracellular levels of the second messenger's cAMP and cGMP is called phosphodiesterases. The first phosphodiesterase inhibitor was recognized in 1970 and up to the present time, 12 dissimilar isoenzymes were recognized [21]. Phosphodiesterase (PDE) is found in every cell in the body. However, the distribution of isoenzymes differs between the tissues. There are 11 types of PDE, some of which highly selective for cGMP such as PDE5, 6 and 9, other types have coupled specificity for both cGMP and cAMP such as PDE1,2 and 11 and others have selective cAMP but sensitive cGMP such as PDE3 and 10. The production of cGMP induces PDE5 activity by allosteric integration of cGMP into the GAF regulatory tandem domain. The enzyme catalytic rate and affinity are increased due to cGMP attachment to the PDE5 GAF domain. At the same time, cGMP-dependent protein kinase (cGK I) phosphorylates PDE5 in its N-terminal region. After phosphorylation, the most likely cause of prolonged PDE5 activation is increased sympathy of cGMP. The regulation of the signaling sensitivity of NO/cGMP in platelets is controlled by a feedback mechanism, and manifests as a temporary form of cGMP response (rapid increase in cGMP and hydrolysis) [22]. Furthermore, this mechanism is also explaining the desensitization induction by NO - the decreased cGMP quantity next to second stimulation by NO- (26). The efficiency of PDE5 inhibitors depends on the manufacturing capacity of cGMP, and

the fact that PDE5 closely interacts with NO/cGMP signaling is well accepted. The effectiveness of PDE5 inhibitors depends on the ability of cGMP to construct. This is because PDE5 enzymes are closely related to NO/cGMP signaling. Nitric oxide synthase (NOS) inhibition retract the synthesis of NO and subsequent cGMP production, thus inhibiting of the vascular dilation effect through PDE5 inhibition. Inhibition of PDE expected to be less applicable in locations related with diminished NOS action, such as endothelial function impairment. On the other hand, sildenafil-PDE5 inhibitors-can effectively enhance the effects of NO-producing compounds. Therefore, patients taking organic nitrates and sildenafil have severely reduced blood pressure (24). In Rabbits, inhibition of PDE-5 produce important IOP reduction, such effect endorsed by topical administration of sildenafil ophthalmic solution (0.3%) in both normotensive and corticosteroid induced hypertensive rabbit's eyes[27], this study suggests that sildenafil could be considered as an inhibitory modulator of AH production by decreasing intracellular Ca^{+2} [28]. In addition to the above, sildenafil can also increase AH outflow and uveoscleral outflow [29]. Other suggested mechanism by which sildenafil can inhibit AH secretion by blocking NKA pump in NPE as a result of cGMP elevation [30]. This mechanism is consistent with Ellis and colleagues who reported that elevated cGMP inhibits NKA in the choroid plexus and CB [31]. Therefore, this study suggests that increasing cGMP as a result of using sildenafil may reduce IOP.

NA, K-ATPASE enzyme

About 60 years ago, Skou ~~was discovered as an NKA pump, which can be used as a molecular tool to transport Na + and K + through cell membranes.~~ NKA is a part of the P-type ATPase family. NKA is a protein complex molecule widely distributed in cell membranes, with two non-covalently linked subunits α and β [32]. Subunit α which contains ATP and other ligand attachment sites is designed as a catalytic subunit. The role of β subunit is vital for the membrane focusing and full participation of the NKA. Four protein isoforms of NKA have been recognized. In 1987, Michael Marks and Nicholas Seeds were found to have various isoforms. Digoxin and ouabain (plant-derived cardiac steroids) are known to be inhibitors of NKA pumps (34). Over 200 years, digoxin has used widely for congestive heart failure management. However, it was not until the detection of NKA pumps in the 1950s that the exact mechanism by which digitalis soared to inhibit NKA was documented [35]. The pigmented cells of the ciliary epithelium express $\alpha 1\beta 1$ isoform of, whereas the nonpigmented cells of the ciliary epithelium show $\alpha 2\beta 3$ isoform of NKA. With limited rare local and systemic toxicity, the selective inhibition of $\alpha 2$ isoform can effectively go

Comment [kk6]: (3 words to be deleted, they change the meaning)

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

lower . Earlier trials using recombinant human isoforms of $\alpha_1\beta_1$, $\alpha_2\beta_1$, and $\alpha_3\beta_1$ displayed digoxin with selective α_2 activity due to its trisdigitoxose moiety[36]. This fact led to a likelihood that alteration of the trisdigitoxose might elevate the selectivity toward α_2 . Several derivatives display greater selectivity for α_2 over α_1 , nearly 8-fold. In rabbits, the effect of topical administration of cardiac glycosides on IOP has been considered by inhibiting or reversing the ability to use A3 adenosine receptor selective agonists to induce a critical increase in IOP [37]. Digoxin derivatives with relatively α_2 -selective activity shows more IOP lowering effect when compared to that of digoxin, digoxigenin, or ouabain. Therefore, due to the main role of α_2 in the production of AqH, α_2 -selective derivatives of digoxin may become a new therapy for regulating IOP [36]. Furthermore, a study done by Waleed et al. endorsed that topical application of digoxin (0.00625%) ophthalmic solution reduces IOP significantly in both normotensive and betamethasone induced hypertensive rabbits eyes [17]. Other studies have shown that the administration of low and non-toxic concentrations of ouabain increases AqH outflow. The researchers studied other mechanisms of action of ouabain's effectiveness. Simon and colleagues demonstrated the ability of ouabain to reduce IOP (38) by attaching and preventing the NKA pump in the ciliary process of the eye[38]. NKA enhances the transfer of two K^+ from the outside space into the cell and the removal of three Na^+ from the cell, while hydrolyzing ATP to ADP and inorganic phosphate (Pi). The emergence of electrochemical gradients represents the main energy source used by transportation equipment to drive ions through AqH to form the required transepithelial transportation [39]. In addition, research shows that the combination of ouabain and NKA may control cytoskeletal proteins, thereby changing the morphology of cells [40].

Calcium channel blocker

Calcium is a vital intracellular messenger [41] and influx Ca^{2+} could have many effects on the dynamics of AqH, comprising ciliary perfusion as well as hydrostatic and osmotic component. Calcium channel blockers (CCBs) are drugs often used to treat hypertension and coronary artery disease[42] and to reduce vascular tone by reducing Ca^{2+} influx leads to vasodilation and raising in the local blood flow in many tissues constituting the head of ocular nerve. It can also show another beneficial effect of CCB by inhibiting the synthesis of extracellular matrix collagen [43]. ~~Trabecular meshwork cell relaxation~~ increases the outflow facility of AqH by blocking the L-type channel also produced by CCB. Perfusion experiment work with verapamil administration in the incised human eye showed a dose-related increase in the outflow facility(48). Most studies conducted in laboratory animals and humans

Comment [kk7]: Although the systemic use of Calcium Channel Blockers for treatment of systemic hypertension is possibly associated with increased risk of primary open – angle glaucoma, topical use enhances Trabecular meshwork cell relaxation, which
Please add the Reference:
Khawaja, Anthony P. Calcium Channel Blockers and Risk of Primary Open-angle Glaucoma. Journal of Glaucoma, March 2019 - Volume 28 - Issue 3 - p e49-e50.

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

believe that local CCB (eg, nimodipine, verapamil, diltiazem, nifedipine, or flunarizine) reduces IOP. "Topical diltiazem can minimize intraocular pressure (approximately 5 mmHg) in normal rabbits [46]; (4 mmHg) in the betamethasone-induced ocular hypertensive rabbits [47]. Topical application of diltiazem can respectively reduce the intraocular pressure of normal, betamethasone induced intraocular hypertension and water load induced intraocular pressure (6 mmHg; 4 mmHg and 5 mmHg) [48], while normal cynomolgus monkey's IOP decreased the most (4 mmHg, [49]. Maximum intraocular pressure reduction after local application of nifedipine is (6 mmHg) in normal rabbits [46] and (5 mmHg) in cynomolgus monkeys. Netland and colleagues showed that verapamil infusion can reduce intraocular pressure in normal people and cynomolgus monkeys, and has a beneficial effect on the posterior bulb or optic nerve head circulation [50]. Other studies have shown that verapamil can reduce IOP in normal people by about 3 mmHg [51]; normal cynomolgus monkeys are 2.6 mmHg [49]; 49); and 6 mmHg in normal rabbits [46]. In 12 patients with intraocular hypertension, a single local application of verapamil 0.125% can encourage a reduction in IOP of 3 to 4 mm Hg for 10 hours, while a simple drop was detected in normal IOP volunteers (≈ 1.5 mm Hg) [52]. In patients with high intraocular pressure, topical application of 0.125% verapamil for 14 days can reduce IOP by about 7.0 ± 2.9 mm Hg [53]. Sunil and his colleagues showed that topical application of diltiazem (0.5%) and diltiazem (0.125%) in corticosteroid-induced glaucoma for 12 days can reduce intraocular pressure by about 8.0 and 9.6 mmHg, respectively [54]. Researcher Waleed and colleagues showed in a study that after one day of topical eye drops (0.5%) of nimodipine can reduce IOP by (9.09%) in normotensive and (19.29%) in betamethasone -induced hypertensive rabbits eyes. Also, the peak IOP decline achieved after 4 days of installation in both normotensive (16.16%) and hypertensive (22.86%) models [55]. Flunarizine with sodium channel blocking activity, L-type and T-type calcium channel blockers have good ocular bioavailability, and the maximum IOP reduction when applied topically is 2-5 mmHg ;10 mmHg; 4 mmHg [49] and 5 mmHg in normal rabbits (58), chymotrypsin- induced ocular hypertensive rabbits [57]; normal cynomolgus monkeys [59] and unilateral laser-induced ocular hypertensive cynomolgus monkeys [60] respectively. Additionally, due to its inhibition of calcium and sodium influx, flunarizine shows a protective effect on retinal injury that induced by intravitreal injection of N-methyl-D- aspartic acid (NMDA) [56]. Similarly, for patients with normal tension glaucoma (NTG), oral flunarizine also shows important significance for retrobulbar hemodynamics. In addition, flunarizine inhibits the contraction of human trabecular meshwork cells induced by ET-1. CCB interferes with gap junctions or cation transport in ciliary epithelial cells, resulting in reduced AqH synthesis [61].

Renin-angiotensin system

The circulating renin-angiotensin system (RAS) plays an important role in regulating blood pressure and fluid balance. The peptide angiotensin II (Ang II), the angiotensin-converting enzyme (ACE1) and the angiotensin receptor type1 (AT1) are chiefly guided by RAS. Vasoconstriction, stimulation of the sympathetic nervous system and stimulation of aldosterone production are the main effects of Ang II through which it controls the blood pressure [62]. Tissue AT1 receptor is the main target of Ang II. As stated in the new confirmation, it is believed that the increase in blood pressure in the elderly is not only regulated by circulating RAS, but that autocrine or paracrine of local RAS also plays a vital role in controlling blood pressure. Therefore, not only Ang II and ACE1 found in tissues are involved in controlling blood pressure, but also angiotensin (1-7) [63]. As a result from mentioned studies, the treatment of hypertensive patient must focus on desired actions that resulted from stimulation of Ang(1-7)- ACE2 Mas axis, as well as suppression of the normal AT1- Ang II- ACE1 axis(70). Recently, RAS in the eye tissue of non-pigmented and colored epithelial cells plays an important role in the formation and drainage of AqH. Ang II opens potassium channels and stimulates aldosterone production by triggering the Ca²⁺ signaling system. The volume of cell loss that accompanies the mentioned effects indicates that Ang II has secretagogue activity in NPEC [66]. Additionally, In addition, the activation of Na⁺/H⁺ exchange by Ang II will lead to an increase in Na⁺ cytoplasmic concentration. There is a link between the sodium and the pathological factors in epithelia of both renal tubules and CB, which give a reason of the concurrence of systemic hypertension and glaucoma [66]. Furthermore, RAS plays an essential role in the outflow of AqH from TM. Inducing cell proliferation in bovine and in vitro by stimulating collagen synthesis is the recommended role of Ang II [67].

~~Intracamerally administration of Ang II able to reduce the uveoscleral outflow[68]. The application of both natural and synthetic Ang II to human eyes and cat eyes can reduce IOP, with the same reduction effect observed in in vivo and in vitro studies. Iris artery vasoconstriction is the mechanism by which it reduces the IOP[68]. Human studies have shown that although the effect of lowering blood pressure is only shown in patients with arterial hypertension, ACE1 inhibitors or AT2 receptor blockers can reduce intraocular pressure in glaucoma and normal tension glaucoma patients In both glaucoma models (acute and chronic), perindopril has an ocular hypotensive effect on rabbits [69]. ACE1 inhibitors can diminish the AqH formation in the CB by reducing blood flow. ACE1 inhibitors promote the formation of prostaglandins by inhibiting the breakdown of bradykinin, which may result in decreased intraocular pressure by promoting the outflow of the uveal scleral area (80). The inhibiting effects on~~

Comment [kk8]: Furthermore, intracameral administration of Ang II is...

Formatted: Strikethrough

bradykinin breakdown by ACE inhibitors also stimulate prostanoids vasodilation effects and the NO pathway, as well as decrease the synthesis of the endothelin-1, a vasoconstrictive peptide that shown to display contraction activity in both porcine and in the human ophthalmic ciliary epithelium(78). Specifically, for glaucoma losartan has ability to decrease normal or elevated IOP in humans IOP [73]. Similarly, in animal models that induce an increase in IOP, olmesartan can reduce IOP (84). In addition to reducing the effect of IOP, the neuroprotective activity shown during the use of ARB in the glaucoma environmen. Candesartan has been displayed to reduce RGC loss in rats with increased intraocular pressure and normal glaucoma (85). Furthermore, losartan when given to the mouse eyes with raised IOP shows a neuroprotective activity on RGCs [79]. A study showed that after one day of infusion of telmisartan eye drops (1%), significantly reduced IOP(12%) in betamethasone- hypertensive rabbit eyes and reached a maximum IOP reduction (20.33%) after 7 days of infusion[80]. Captopril administration continuously to the rat with chronic model of glaucoma showed an important neuroprotection activity against RGC loss [77]. In addition to its role in balancing intraocular fluids, local RAS also plays a key role in the development of diabetic retinopathy. Recently, a prorenin receptor found in Mueller's retinal cells is involved in the development of neovascular pathology, and the inhibitory effect of these receptors has shown beneficial effects on diabetic retinopathy [81]. From the mentioned findings, we conclude the role of RAS in reducing IOP. New options for glaucoma treatment focus on increasing the activity of ACE2, Ang (1-7) and Mas receptors. What's more, RAS acting compounds exert a -beneficial role in neuroprotection activities.

Conclusion

This review focuses on novel treatment strategies and possible expected mechanisms to reduce IOP. Many new targets for glaucoma treatment submitted, but up to now the merely available compounds for glaucoma treatment focusing on lowering IOP, which are diminishing the disease progression rate. Accordingly, physicians may soon have an enlarged number of curative choices to select from, together with possibly potent treatments. Any change from treatment of the disease to new therapies must be given too much attention.

References

1. Tham Y-C, Li X, Wong TY, Quigley HA, Aung T, Cheng C-Y. Global prevalence of glaucoma and projections of glaucoma burden through 2040: a systematic review and meta-analysis.

Comment [kk9]: Taking everything into consideration, in the foreseeable future, ophthalmologists is expected to have an enlarged number of glaucoma treatment options and even curative.

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Formatted: Strikethrough

Comment [kk10]: Beyond doubt, too much attention must be paid to any replacement of classic treatments by new therapies.

Formatted: Strikethrough

Formatted: Strikethrough

- Ophthalmology. 2014;121(11):2081–90.
2. Chang EE, Goldberg JL. Glaucoma 2.0: neuroprotection, neuroregeneration, neuroenhancement. *Ophthalmology*. 2012;119(5):979–86.
 3. Germano RAS, Finzi S, Challa P, Susanna Junior R. Rho kinase inhibitors for glaucoma treatment-Review. *Arq Bras Oftalmol*. 2015;78(6):388–91.
 4. Wiederholt M, Thieme H, Stumpff F. The regulation of trabecular meshwork and ciliary muscle contractility. *Prog Retin Eye Res*. 2000;19(3):271–95.
 5. Beidoe G, Mousa SA. Current primary open-angle glaucoma treatments and future directions. *Clin Ophthalmol (Auckland, NZ)*. 2012;6:1699.
 6. Aliancy J, Stamer WD, Wirostko B. A review of nitric oxide for the treatment of glaucomatous disease. *Ophthalmol Ther*. 2017;6(2):221–32.
 7. Verkman AS. Role of aquaporin water channels in eye function. *Exp Eye Res*. 2003;76(2):137–43.
 8. Chang JYH, Stamer WD, Bertrand J, Read AT, Marando CM, Ethier CR, et al. Role of nitric oxide in murine conventional outflow physiology. *Am J Physiol Physiol*. 2015;309(4):C205–14.
 9. Nathanson JA. Direct application of a guanylate cyclase activator lowers intraocular pressure. *Eur J Pharmacol*. 1988;147(1):155–6.
 10. Dismuke WM, Sharif NA, Ellis DZ. Human trabecular meshwork cell volume decrease by NO-independent soluble guanylate cyclase activators YC-1 and BAY-58-2667 involves the BKCa ion channel. *Invest Ophthalmol Vis Sci*. 2009;50(7):3353–9.
 11. Heyne GW, Kiland JA, Kaufman PL, B'Ann TG. Effect of nitric oxide on anterior segment physiology in monkeys. *Invest Ophthalmol Vis Sci*. 2013;54(7):5103–10.
 12. Kotikoski H, Vapaatalo H, Oksala O. Nitric oxide and cyclic GMP enhance aqueous humor outflow facility in rabbits. *Curr Eye Res*. 2003;26(2):119–23.
 13. Kee C, Kaufman PL, Gabelt BT. Effect of 8-Br cGMP on aqueous humor dynamics in monkeys. *Invest Ophthalmol Vis Sci*. 1994;35(6):2769–73.
 14. Zimmerman AL, Yamanaka G, Eckstein F, Baylor DA, Stryer L. Interaction of hydrolysis-resistant analogs of cyclic GMP with the phosphodiesterase and light-sensitive channel of retinal rod outer segments. *Proc Natl Acad Sci*. 1985;82(24):8813–7.

15. Ghosh S, Freitag AC, Martin-Vasallo P, and Coca-Prados M. Cell Distrib Differ gene Expr three alpha subunit isoforms Na, K-ATPase Ocul ciliary Ep J Biol Chem. 1990;265:2935–40.
16. Shahidullah M, Wilson WS, Yap M, To C. Effects of ion transport and channel-blocking drugs on aqueous humor formation in isolated bovine eye. Invest Ophthalmol Vis Sci. 2003;44(3):1185–91.
17. Abdulsahib WK, Abood SJ. Effect of digoxin ophthalmic solutions on the intraocular pressure in rabbits.
18. Wareham LK, Buys ES, Sappington RM. The nitric oxide-guanylate cyclase pathway and glaucoma. Nitric Oxide. 2018;77:75–87.
19. Shahidullah M, Mandal A, Wei G, Delamere NA. Nitric Oxide Regulation of Na, K-ATPase Activity in Ocular Ciliary Epithelium Involves Src Family Kinase. J Cell Physiol. 2014;229(3):343–52.
20. Shahidullah M, Yap M, To C. Cyclic GMP, sodium nitroprusside and sodium azide reduce aqueous humour formation in the isolated arterially perfused pig eye. Br J Pharmacol. 2005;145(1):84–92.
21. Supuran CT, Mastrolorenzo A, Barbaro G, Scozzafava A. Phosphodiesterase 5 inhibitors-drug design and differentiation based on selectivity, pharmacokinetic and efficacy profiles. Curr Pharm Des. 2006;12(27):3459–65.
22. Mullershausen F, Friebe A, Feil R, Thompson WJ, Hofmann F, Koesling D. Direct activation of PDE5 by cGMP: long-term effects within NO/cGMP signaling. J Cell Biol. 2003;160(5):719–27.
23. Schlaich MP, Sobotka PA, Krum H, Lambert E, Esler MD. Renal sympathetic-nerve ablation for uncontrolled hypertension. N Engl J Med. 2009;361(9):932–4.
24. Groneberg D, König P, Wirth A, Offermanns S, Koesling D, Friebe A. CLINICAL PERSPECTIVE. Circulation. 2010;121(3):401–9.
25. Webb DJ, Muirhead GJ, Wulff M, Sutton JA, Levi R, Dinsmore WW. Sildenafil citrate potentiates the hypotensive effects of nitric oxide donor drugs in male patients with stable angina. J Am Coll Cardiol. 2000;36(1):25–31.
26. Mergia E, Stegbauer J. Role of Phosphodiesterase 5 and Cyclic GMP in Hypertension. Curr Hypertens Rep [Internet]. 2016 Apr;18(5):39. Available from: <https://pubmed.ncbi.nlm.nih.gov/27079836>

27. Abdulsahib WK, Al-Zubaidy A, Sahib HB, Kathem SH. Tolerable ocular hypotensive effect of topically applied sildenafil in ocular in normotensive and betamethasone-induced hypertensive rabbits. *Int J Pharm Sci Rev Res.* 2015;35(1).
28. McLaughlin CW, Peart D, Purves RD, Carré DA, Peterson-Yantorno K, Mitchell CH, et al. Timolol may inhibit aqueous humor secretion by cAMP-independent action on ciliary epithelial cells. *Am J Physiol Physiol.* 2001;281(3):C865–75.
29. Samuelsson-Almén M, Nilsson SFE, Mäepea O, Bill A. Effects of atrial natriuretic factor (ANF) on intraocular pressure and aqueous humor flow in the cynomolgus monkey. *Exp Eye Res.* 1991;53(2):253–60.
30. Kleineller A, Fambrough DM, Benos DJ, Civan MM. *The Eye's Aqueous Humor: From Secretion to Glaucoma.* Academic Press; 1997.
31. Ellis DZ, Nathanson JA, Sweadner KJ. Carbachol inhibits Na⁺-K⁺-ATPase activity in choroid plexus via stimulation of the NO/cGMP pathway. *Am J Physiol Physiol.* 2000;279(6):C1685–93.
32. Sweadner KJ. Isozymes of the Na⁺/K⁺-ATPase. *Biochim Biophys Acta (BBA)-Reviews Biomembr.* 1989;988(2):185–220.
33. Marks MJ, Seeds NW. A heterogeneous ouabain-ATPase interaction in mouse brain. *Life Sci.* 1978;23(27–28):2735–44.
34. Akera T, Brody TM. Inotropic action of digitalis and ion transport. *Life Sci.* 1976;18(2):135–41.
35. Cui X, Xie Z. Protein Interaction and Na/K-ATPase-Mediated Signal Transduction. *Molecules* [Internet]. 2017 Jun 14;22(6):990. Available from: <https://pubmed.ncbi.nlm.nih.gov/28613263>
36. Katz A, Tal DM, Heller D, Haviv H, Rabah B, Barkana Y, et al. Digoxin Derivatives with Enhanced Selectivity for the α2 Isoform of Na, K-ATPase EFFECTS ON INTRAOCULAR PRESSURE IN RABBITS. *J Biol Chem.* 2014;289(30):21153–62.
37. Socci RR, Chu E, Bayorh MA, Chu T-C. 4-aminopyridine transiently increases intraocular pressure in rabbits. *Pharmacology.* 2003;69(2):108–14.
38. Simon KA, Bonting SL, Hawkins NM. Studies on sodium-potassium-activated adenosine triphosphatase: II. Formation of aqueous humour. *Exp Eye Res.* 1962;1(3):253–61.
39. Delamere NA, Socci RR, King KL, Bhattacharjee P. The influence of 12 (R)-

- hydroxyeicosatetraenoic acid on ciliary epithelial sodium, potassium-adenosine triphosphatase activity and intraocular pressure in the rabbit. *Invest Ophthalmol Vis Sci.* 1991;32(9):2511–4.
40. Jung J, Yoon T, Choi EC, Lee K. Interaction of cofilin with triose-phosphate isomerase contributes glycolytic fuel for Na, K-ATPase via Rho-mediated signaling pathway. *J Biol Chem.* 2002;277(50):48931–7.
41. Kathem SH, Nauli SM. Mechanosensory function of Cav1. 2 calcium channel in renal primary cilia. *FASEB J.* 2016;30(1_supplement):1183–8.
42. Fadheel OQ, Al-Jawad FH, Abdulsahib WK, Ghazi HF. Effect of Felodipine against Pilocarpine induced Seizures in Rats. *Int J Pharm Sci Rev Res.* 2018;52:54–60.
43. Shayegan MR, Bolorian AA, Kianoush S. Comparative study of topical application of timolol and verapamil in patients with glaucoma within 6 months. *J Ocul Pharmacol Ther.* 2009;25(6):551–4.
44. Green K, Kim K. Papaverine and verapamil interaction with prostaglandin E2 and Δ 9-tetrahydrocannabinol in the eye. *Exp Eye Res.* 1977;24(2):207–12.
45. Podos SM. The effect of cation ionophores on intraocular pressure. *Invest Ophthalmol Vis Sci.* 1976;15(10):851–4.
46. Santafe J, Jesu M, Segarra J, Melena J. A long-lasting hypotensive effect of topical diltiazem on the intraocular pressure in conscious rabbits. *Naunyn Schmiedebergs Arch Pharmacol.* 1997;355(5):645–50.
47. Melena J, Santafé J, Segarra J. The effect of topical diltiazem on the intraocular pressure in betamethasone-induced ocular hypertensive rabbits. *J Pharmacol Exp Ther.* 1998;284(1):278–82.
48. Santafé J, de Ibarreta MJM, Segarra J, Melena J. The effect of topical diltiazem on ocular hypertension induced by water loading in rabbits. *Gen Pharmacol Vasc Syst.* 1999;32(2):201–5.
49. Siegner SW, Netland PA, Schroeder A, Erickson KA. Effect of calcium channel blockers alone and in combination with antiglaucoma medications on intraocular pressure in the primate eye. *J Glaucoma.* 2000;9(4):334–9.
50. Erickson KA, Schroeder A, Netland PA. Verapamil increases outflow facility in the human eye. *Exp Eye Res.* 1995;61(5):565–7.
51. Abreu MM, Kim YY, Shin DH, Netland PA. Topical verapamil and episcleral venous pressure.

- Ophthalmology. 1998;105(12):2251–5.
52. Netland PA, Grosskreutz CL, Feke GT, Hart LJ. Color Doppler ultrasound analysis of ocular circulation after topical calcium channel blocker. *Am J Ophthalmol*. 1995;119(6):694–700.
 53. Goyal JK, Khilnani G, Sharma DP, Singh J. The hypotensive effect of verapamil eye drops on ocular hypertension. *Indian J Ophthalmol*. 1989;37(4):176.
 54. Ganekal S, Dorairaj S, Jhanji V, Kudlu K. Effect of topical calcium channel blockers on intraocular pressure in steroid-induced glaucoma. *J Curr glaucoma Pract*. 2014;8(1):15.
 55. Abdulsahib WK, Fadhil OQ, Tizkam HH. Effect of topically applied nimodipine on the intraocular pressure on ocular normotensive and betamethasone-induced hypertensive eyes in rabbits. *Int J Res Pharm Sci*. 2019 Oct 1;10(4):2727–32.
 56. Osborne NN, Wood JPM, Cupido A, Melena J, Chidlow G. Topical flunarizine reduces IOP and protects the retina against ischemia-excitotoxicity. *Invest Ophthalmol Vis Sci*. 2002;43(5):1456–64.
 57. Maltese A, Bucolo C. Pharmacokinetic profile of topical flunarizine in rabbit eye and plasma. *J Ocul Pharmacol Ther*. 2003;19(2):171–9.
 58. Pauwels PJ, Leysen JE, Janssen PAJ. Ca⁺⁺ and Na⁺ channels involved in neuronal cell death. Protection by flunarizine. *Life Sci*. 1991;48(20):1881–93.
 59. Campana G, Bucolo C, Murari G, Spampinato S. Ocular Hypotensive Action of Topical Flunarizine in the Rabbit: Role of ζ 1 Recognition Sites. *J Pharmacol Exp Ther*. 2002;303(3):1086–94.
 60. Wang R-F, Gagliuso DJ, Podos SM. Effect of flunarizine, a calcium channel blocker, on intraocular pressure and aqueous humor dynamics in monkeys. *J Glaucoma*. 2008;17(1):73–8.
 61. Mito T, Delamere NA, Coca-Prados M. Calcium-dependent regulation of cation transport in cultured human nonpigmented ciliary epithelial cells. *Am J Physiol Physiol*. 1993;264(3):C519–26.
 62. Fyhrquist F, Metsärinne K, Tikkanen I. Role of angiotensin II in blood pressure regulation and in the pathophysiology of cardiovascular disorders. *J Hum Hypertens*. 1995;9:S19-24.
 63. Ferreira AJ, Raizada MK. Are we poised to target ACE2 for the next generation of antihypertensives? *J Mol Med*. 2008;86(6):685.
 64. Ferrario CM. ACE 2: More of Ang 1-7 or less Ang II? *Curr Opin Nephrol Hypertens*. 2011;20(1):1.
 65. Paulis L, Unger T. Novel therapeutic targets for hypertension. *Nat Rev Cardiol*. 2010;7(8):431.

66. Cullinane AB, Leung PS, Ortego J, Coca-Prados M, Harvey BJ. Renin-angiotensin system expression and secretory function in cultured human ciliary body non-pigmented epithelium. *Br J Ophthalmol*. 2002;86(6):676–83.
67. Vaajanen A. Expression and function of angiotensins in the regulation of intraocular pressure-an experimental study. 2009;
68. Vaajanen A, Vapaatalo H, Kautiainen H, Oksala O. Angiotensin (1-7) reduces intraocular pressure in the normotensive rabbit eye. *Invest Ophthalmol Vis Sci*. 2008;49(6):2557–62.
69. Mehta A, Iyer L, Parmar S, Shah G, Goyal R. Oculohypotensive effect of perindopril in acute and chronic models of glaucoma in rabbits. *Can J Physiol Pharmacol*. 2010;88(5):595–600.
70. LOTTI VJ, PAWLOWSKI N. Prostaglandins mediate the ocular hypotensive action of the angiotensin converting enzyme inhibitor MK-422 (enalaprilat) in African green monkeys. *J Ocul Pharmacol Ther*. 1990;6(1):1–7.
71. Loftsson T, Thorisdottir S, Fridriksdottir H, Stefansson E. Enalaprilat and enalapril maleate eyedrops lower intraocular pressure in rabbits. *Acta Ophthalmol*. 2010;88(3):337–41.
72. Haefliger IO, Flammer J, Lüscher TF. Nitric oxide and endothelin-1 are important regulators of human ophthalmic artery. *Invest Ophthalmol Vis Sci*. 1992;33(7):2340–3.
73. Costagliola C, Verolino M, De Rosa ML, Iaccarino G, Ciancaglini M, Mastropasqua L. Effect of oral losartan potassium administration on intraocular pressure in normotensive and glaucomatous human subjects. *Exp Eye Res*. 2000;71(2):167–71.
74. Inoue T, Yokoyama T, Mori Y, Sasaki Y, Hosokawa T, Yanagisawa H, et al. The effect of topical CS-088, an angiotensin AT1 receptor antagonist, on intraocular pressure and aqueous humor dynamics in rabbits. *Curr Eye Res*. 2001;23(2):133–8.
75. Shah GB, Sharma S, Mehta AA, Goyal RK. Oculohypotensive effect of angiotensin-converting enzyme inhibitors in acute and chronic models of glaucoma. *J Cardiovasc Pharmacol*. 2000;36(2):169–75.
76. Wang R-F, Podos SM, Mittag TW, Yokoyama T. Effect of CS-088, an angiotensin AT1 receptor antagonist, on intraocular pressure in glaucomatous monkey eyes. *Exp Eye Res*. 2005;80(5):629–32.

77. Semba K, Namekata K, Guo X, Harada C, Harada T, Mitamura Y. Renin–angiotensin system regulates neurodegeneration in a mouse model of normal tension glaucoma. *Cell Death Dis.* 2014;5(7):e1333–e1333.
78. Yang H, Hirooka K, Fukuda K, Shiraga F. Neuroprotective effects of angiotensin II type 1 receptor blocker in a rat model of chronic glaucoma. *Invest Ophthalmol Vis Sci.* 2009;50(12):5800–4.
79. Quigley HA, Pitha IF, Welsbie DS, Nguyen C, Steinhart MR, Nguyen TD, et al. Losartan treatment protects retinal ganglion cells and alters scleral remodeling in experimental glaucoma. *PLoS One.* 2015;10(10).
80. Mohammed QH, Haitham K, Waleed KA. Effect of Telmisartan on Intra-Ocular Pressure in induced Open Angle Glaucoma in Rabbits. *Int J Sci Res.* 2017;6(10):1656–9.
81. Kida T, Ikeda T, Nishimura M, Sugiyama T, Imamura Y, Sotozono C, et al. Renin-angiotensin system in proliferative diabetic retinopathy and its gene expression in cultured human Müller cells. *Jpn J Ophthalmol.* 2003;47(1):36–41.
82. Tanna AP, Lin AB. Medical therapy for glaucoma: what to add after a prostaglandin analogs? *Curr Opin Ophthalmol.* 2015;26(2):116–20.
83. Vaajanen A, Vapaatalo H. Local ocular renin–angiotensin system—a target for glaucoma therapy? *Basic Clin Pharmacol Toxicol.* 2011;109(4):217–24.

Table 1: Anti-glaucomatous agents and their influences on aqueous humour pathway [82]

Compound	IOP (%)	Aqueous production (%)	Aqueous out-flow (%)
β-blocker (non-selective)	↓ 25	↓ 33	
β-blocker (β1-selective)	↓ 20	↓ 24	
Direct miotic agents	↓ 20		↓ 24
adrenergic agonist (non-selective)	↓ 20		↓

α 2-agonist (selective)	↓25	↓35
Inhibitors of Carbonic anhydrase	↓25	↓35
Analogues of prostaglandin	↓30	↓100




Figure 1: Anatomy of the human aqueous humour pathway [83].




Figure 2 :Anatomy of a ciliary process [83]

UNDER PEER REVIEW