

Effect of Probiotics on Growth Performance and Carcass Characteristics in Broilers at Farmer's Door

Original Research Article

ABSTRACT

Aims: An experiment was conducted to investigate the comparative efficacy of two probiotics of different origins (yeast and bacterial based) on the growth performance and carcass characteristics of broilers chicken.

Place and Duration of Study: The study was conducted at Krishi Vigyan Kendra, Purnea between January, 2020 to March, 2020.

Methods: Two hundred and forty, day old chicks were kept and were randomly divided into 4 groups (A, B, C and D) and each group had 2 replicates of 20 chicks. Four (A, B, C and D) iso-caloric and iso-nitrogenous (ME 3000 kcal/ kg and CP 21.56%) broiler starter and finisher (ME 3000 kcal/ kg and CP 18.75%) diets were formulated containing *Saccharomyces Cerevisiae*, *Bacillus cereus toyoi* and mixture of both @ 0.2 % except group A which was control. Birds were raised for first week on commercial diet. Broiler starter and finisher diets were fed from 8-28 and 29-42 days, respectively.

Result: A significant ($P < 0.05$) effect was observed on overall performance of the birds fed diet containing probiotics. Probiotics in feed at 0.2% *Saccharomyces Cerevisiae* supplementation improved body weight gain (+12.7%), feed intake (3.8%) and feed conversion efficiency (-8.7%) compared with the control diet. Among groups, no significant differences were observed in carcass traits but highest dressing percentage were recorded in group-B followed by group-C than group-D and lowest in control group.

Conclusion: The result indicated that the addition of probiotics in feed containing 0.2% *Saccharomyces cerevisiae* improved broiler growth performances and it is beneficial to be used as supplement in feed of broiler chickens.

Keywords: *Saccharomyces Cerevisiae*, *Bacillus cereus toyoi*, growth performance, carcass characteristics, broilers

1. INTRODUCTION

There is a worldwide attempt to reduce antibiotic use in animal production because increased microbial resistance to antibiotic and residues in animal products can be harmful to consumers

*Corresponding Author's Email : hemantdec20@gmail.com

[1]. In recent years, several investigations were performed to find out the different alternatives of growth-promoting antimicrobials [2]. Those strategies were mainly focused on inhibition of proliferation of pathogenic bacteria and facilitation of beneficial gut micro-flora resulting in improved immunity, health status and performance of animals [3]. Probiotics are live microorganisms that affect the host animal by improving its intestinal microbial balance. Probiotic supplementation results in improved microbial population in the gut due to pathogen inhibition. Mechanisms of pathogen inhibition may include enhanced immune system, prevention of amine synthesis, neutralization of enterotoxin produced by pathogenic bacteria, competition for available nutrients, and bactericidal activity by secretion of inhibitory substances or competition for adhesion receptors to intestinal epithelium [4]. Healthy animals generally maintain a balanced microbial population that plays an important role in the growth and health of animals [5]. Some studies show that probiotics supplementation in feed of chickens improves the performance and it has been reported that probiotics were the most effective growth promoter [6].

The present study was carried out to evaluate effect of probiotics on growth performance and carcass characteristics in broilers.

2. MATERIALS AND METHODS

Day-old broiler chicks (n=240) were kept for experiment. Chicks were weighed on first day and checked for their physical health. The four poultry farmers from Purnea district were selected to conduct the present experiment in their farm. The 4 poultry farmers were divided into 4 groups having 40 chicks in each group. Trial was conducted in deep litter housing system and the managerial practices were kept control throughout the experimental period. To provide energy and to overcome the stress of transportation chicks were offered with a sugar solution (250 g sugar/liter water). Then, chicks were put in the brooding pen. The brooding temperature was maintained at 90°F during 1st week and it was then gradually lowered by 5 °F every week till it reached 75°F. In the first week commercial starter diet was fed to chicks. On 8th day, all birds were weighed individually and were randomly divided into four experimental groups (A, B, C and D) having 40 chicks in each group which were further divided into two replicates (20 chicks/replicate). Birds were vaccinated against Newcastle disease and Infectious Bursal disease.

Table1. Basal composition and analyzed results of the experimental diets

Ingredients	Starter diets				Finisher diets			
	A	B	C	D	A	B	C	D
Maize	59.5	59.3	59.3	59.3	42.00	42.00	42.00	42.00
Wheat	-	-	-	-	16.44	16.24	16.24	16.24
Soybean Meal	30.59	30.59	30.59	30.59	28.75	28.75	28.75	28.75
Fish Meal	4.58	4.58	4.58	4.58	-	-	-	-
Oil	2.5	2.5	2.5	2.5	2.5	2.5	2.5	2.5
Limestone	1.01	1.01	1.01	1.01	1.08	1.08	1.08	1.08
Dicalcium phosphate	0.87	0.87	0.87	0.87	1.08	1.08	1.08	1.08
Salt	0.27	0.27	0.27	0.27	0.3	0.3	0.3	0.3
Vitamin mix	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25
Mineral mixture	0.25	0.25	0.25	0.25	0.25	0.25	0.25	0.25
DL-methionine	0.18	0.18	0.18	0.18	0.07	0.07	0.07	0.07
<i>Saccharomyces cerevisiae</i>	-	0.2	-	-	-	0.2	-	-
<i>Bacillus cereus toyoi</i>	-	-	0.2	-	-	-	0.2	-
<i>Saccharomyces cerevisiae</i> + <i>Bacillus cereus toyoi</i>	-	-	-	0.1 + 0.1	-	-	-	0.1 + 0.1
Calculated value								
ME , Kcal/Kg	3000	3000	3000	3000	3000	3000	3000	3000
Crude protein, %	21.56	21.56	21.56	21.56	18.75	18.75	18.75	18.75

Two diets were prepared. Starter diet having CP 21.56 % and ME 3000 kcal/kg and in finisher diet CP 18.75 % and ME 3000 kcal/kg (Table 1). From 8-28 day birds were fed starter diet and finisher diet from 29-42 days. Birds of group A were fed diet without any supplementation (control group), whereas birds of group B, C and D were fed starter and finisher diets containing 0.2% *Saccharomyces cerevisiae*, *Bacillus cereus toyoi* and blend of both *Saccharomyces cerevisiae* and *Bacillus cereus toyoi* (0.1% + 0.1%), respectively.

During the trial weekly feed intake, weekly body weight was recorded and feed conversion ratio (FCR) of the birds was calculated. At the end of the experiment, 2 birds from each replicate were randomly selected and slaughtered to determine dressing percentage, breast and thigh meat yield and giblet organs weight. The data collected were used to calculate dressing percentage and organ weights (g organ weight/100 g body weight). The data was analyzed statistically [7].

3. RESULTS AND DISCUSSION

Weight gain at different periods is presented in table-2. Up to 28th day showed a significant increase in weight gain of broilers fed diets containing *Saccharomyces cerevisiae* (1265 gm) and *Bacillus cereus toyoi* (1245 gm) as compare to control group (1105 gm). At 42nd day highest weight gain was observed in group-B (2457 gm) followed by group-C (2419 gm) than Group D (2258 gm) and lowest in group A (2180 gm). All four groups were significantly differing to each other. These results confirmed the previous findings, who reported that dietary inclusion of probiotics in the diets of broilers showed improved body weight gain [8,9,10,11,12]. Therefore improvement in body weight gain of the birds in this study may be attributed to improved digestibility of crude protein, which may have contributed in better growth of the birds.

The growth performance of broiler starter, finisher and overall from the first day of experiment upto 42 day was recorded and presented in Table 2. Results showed improved feed intake due to the addition of *Saccharomyces cerevisiae*, *Bacillus cereus toyoi* and mixture of both at 0.2 % of feed compared to the control. The highest feed intake at 42nd day was observed in birds fed diet supplemented with *Saccharomyces cerevisiae* (4382 gm) followed by those fed diet supplemented with *Bacillus cereus toyoi* (4378 gm) and birds fed diet supplemented with mixture of both *Saccharomyces cerevisiae* and *Bacillus cereus toyoi* (4376 gm). All three supplemented groups were non-significant among themselves and significantly higher than control. Nawaz *et al.* also reported the increased feed intake in broilers fed diet supplemented with of *Saccharomyces cerevisiae* [8]. Results of the present study supported by the findings of who reported increased feed intake in broilers fed diet supplemented with different levels of *Saccharomyces cerevisiae* [13]. Results were also in accordance with those of Shareef *et al.* who used 1.0, 1.5 and 2.0% *Saccharomyces cerevisiae* in broiler diet and found a significant increase in feed intake [14]. Improved feed intake of those groups which are supplemented with probiotics in broiler feed [15].

Table 2. Growth performance and nutrient digestibility of broilers fed diets supplemented with *Saccharomyces cerevisiae*, *Bacillus cereus toyoi* and mixture of both *Saccharomyces cerevisiae* and *Bacillus cereus toyoi* (8-42 days)

Group	Weight gain (g)			Feed intake (g)			Feed conversion ratio		
	8-28	29-42	8-42	8-28	29-42	8-42	8-28	29-42	8-42
A	1105 ^c	1075 ^b	2180 ^d	1970 ^b	2249 ^b	4219 ^b	1.78 ^b	2.09 ^b	1.95 ^b
B	1265 ^a	1192 ^a	2457 ^a	2077 ^a	2305 ^a	4382 ^a	1.64 ^a	1.93 ^a	1.78 ^a
C	1245 ^a	1174 ^a	2419 ^b	2066 ^{ab}	2312 ^a	4378 ^a	1.66 ^{ab}	1.98 ^a	1.81 ^a
D	1180 ^b	1078 ^b	2258 ^c	2025 ^b	2351 ^a	4376 ^a	1.72 ^b	2.18 ^b	1.94 ^b
SEm±	9.68	6.34	7.84	12.16	12.46	22.76	0.02	0.01	0.02
C.D. (5%)	25.97	24.91	30.79	47.75	48.49	89.35	0.07	0.06	0.06

Mean under same superscript did not differ significantly

SEm= Standard error of mean

C.D. value= Critical difference value

The different values of FCR showed improvement in broilers fed diets containing *Saccharomyces cerevisiae* group and *Bacillus cereus toyoi* group compared to birds fed diets without probiotic supplementation (Table 2). The most efficient FCR was observed in birds of group B (1.78:1) fed diet containing *Saccharomyces cerevisiae*, followed by those of group C fed diet containing *Bacillus cereus toyoi* (1.81:1) and group D (1.94:1) fed diet containing mixture of both *Saccharomyces cerevisiae* + *Bacillus cereus toyoi* at the end of experiment. These results are in agreement with the findings [16] when they fed *Lactobacillus acidophilus* based probiotic to broiler chicks and observed that chicks fed probiotic cultures showed improved FCR than those control chicks. Dietary inclusion of yeast at 1.5%/kg of diet resulted in improved FCR [13]. Improved FCR of the birds using the yeast culture may be attributed to the digestion of crude protein, which enhanced growth of the birds resulting in improved efficiency of feed utilization. Improved FCR in broilers fed diet supplemented with of *Saccharomyces cerevisiae* [8]

Different values of carcass traits are presented in table-3. Among groups, no significant differences were observed in carcass traits but highest dressing percentage was recorded in group-B (66.94%) followed by group-C (66.14%) than group-D (65.55%) and lowest in control

group (65.01%). Similar trend were also found in breast meat percentage and thigh meat percentage were highest in group -B followed by group -C, group-D and group –A. Better dressing percentage in broilers reported when supplementation with probiotics [8,17]. Mutassim reported that supplementation of yeast increased breast meat yield in broilers [18]. The higher dressing percentage in birds fed diet containing yeast (*Saccharomyces cerevisiae*) may be due to higher body weight gain in the birds of this group compared to other treatment groups.

Table 3. Carcass characteristics of broilers fed diet *Saccharomyces cerevisiae*, *Bacillus cereus toyoi* and mixture of both *Saccharomyces cerevisiae* and *Bacillus cereus toyoi*

Group	Dressing (%)	Breast meat (%)	Thigh meat (%)	Abdominal fat (%)
A	65.01	20.31	20.26	2.87
B	66.94	21.32	21.89	1.99
C	66.14	20.99	20.83	2.05
D	65.55	20.71	20.51	2.47
SEm±	0.82	0.47	0.56	0.01
C.D. (5%)	2.74	1.61	1.56	0.05

4. CONCLUSION:

Under the conditions of the present study, probiotic supplementation at 0.2% *Saccharomyces cerevisiae* in broiler feed was effective in improving weight gain, feed intake, feed conversion efficiency and carcass traits. The results indicated that supplementation of current probiotic is beneficial in feed of broiler but optimal concentration of probiotics in broiler feed deserves further more investigations.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Yeo j and Kim KI. Effect of feeding diets containing an antibiotic, a probiotic or yucca extract on growth and intestinal urease activity in broiler chicks. *Poul. Sci.* 1997; **76**:381-385.

2. Huyghebaert G, Ducatelle R and Immerseel FV. An update on alternatives to antimicrobial growth promoters for broilers. *Vet J* . 2011; **187**: 182-188.
3. Adil S and Magray SN. Impact and manipulation of gut microflora in poultry: A review. *J Anim Vet Adv*. 2012; **11**: 873-877.
4. Lee K, Lillehoj HS and Siragusa GR. Direct-fed microbials and their impact on the intestinal microflora and immune system of chickens. *J Poultry Sci*. 2010; **47**: 106-114.
5. Ahmad A (2006). Effect of probiotic on broilers performance. *Int.J.Poult.Sci*. **5**: 593-597.
6. Mansoub HN. Comparative effect of butyric acid, probiotic and garlic on performance and serum composition of broiler hickens. *Am-Euras.J.Agric. & Environ.Sci*. 2011; **11**:507-511.
7. Snedecor GW and Cochran WG. Statistical Method. 8th end, Oxford and IBH Publishing Co. New Delhi.1994.
8. Nawaz H, Abbas IM, Ali M and Haq A. Effect of probiotics on growth performance, nutrient digestibility and carcass characteristics in broilers. *Jour. of Anim. & Plant Sci*, 2016; **26(3)**: 2016, Page: 599-604.
9. Falaki M, Shams Shargh M, Dastar B, Zrehdaran S. Effects of different levels of probiotic and prebiotic on performance and carcass characteristics of broiler chickens. *J. Anim. Vet. Adv.*, 2010; **9(18)**: 2390- 2395.
10. Zhang AW, Lee BD, Lee SK, Lee KW, Ann GH, Song KB and Lee CH. Effects of graded levels of dietary *Saccharomyces cerevisiae* on growth performance and meat quality in broiler chickens. *Asian-Aust. J. Anim. Sci*. 2005; **18**: 699-703.
11. Angel RR, Dalloul A and Doerr J. Performance of broiler chickens fed diets supplemented with a direct-fed microbial. *Poult. Sci*. 2005; **84**: 1222-1231.
12. Santin, E, Paulillo AC, Nakagui LSO, Alessi AC, Polverio WJC and Maiorka A. Evaluation of cell wall yeast as adsorbent of ochratoxin in broiler diets. *Int. J. Poult. Sci*. 2003; **2**: 465- 468.
13. Paryad, A and Mahmoudi M. Effect of different levels of supplemental yeast (*Saccharomyces cerevisiae*) on performance, blood constituents and carcass characteristics of broiler chicks. *Afr. J. Agri. Res*. 2008; **3**: 835-842.
14. Shareef AM and Al-Dabbagh ASA. Effect of probiotic (*Saccharomyces cerevisiae*) on performance of broiler chicks. *Iraqi J. Vet. Sci*. 2009; **23**: 23-29.

15. Shabani R, Nosrati M, Javandel F, Ali AAG and Kioumars H. The effect of probiotics on growth performance of broilers. *Anna. of Bio. Res.*, 2012; **3 (12)**:5450-5452.
16. Ramesh BK, Satyanarayana ML, Gowda RNS, Vijayasarithi SK and Rao S. Effect of *Lactobacillus acidophilus* on the growth of *Salmonella gallinarum* infected broilers. *Ind. J. Poult.Sci.*2000; **35**: 338-340.
17. Adejumo DO, Onifade AA and Afonja SA. Supplemental effects of dried yeast (Yea-sacc 1026 P®) in a low protein diet on growth performance, carcass characteristics and organ weights of broiler chicken. *Tropical Vet.* 2004; **22**: 72-77.
18. Mutassim, MA. Effects of feeding dry fat and yeast culture on broiler chicken performance. *Turk. J. Vet. Anim. Sci.* 2013; 37: 31-37.