

Ethno-ecological variability in the consumption of leafy green plants in the Republic of Benin

ABSTRACT

Beninese's dietary habits related to Amaranth (*Amaranthus cruentus*), African basil (*Ocimum gratissimum*) and African eggplant (*Solanum macrocarpon*) were studied in eleven districts of Republic of Benin. Data were collected from five hundred households on their consumption habits. Results showed that these leafy vegetables are consumed, either alone or accompanied by other leafy vegetables, depending on seasonal availability, health-promotion, and taste. Diversification and greater access to leafy green vegetables could improve individuals' nutritional profile.

KEYWORDS: Benin, dietary habits, food consumption, local criteria, green leafy vegetables.

1. INTRODUCTION

A diet rich in a diversity of plant resources has shown to enhance one's health and nutritional profile (Visweswari *et al.*, 2013). In Sub-Saharan Africa (SSA), the consumption of indigenous leafy vegetables plays an important role in the nutrition and health of rural poor because of general availability, richness in micronutrients, and for their medicinal virtues (Achigan-Dako *et al.*, 2010; Oluwalana *et al.*, 2011; Sossa-

Vihotogbé *et al.*, 2013). In the Republic of Benin, biodiversity inventories of traditional leafy vegetables reveal that more than 180 plant species are regularly cultivated, or semi-cultivated as wild plants, and play a significant role in daily nutrition (Dansi *et al.*, 2008). Most prominent among these plant species are Amaranth (*Amaranthus cruentus*, *Amaranthaceae*), African basil (*Ocimum gratissimum*, *Lamiaceae*) and African eggplant (*Solanum macrocarpon*, *Solanaceae*), which are valued for their nutritional and medicinal benefits (Dansi *et al.*, 2008; Achigan-Dako *et al.*, 2010; Vodouhè *et al.*, 2012a). For example, African basil is often used as spice and aromatic herbs for culinary purposes (Akpo-Djenontin *et al.*, 2016) and is also used in folk medicine for the treatment of upper respiratory tract infections, diarrhoea, skin diseases, pneumonia, cough, conjunctivitis intestinal worms and gastrointestinal disorders (Achigan-Dako *et al.*, 2010; Mahapatra *et al.*, 2010; Rahmatullah *et al.*, 2010). The leaves of Amaranth are used for treatment of colds, coughs, stomach aches, diarrhoea, skin rashes, back aches, constipation, fever, haemorrhage, kidney complaints and anaemia (Achigan-Dako *et al.*, 2010; Nana *et al.*, 2012). Medicinal uses of African eggplant include the treatment of tooth aches, allergic rhinitis, skin diseases, rheumatic diseases, joint pains, throat and stomach problems gastro-oesophageal reflux diseases, constipation, ulcers, diabetes and for weight reduction (Achigan-Dako *et al.*, 2010; Dougnon *et al.*, 2012; Muhammad and Shinkafi, 2014). These leafy vegetables exhibit antioxidant properties that offer protection against oxidative-stress related diseases (Adewale *et al.*, 2014; Acho *et al.*, 2015).

As popular as Amaranth, African basil and African eggplant are in the Republic of Benin, there is a lack of information on consumer's dietary habits and preservation techniques related to these plant resources. Although general information exists on their regional distribution, as well as their nutritional and medicinal values (Dansi *et al.*, 2008; Achigan-Dako *et al.*, 2010), ethno-nutritional information for these plant

resources is poorly documented (Vodouhè *et al.*, 2012a). Moreover, there has been no previous research conducted on the ethno-food knowledge, endogenous processing, or preservation methods of these species. Given the diversity of socio-cultural groups and agro-ecological zones in Benin (Achigan-Dako *et al.*, 2010; Yabi and Afouda, 2012; Segnon and Achigan-Dako, 2014) a rich knowledge exists on the various dietary habits and unique characteristics of these indigenous leafy green vegetables. Yet this knowledge is poorly documented. This study aimed at characterizing the diversity among local dietary habits through the administration of a household survey on the consumption of Amaranth, African basil and African eggplant across different age, cultural and geographical groups in Benin. This research is a prerequisite for the future promotion and development of indigenous leafy green vegetables for their health and food security benefits.

2. METHODOLOGY

2.1. Study area

This study was conducted in the Republic of Benin in West Africa (Figure 1: Benin map showing the geographical location of the survey sites). Benin is divided into three geographical areas (south, centre and north) and three major phyto-geographical regions (Guinean, Sudano-Guinean, and Sudanian) corresponding to the geographical areas (Achigan-Dako *et al.*, 2010; Segnon and Achigan-Dako, 2014). Southern and central areas are relatively humid agro-ecological zones with two rainy seasons, with a mean annual rainfall between 1100 to 1400 mm/year. The northern area comprises an arid and semi-arid agro-ecological zone characterized by one rainy season with irregular annual rainfall of 800 and 950 mm/year (Yabi and Afouda, 2012). Mean annual temperatures range from 26 to 28 °C across Benin but often reach 35 to 40 °C in far northern localities (Yabi and Afouda, 2012). Vegetation patterns show a gradient from south to north resulting from a combination of climate and

soils (Achigan-Dako *et al.*, 2010; Segnon and Achigan-Dako, 2014). The Guinean zone is a semi-deciduous rainforest zone with ferralitic soils located in the south while, Sudanian zone is a woodland and savanna region with ferruginous soils located in north. The Sudano-Guinean zone is a transitional zone characterized by a vegetation mosaic of forest islands, gallery forests and savannas (Achigan-Dako *et al.* 2010; Segnon and Achigan-Dako 2014). Inhabiting these regions are approximately 42 distinct socio-linguistic groups, the largest in terms of population being the Adja, Fon, Bariba, Berba, Dendi, Fulani, Goun, Idatcha, Lopka-Yom, Nagot-Yoruba and Waama (Achigan- Dako *et al.*, 2010; Segnon and Achigan-Dako, 2014).

2.2.Sociocultural characteristics of Republic of Benin population

Benin is constituted with several ethnic groups which numerical importance is variable. At the national level, the Fon group is the most important with 39.2% of Benin's population followed by Adja (15.2%), Yoruba (12.3%), Bariba (9.2%), Peulh (7%), Otamari (6.1%) Yoa (4%), Dendi (2.5%) and other ethnic groups (1.6%). The different ethnic groups are unequally distributed over the national territory. Socio-cultural groups dominated by Fon are strongly localized in the south of the country. In the north, Bariba, Dendi and Peulh coexist. The Adja are concentrated in the Southwest. In the center and South-East of the country, there are Yoruba and related. The most widely practiced religions in Benin are Catholicism (27.1%), Islam (24.4%) and Vodoun (17.3%).

2.3. Sampling

Field data were collected in October 2015 through December 2016 in eleven districts across three geographical areas (south: Abomey-Calavi, Bohicon, Cotonou, Djidja and Seme-kpodji; north-eastern: N'Dali, Parakou and Tchaourou and north-western: Boukoumbe, Djougou and Ouake); each characterised by phyto-geographical and socio-linguistic diversity (**Figure 1**: Benin map showing the geographical location of the survey sites; **Table 1**: List of villages surveyed and their socio-cultural groups). District selection was based on previous works which showed distribution of leafy vegetables in the country (Dansi *et al.*, 2008; Achigan-Dako *et al.*, 2010). Additional selection criteria included the existence of leafy vegetable fields-schools in some villages and a willingness to collaborate with research team. Thus, 64 villages, representing 24 socio-cultural groups (Adja, Ahoussa, Ani, Bariba, Biali, Ditamari, Dendi, Fon, Fulani, Goun, Gourmantche, Idatcha, Itcha, Kabiere, Lama, Lokpa, Mahi, Minan, Nagot, Pedah, Waama, Xlwa, Yom and Yoruba) were included in this study. (Figure 1: Benin map showing the geographical location of the survey sites; Table 1: List of villages surveyed and their socio-cultural groups). Households were randomly selected in each village, using bottle twirling method as described by Sossa-Vihotogbé *et al.*, (2012). From these groups, 500 women from various age and socio-cultural groups ($N=24$) and localities were identified. The survey was directed towards girls and women because of their knowledge of household food consumption patterns (Vorster *et al.*, 2007). Those interviewed include married, single, widowed or divorced women with basic and secondary education levels. Surveys were administered in the homes of informants, and in some cases with the aid of local interpreters.

2.4. Data collection

Questionnaires were administered in person to 500 respondents. Those respondents were asked to identify: (i) household uses of leafy

vegetables; (ii) consumption reasons; (iii) selection criteria; (iv) consumption frequencies; (v) processing methods; and (vi) traditional preservation techniques. The information recorded was based on local knowledge of Amaranth, African basil and African eggplant in relation with age, cultural and geographical groups of respondents.

3. RESULTS

3.1. Uses of leafy vegetables

Various species of traditional leafy vegetables were consumed within study areas (**Table 2:** Dietary uses of traditional leafy vegetables consumed in study areas and most of the households surveyed consumed a minimum of five and a maximum of 17 different leafy green vegetables. Results indicated that 95% of surveyed households consume Amaranth, 95% consume African basil, 96% consume African eggplant, and 88% consume all three. According to geographic areas, consumption of Amaranth was highest in the north (east: 100% and west: 97%) than south (91%). In contrast, consumption of African basil and African eggplant were higher in the south (98% and 99%, respectively) and northwest (98% and 99%, respectively) than north-east (88% and 88%, respectively). Depending on age groups, respondents over the age of 51 have a high consumption of Amaranth (98%), African basil (100%) and African eggplant (98%) when compared to the other age groups. The age group of 18-30 eighteen consumed more Amaranth (97%) than age group of 31-50 (92%). Inversely, level of consumption of African basil and African eggplant 18-30 year olds (94% and 95%, respectively) was similar to that of the 31-50 year old age group (95% for both vegetables).

All socio-cultural groups surveyed consumed the three leafy vegetables except Gourmantche socio-cultural group who did not consume African basil. Ani, Kabiere, Lama, Lokpa, Mina, Pedah, Waama, Xlwa and Yom socio-cultural groups were unanimous about these leafy vegetables consumption while some variations was found within the other socio-cultural groups.

According to the respondents, each of these vegetables is prepared alone (100 %) or in combination with one or both of the others (41%) or with other leafy vegetables species, such as leaves of Okra (*Abelmoschus esculentus*), African baobab (*Adansonia Digitata*), Jute mallow (*Corchorus olitorius*), Cassava (*Manihot esculenta*), Black benniseed (*Sesamum radiatum*) or Bitter leaf (*Vernonia amygdalina*).

3.2.Reasons for consumption

Consumption of the investigated vegetables was rooted in households' dietary habits or those of their in-laws and it is transmitted generationally. Cultural mixing has also changed dietary habits of Bariba socio-cultural group who, previously, consumed African eggplant fruits only rather than its leaves. However, a dietary change has been occurred among Baraiba households and now consuming African eggplant leaves as well as fruits. Bariba informants noted that this change is a result of greater interaction with Ditamari, Fon, and Nagot cultural groups, and the process of knowledge diffusion that has occurred. The reasons for this dietary change included favourable taste when added to traditional sauces as well as nutritive value (48%) related to their content in fibres, vitamins and minerals as well as their therapeutic properties (75%) against digestive disorders. For example, wrestlers from the Lokpa socio-cultural group consume African eggplant and its leaves, along with leaves of Okra and Cowpea (*Vigna unguiculata*), early in the morning of struggle festivals to acquire more strength before competition.

Consumption of African basil was found to be common among women within Bariba, Ditamari, Fon, Idatcha, Lama, Lokpa, Nagot, Waama, Yom and Yoruba socio-cultural groups. Consumption was noted to be most prevalent during periods of menstruation, pregnancy, and breastfeeding. Its therapeutic properties were multiple and were linked to its ability to remove blood clots, facilitate childbirth, clear waste after childbirth, treat postpartum infections, wound healing, and stimulate milk secretion.

In contrast, the non-consumption of the studied vegetables was related to lack of knowledge on processing methods (3.20%), bitter taste of African eggplant (2.40%), unpleasant odour of Amaranth (1.40%) and African basil (1.20%), and totemic considerations (0.40 %).

3.3.Selection criteria

The supply of leafy vegetables was secured through sharing (15.80 %), by purchase at markets (76.40 %), or at other production sites (home gardens: 35.60 %, gardening sites: 16.40 % and fields: 15.40 %). Amaranth and African eggplant were most often purchased at markets (70.60 % and 71.40 %, respectively), fields (15% and 12%, respectively) and gardening sites (15% for both species). Conversely, African basil is most often obtained from home gardens (33.80 %) and through sharing (15.40 %). The availability of these vegetables varied according to seasons and species although their overall availability is greater during the rainy season. Amaranth (87%) and African eggplant (77%) were more available throughout the year than is African basil (65%). The criteria used in the selection of these vegetables included therapeutic properties (72%), taste (68%), availability\seasonality (55%), time required for processing (32%) and cost (15%). Based on a scale of 1 to 5, respondents were asked to identify the main attribute that influence selection. Mean scores revealed that the main criterion for Amaranth was Availability (2.59), therapeutic properties (3.29) for African basil, and taste (2.60) for and African eggplant.

Therapeutic properties were an important selection criterion to Bariba, Gourmantche, Lokpa and Yom socio-cultural groups who lived in north-western part of the country. Inversely, respondents from Ani, Goun, Idatcha, Itcha, Nagot and Yoruba socio-cultural groups, living in southern areas and having between eighteen and thirty years old considered facility to transform of the leafy vegetables for their selection. Taste and cost were important criteria for Adja, Ahoussa, Fon, Kabiere, Mina and Xlwa socio-cultural groups, regardless their age and places of dwelling.

3.4.Frequency of consumption and processing methods

Consumption frequencies varied according to vegetables species but most respondents reported that consumption occurred once or twice per week. At these frequencies, consumption of Amaranth (36% for both frequencies), African basil (once per week: 38% and two or three times per week: 29%) and African eggplant (once per week: 32% and two or three times per week: 45.00 %) occurred regularly. The others consumption frequencies mentioned were 'four or five times per week' (Amaranth: 11%, African basil: 7% and African eggplant: 9%), 'once per day' (Amaranth: 1.4%, African basil: 2.4% and African eggplant: 0.8%) and 'once or twice per month' (Amaranth: 7%, African basil: 13% and African eggplant: 8%). These vegetables were used in processed form for sauces preparation with direct cooking (Amaranth: 62%, African basil: 49% and African eggplant: 75%) and pre-cooking (Amaranth: 47%, African basil: 28% and African eggplant: 36%) as the main processing methods. Fresh leaves are crushed on stone grinder (Amaranth: 4%, African basil: 40% and African eggplant: 7%) to create a paste which is used for sauces preparation. Dried leaves (Amaranth: 1%, African basil: 6% and African eggplant: 1%) and leaf powders (Amaranth: 5%, African basil: 3% and African eggplant: 1%) were also reported as ingredients used for sauces preparation.

Fresh leaves, crushed leaves, dried leaves and leaf powders were directly added to onion-based sauces (Amaranth) or tomato based sauce (Amaranth, African basil and African eggplant) supplemented with powder of Egusi (*Citrullus lanatus*) seeds after appropriate treatments making them palatable (Table 2: Dietary uses of traditional leafy vegetables consumed in study areas)

Scientific names (English names)	Local names ¹	Dietary utilization		
		Form of used	Processing methods	Beneficial effects
<i>Abelmoschus esculentus</i> (Okra)	Bariba: Kôbusa, Ditamari: Tinoufanti, Fon: Féviman, Lokpa: Maatou, Nagot: Ewéila.	Fresh leaves, dried leaves or dried leaf powders	Cooking in boiling water containing softening ingredient (potash or kanmu: alkaline rock salt) to get a slimy soup.	Very nutritious, stimulate appetite and manage digestive disorders.
<i>Adansonia digitata</i> (African baobab)	Bariba: Sônnawourousou, Ditamari: Titonankanti, Fon: Azizonman, Lokpa: Kôtôlassa, Nagot: Ewéotché.	Fresh leaves, dried leaves or dried leaf powders	Cooking in boiling water containing kanmu or potash to get a slimy soup.	Very nutritious, manage digestive disorders.
<i>Amaranthus cruentus</i> (Amaranth)	Bariba: Affônou, Ditamari: Aléfô, Fon: Fotètè, Lokpa: Aléfô, Nagot: Effô tètè.	Fresh leaves or pre-cooked (in salted boiling water)	Raw leaves are used as an ingredient of salad, addition of raw or pre-cooked form into tomato or onion-based sauce to get a vegetable soup.	Very nutritious, stimulate appetite and manage digestive disorders.
<i>Celosia argentea</i> (Cockscomb)	Bariba: Affônou, Ditamari: Tinonyawoti, Fon: Sôman, Nagot: Odjogodo.	Fresh leaves or pre-cooked (in salted boiling water)	Addition into tomato or onion-based sauce to get a vegetable soup.	Very nutritious.
<i>Ceratotheca sesamoides</i> (False sesame)	Bariba: Kpééwori, Ditamari: Siwadouanwe, Fon: Agbô, Lokpa: Hounoume, Nagot: Idjabô.	Fresh leaves or dried leaf powders	Addition into boiling water containing kanmu or potash to get a slimy soup.	Very nutritious, manage digestive disorders.
<i>Corchorus olitorius</i> (Jute mallow)	Bariba: Yôyôkou, Ditamari: Tifanwounti, Fon: Ninnouwi, Lokpa: Ayôyôkou, Nagot: Owoyô.	Fresh leaves	Addition into boiling water containing kanmu or potash to get a slimy soup.	Very nutritious, restore appetite and manage digestive disorders.

Scientific names (English names)	Local names ¹	Dietary utilization		
		Form of used	Processing methods	Beneficial effects
<i>Hibiscus sabdariffa</i> (Roselle)	Bariba: Sinri, Ditamari: Tikouan n'ti, Fon: Sinku, Lokpa: Ankpaman, Nagot: Kpakpa.	Pre-cooked leaves (in boiling water containing kanmu or potash)	Addition into tomato-based sauce to get a vegetable soup.	Very nutritious, manage digestives disorders.
<i>Justicia tenella</i>	Bariba: Kourôkountônu, Ditamari: Tinoukounti, Lokpa: Tiletoussi, Nagot: Djagoudjagou.	Fresh leaves or pre- cooked (in boiling water containing kanmu or potash)	Addition into tomato-based sauce to get a vegetable soup.	Very nutritious
<i>Manihot esculenta</i> (Cassava)	Bariba: Kpakiwourousou, Ditamari: Tikôkônouwôti, Fon: Finyinman, Lokpa: Agbedehatou, Nagot: Ewékpaki.	Pre-cooked leaves (in boiling water containing kanmu or potash)	Addition into tomato-based sauce to get a vegetable soup.	Very nutritious.
<i>Moringa oleifera</i> (Drum-stick tree)	Bariba: Yorouyara, Ditamari: Mounpèkom, Fon: Kpatiman, Lokpa: Lôtaha, Nagot: Lagalanga.	Fresh leaves or dried leaf powders	Fresh leaves are used to make teas and as an ingredient of salad and vegetable soup. Dried leaf powders are used as diet supplements.	Very nutritious, stimulate appetite and manage digestives disorders.
<i>Ocimum gratissimum</i> (African basil)	Bariba: Danbakarou, Ditamari: Tinassiyinti, Fon: Tchiayo, Lokpa: Assôhou, Nagot: Simonba.	Fresh leaves, crushed or pre- cooked (in boiling water containing kanmu or potash)	Raw leaves are used to make teas and as a seasoning, addition of crushed or pre- cooked leaves into tomato or onion-based sauce to get a vegetable soup.	Very nutritious, manage digestives disorders.
<i>Sesamum radiatum</i> (Black benniseed)	Bariba: Dossi, Ditamari: Tiwadounati, Fon: Akanmanku, Lokpa: Touhoonôme, Nagot: Dossé, Goolowo.	Fresh leaves or dried leaf powders	Cooking in boiling water containing kanmu or potash to get a slimy soup.	Very nutritious, manage digestives disorders.
<i>Solanum macrocarpon</i> (African eggplant)	Bariba: Sambinouwourousou, Ditamari: Tikawounfanti, Fon: Gboman, Lokpa: Mètècou,	Pre-cooked leaves (in boiling water containing kanmu	Addition into tomato or onion- based sauce to get a vegetable soup.	Very nutritious, manage digestives disorders.

Scientific names (English names)	Local names ¹	Dietary utilization		
		Form of used	Processing methods	Beneficial effects
	Nagot: Katakoukpakou.	or potash)		
<i>Talinum triangulare</i> (Water leaf)	Bariba: Odôndôn, Ditamari: Yêmontouo, Fon: Aglassoeman, Lokpa: Kamplékankann'dê, Nagot: Odôndôn.	Fresh leaves or pre-cooked (in boiling water containing kanmu or potash) and dried leaves	Addition of pre-cooked leaves or dried leaves into tomato-based sauce to get a vegetable soup.	Very nutritious, manage digestives disorders.
<i>Vernonia amygdalina</i> (Bitter leaf)	Bariba: Touwan, Ditamari: Souwaka, Fon: Amanvivè, Lokpa: Souwaka, Nagot: Eéwoh.	Pre-cooked leaves (in boiling water containing kanmu or potash)	Addition into tomato-based sauce to get a vegetable soup.	Very nutritious, manage digestives disorders.
<i>Vigna unguiculata</i> (Cowpea)	Bariba: Suiwourousou, Ditamari: Titouti, Fon: Ayiman, Lokpa: Tchaasé, Nagot: Oson.	Pre-cooked leaves (in boiling water containing kanmu or potash)	Addition into tomato-based sauce to get a vegetable soup.	Very nutritious, stimulate appetite and manage digestives disorders.
<i>Vitex doniana</i> (Black plum)	Bariba: Kounonkou, Ditamari: Timantoun n'ti, Fon: Fonman, Nagot: Ewa.	Fresh leaves or pre-cooked (in boiling water containing kanmu or potash)	Fresh leaves are used to make teas, addition of pre-cooked form into tomato-based sauce to get a vegetable soup.	Very nutritious, manage digestives disorders.

Sources: survey in October 2015 and December 2016 and additional information from Dansi *et al.* (2008) and Achigan-Dako *et al.* (2010)

¹: choice of local languages is related to the main cultural groups encountered [Bariba (17.40 %), Ditamari (11.60 %), Fon (19.40 %), Lokpa (9.20 %) and Nagot (9.60 %)].

Table 3: Various types of sauces obtained from each leafy vegetable). This processing method is used by most of socio-cultural groups encountered except Adja and Gourmantche socio-cultural groups who did not apply it to any of these vegetables. Similarly, Itcha, Pedah and Xlwa socio-cultural groups did not apply it to African basil and African eggplant while Kabiere and Mahi people did not apply it to African basil alone and Yoruba people to African eggplant alone. Pre-cooking in boiling water is applied to fresh leaves and requires appropriate treatments before pre-cooked leaves can be added to a tomato-based sauce supplemented with Egusi seeds powder (Table 2: Dietary uses of traditional leafy vegetables consumed in study areas

Scientific names (English names)	Local names ¹	Dietary utilization		
		Form of used	Processing methods	Beneficial effects
<i>Abelmoschus esculentus</i> (Okra)	Bariba: Kôbusa, Ditamari: Tinoufanti, Fon: Féviman, Lokpa: Maatou, Nagot: Ewéila.	Fresh leaves, dried leaves or dried leaf powders	Cooking in boiling water containing softening ingredient (potash or kanmu: alkaline rock salt) to get a slimy soup.	Very nutritious, stimulate appetite and manage digestives disorders.
<i>Adansonia digitata</i> (African baobab)	Bariba: Sônnawourousou, Ditamari: Titonankanti, Fon: Azizonman, Lokpa: Kôtôlassa, Nagot: Ewéotché.	Fresh leaves, dried leaves or dried leaf powders	Cooking in boiling water containing kanmu or potash to get a slimy soup.	Very nutritious, manage digestives disorders.
<i>Amaranthus cruentus</i> (Amaranth)	Bariba: Affônou, Ditamari: Aléfô, Fon: Fotètè, Lokpa: Aléfô, Nagot: Effô tètè.	Fresh leaves or pre-cooked (in salted boiling water)	Raw leaves are used as an ingredient of salad, addition of raw or pre-cooked form into tomato or onion-based sauce to get a vegetable soup.	Very nutritious, stimulate appetite and manage digestives disorders.
<i>Celosia argentea</i> (Cockscomb)	Bariba: Affônou, Ditamari: Tinonyawoti, Fon: Sôman, Nagot: Odjogodo.	Fresh leaves or pre-cooked (in salted boiling water)	Addition into tomato or onion-based sauce to get a vegetable soup.	Very nutritious.
<i>Ceratotheca sesamoides</i> (False sesame)	Bariba: Kpééwori, Ditamari: Siwadouanwe, Fon: Agbô, Lokpa: Hounoume, Nagot: Idjabô.	Fresh leaves or dried leaf powders	Addition into boiling water containing kanmu or potash to get a slimy soup.	Very nutritious, manage digestives disorders.
<i>Corchorus olitorius</i> (Jute mallow)	Bariba: Yôyôkou, Ditamari: Tifanwounti, Fon: Ninnouwi, Lokpa: Ayôyôkou, Nagot: Owoyô.	Fresh leaves	Addition into boiling water containing kanmu or potash to get a slimy soup.	Very nutritious, restore appetite and manage digestives disorders.

Scientific names (English names)	Local names ¹	Dietary utilization		
		Form of used	Processing methods	Beneficial effects
<i>Hibiscus sabdariffa</i> (Roselle)	Bariba: Sinri, Ditamari: Tikouan n'ti, Fon: Sinku, Lokpa: Ankpaman, Nagot: Kpakpa.	Pre-cooked leaves (in boiling water containing kanmu or potash)	Addition into tomato-based sauce to get a vegetable soup.	Very nutritious, manage digestives disorders.
<i>Justicia tenella</i>	Bariba: Kourôkountônu, Ditamari: Tinoukounti, Lokpa: Tiletoussi, Nagot: Djagoudjagou.	Fresh leaves or pre- cooked (in boiling water containing kanmu or potash)	Addition into tomato-based sauce to get a vegetable soup.	Very nutritious
<i>Manihot esculenta</i> (Cassava)	Bariba: Kpakiwourousou, Ditamari: Tikôkônouwôti, Fon: Finyinman, Lokpa: Agbedehatou, Nagot: Ewékpaki.	Pre-cooked leaves (in boiling water containing kanmu or potash)	Addition into tomato-based sauce to get a vegetable soup.	Very nutritious.
<i>Moringa oleifera</i> (Drum-stick tree)	Bariba: Yorouyara, Ditamari: Mounpèkom, Fon: Kpatiman, Lokpa: Lôtaha, Nagot: Lagalanga.	Fresh leaves or dried leaf powders	Fresh leaves are used to make teas and as an ingredient of salad and vegetable soup. Dried leaf powders are used as diet supplements.	Very nutritious, stimulate appetite and manage digestives disorders.
<i>Ocimum gratissimum</i> (African basil)	Bariba: Danbakarou, Ditamari: Tinassiyinti, Fon: Tchiayo, Lokpa: Assôhou, Nagot: Simonba.	Fresh leaves, crushed or pre- cooked (in boiling water containing kanmu or potash)	Raw leaves are used to make teas and as a seasoning, addition of crushed or pre- cooked leaves into tomato or onion-based sauce to get a vegetable soup.	Very nutritious, manage digestives disorders.
<i>Sesamum radiatum</i> (Black benniseed)	Bariba: Dossi, Ditamari: Tiwadounati, Fon: Akanmanku, Lokpa: Touhoonôme, Nagot: Dossé, Goolowo.	Fresh leaves or dried leaf powders	Cooking in boiling water containing kanmu or potash to get a slimy soup.	Very nutritious, manage digestives disorders.
<i>Solanum macrocarpon</i> (African eggplant)	Bariba: Sambinouwourousou, Ditamari: Tikawounfanti, Fon: Gboman, Lokpa: Mètècou,	Pre-cooked leaves (in boiling water containing kanmu	Addition into tomato or onion- based sauce to get a vegetable soup.	Very nutritious, manage digestives disorders.

Scientific names (English names)	Local names ¹	Dietary utilization		
		Form of used	Processing methods	Beneficial effects
	Nagot: Katakoukpakou.	or potash)		
<i>Talinum triangulare</i> (Water leaf)	Bariba: Odôndôn, Ditamari: Yêmontouo, Fon: Aglassoeman, Lokpa: Kamplékankann 'dê, Nagot: Odôndôn.	Fresh leaves or pre-cooked (in boiling water containing kanmu or potash) and dried leaves	Addition of pre-cooked leaves or dried leaves into tomato-based sauce to get a vegetable soup.	Very nutritious, manage digestives disorders.
<i>Vernonia amygdalina</i> (Bitter leaf)	Bariba: Touwan, Ditamari: Souwaka, Fon: Amanvivè, Lokpa: Souwaka, Nagot: Eéwoh.	Pre-cooked leaves (in boiling water containing kanmu or potash)	Addition into tomato-based sauce to get a vegetable soup.	Very nutritious, manage digestives disorders.
<i>Vigna unguiculata</i> (Cowpea)	Bariba: Suiwourousou, Ditamari: Titouti, Fon: Ayiman, Lokpa: Tchaasé, Nagot: Oson.	Pre-cooked leaves (in boiling water containing kanmu or potash)	Addition into tomato-based sauce to get a vegetable soup.	Very nutritious, stimulate appetite and manage digestives disorders.
<i>Vitex doniana</i> (Black plum)	Bariba: Kounonkou, Ditamari: Timantoun n'ti, Fon: Fonman, Nagot: Ewa.	Fresh leaves or pre-cooked (in boiling water containing kanmu or potash)	Fresh leaves are used to make teas, addition of pre-cooked form into tomato-based sauce to get a vegetable soup.	Very nutritious, manage digestives disorders.

Sources: survey in October 2015 and December 2016 and additional information from Dansi *et al.* (2008) and Achigan-Dako *et al.* (2010)

¹: choice of local languages is related to the main cultural groups encountered [Bariba (17.40 %), Ditamari (11.60 %), Fon (19.40 %), Lokpa (9.20 %) and Nagot (9.60 %)].

Table 3: Various types of sauces obtained from each leafy vegetable). Pre-cooking requires use of additives like potash or bicarbonate of soda to maintain green color, to accelerate softening of African basil and African eggplant, and reduce bitterness of African eggplant. Pre-cooking is also practiced by most of socio-cultural groups met except Waama socio-cultural group who did not apply it to any of vegetables. Similarly, Ahoussa, Biali, Fulani and Gourmantche socio-cultural groups did not apply it to African basil. Sauces obtained from pre-cooking method and made of association of different leafy vegetables species require that each vegetable type was pre-cooked separately because of variability in pre-cooking time, thereafter they can be mixed together during processing. Concerning crushed form, fresh leaves were properly processed to obtain a paste which was added to onion-based sauce supplemented with Egusi seeds powder (Table 2: Dietary uses of traditional leafy vegetables consumed in study areas

Scientific names (English names)	Local names ¹	Dietary utilization		
		Form of used	Processing methods	Beneficial effects
<i>Abelmoschus esculentus</i> (Okra)	Bariba: Kôbusa, Ditamari: Tinoufanti, Fon: Féviman, Lokpa: Maatou, Nagot: Ewéila.	Fresh leaves, dried leaves or dried leaf powders	Cooking in boiling water containing softening ingredient (potash or kanmu: alkaline rock salt) to get a slimy soup.	Very nutritious, stimulate appetite and manage digestives disorders.
<i>Adansonia digitata</i> (African baobab)	Bariba: Sônnawourousou, Ditamari: Titonankanti, Fon: Azizonman, Lokpa: Kô Tôlassa, Nagot: Ewéotché.	Fresh leaves, dried leaves or dried leaf powders	Cooking in boiling water containing kanmu or potash to get a slimy soup.	Very nutritious, manage digestives disorders.
<i>Amaranthus cruentus</i> (Amaranth)	Bariba: Affônou, Ditamari: Aléfô, Fon: Fotètè, Lokpa: Aléfô, Nagot: Effô tètè.	Fresh leaves or pre-cooked (in salted boiling water)	Raw leaves are used as an ingredient of salad, addition of raw or pre-cooked form into tomato or onion-based sauce to get a vegetable soup.	Very nutritious, stimulate appetite and manage digestives disorders.
<i>Celosia argentea</i> (Cockscomb)	Bariba: Affônou, Ditamari: Tinonyawoti, Fon: Sôman, Nagot: Odjogodo.	Fresh leaves or pre-cooked (in salted boiling water)	Addition into tomato or onion-based sauce to get a vegetable soup.	Very nutritious.
<i>Ceratotheca sesamoides</i> (False sesame)	Bariba: Kpééwori, Ditamari: Siwadouanwe, Fon: Agbô, Lokpa: Hounoume, Nagot: Idjabô.	Fresh leaves or dried leaf powders	Addition into boiling water containing kanmu or potash to get a slimy soup.	Very nutritious, manage digestives disorders.

Scientific names (English names)	Local names ¹	Dietary utilization		
		Form of used	Processing methods	Beneficial effects
<i>Corchorus olitorius</i> (Jute mallow)	Bariba: Yôyôkou, Ditamari: Tifanwounti, Fon: Ninnouwi, Lokpa: Ayôyôkou, Nagot: Owoyô.	Fresh leaves	Addition into boiling water containing kanmu or potash to get a slimy soup.	Very nutritious, restore appetite and manage digestives disorders.
<i>Hibiscus sabdariffa</i> (Roselle)	Bariba: Sinri, Ditamari: Tikouan n'ti, Fon: Sinku, Lokpa: Ankpaman, Nagot: Kpakpa.	Pre-cooked leaves (in boiling water containing kanmu or potash)	Addition into tomato-based sauce to get a vegetable soup.	Very nutritious, manage digestives disorders.
<i>Justicia tenella</i>	Bariba: Kourôkountônu, Ditamari: Tinoukounti, Lokpa: Tiletoussi, Nagot: Djagoudjagou.	Fresh leaves or pre-cooked (in boiling water containing kanmu or potash)	Addition into tomato-based sauce to get a vegetable soup.	Very nutritious
<i>Manihot esculenta</i> (Cassava)	Bariba: Kpakiwourousou, Ditamari: Tikôkônouwôti, Fon: Finyinman, Lokpa: Agbedehatou, Nagot: Ewékpakpi.	Pre-cooked leaves (in boiling water containing kanmu or potash)	Addition into tomato-based sauce to get a vegetable soup.	Very nutritious.
<i>Moringa oleifera</i> (Drum-stick tree)	Bariba: Yorouyara, Ditamari: Mounpèkom, Fon: Kpatiman, Lokpa: Lôtaha, Nagot: Lagalanga.	Fresh leaves or dried leaf powders	Fresh leaves are used to make teas and as an ingredient of salad and vegetable soup. Dried leaf powders are used as diet supplements.	Very nutritious, stimulate appetite and manage digestives disorders.
<i>Ocimum gratissimum</i> (African basil)	Bariba: Danbakarou, Ditamari: Tinassiyinti, Fon: Tchiayo, Lokpa: Assôhou, Nagot: Simonba.	Fresh leaves, crushed or pre-cooked (in boiling water containing kanmu or potash)	Raw leaves are used to make teas and as a seasoning, addition of crushed or pre-cooked leaves into tomato or onion-based sauce to get a vegetable soup.	Very nutritious, manage digestives disorders.
<i>Sesamum radiatum</i> (Black benniseed)	Bariba: Dossi, Ditamari: Tiwadounati, Fon: Akanmanku, Lokpa: Touhoonôme,	Fresh leaves or dried leaf powders	Cooking in boiling water containing kanmu or potash to get a slimy soup.	Very nutritious, manage digestives disorders.

Scientific names (English names)	Local names ¹	Dietary utilization		
		Form of used	Processing methods	Beneficial effects
	Nagot: Dossé, Goolowo.			
<i>Solanum macrocarpon</i> (African eggplant)	Bariba: Sambinouwourousou, Ditamari: Tikawounfanti, Fon: Gboman, Lokpa: Mètècou, Nagot: Katakoukpakou.	Pre-cooked leaves (in boiling water containing kanmu or potash)	Addition into tomato or onion- based sauce to get a vegetable soup.	Very nutritious, manage digestives disorders.
<i>Talinum triangulare</i> (Water leaf)	Bariba: Odôndôn, Ditamari: Yêmontouo, Fon: Aglassoeman, Lokpa: Kamplékankann'dê, Nagot: Odôndôn.	Fresh leaves or pre- cooked (in boiling water containing kanmu or potash) and dried leaves	Addition of pre-cooked leaves or dried leaves into tomato- based sauce to get a vegetable soup.	Very nutritious, manage digestives disorders.
<i>Vernonia amygdalina</i> (Bitter leaf)	Bariba: Touwan, Ditamari: Souwaka, Fon: Amanvivè, Lokpa: Souwaka, Nagot: Eéwoh.	Pre-cooked leaves (in boiling water containing kanmu or potash)	Addition into tomato-based sauce to get a vegetable soup.	Very nutritious, manage digestives disorders.
<i>Vigna unguiculata</i> (Cowpea)	Bariba: Suiwourousou, Ditamari: Titouti, Fon: Ayiman, Lokpa: Tchaasé, Nagot: Oson.	Pre-cooked leaves (in boiling water containing kanmu or potash)	Addition into tomato-based sauce to get a vegetable soup.	Very nutritious, stimulate appetite and manage digestives disorders.
<i>Vitex doniana</i> (Black plum)	Bariba: Kounonkou, Ditamari: Timantoun n'ti, Fon: Fonman, Nagot: Ewa.	Fresh leaves or pre- cooked (in boiling water containing kanmu or potash)	Fresh leaves are used to make teas, addition of pre-cooked form into tomato-based sauce to get a vegetable soup.	Very nutritious, manage digestives disorders.

Sources: survey in October 2015 and December 2016 and additional information from Dansi *et al.* (2008) and Achigan-Dako *et al.* (2010)

¹: choice of local languages is related to the main cultural groups encountered [Bariba (17.40 %), Ditamari (11.60 %), Fon (19.40 %), Lokpa (9.20 %) and Nagot (9.60 %)].

Table 3: Various types of sauces obtained from each leafy vegetable). This method was known and used by most of socio-cultural groups except Mahi and Mina. Consumption of African basil under this form is used by pregnant and lactating women within Ani, Bariba, Ditamari, Fon, Idatcha, Lokpa, Nagot and Yoruba socio-cultural groups. Use of dried leaf and leaf powders for sauces preparation was common within Dendi, Ditamari, Fon, Lokpa and Yom socio-cultural groups. Many others ingredients were used during the preparation of sauces to enhance taste and thickness (spices, small fry fish, fermented African locus bean and Egusi seeds powder). These sauces accompanied cereal-based dishes (maize, millet, sorghum or rice paste; Akassa; Lio; rice; etc.) or tubers-based dishes (Amala, Eba, pounded yam, etc.).

Processing methods of each leafy vegetable require some unit operations (Table 2: Dietary uses of traditional leafy vegetables consumed in study areas

Scientific names (English names)	Local names ¹	Dietary utilization		
		Form of used	Processing methods	Beneficial effects
<i>Abelmoschus esculentus</i> (Okra)	Bariba: Kôbusa, Ditamari: Tinoufanti, Fon: Féviman, Lokpa: Maatou, Nagot: Ewéila.	Fresh leaves, dried leaves or dried leaf powders	Cooking in boiling water containing softening ingredient (potash or kanmu: alkaline rock salt) to get a slimy soup.	Very nutritious, stimulate appetite and manage digestives disorders.
<i>Adansonia digitata</i> (African baobab)	Bariba: Sônnawourousou, Ditamari: Titonankanti, Fon: Azizonman, Lokpa: Kôtôlassa, Nagot: Ewéotché.	Fresh leaves, dried leaves or dried leaf powders	Cooking in boiling water containing kanmu or potash to get a slimy soup.	Very nutritious, manage digestives disorders.
<i>Amaranthus cruentus</i> (Amaranth)	Bariba: Affônou, Ditamari: Aléfô, Fon: Fotètè, Lokpa: Aléfô, Nagot: Effô tètè.	Fresh leaves or pre-cooked (in salted boiling water)	Raw leaves are used as an ingredient of salad, addition of raw or pre-cooked form into tomato or onion-based sauce to get a vegetable soup.	Very nutritious, stimulate appetite and manage digestives disorders.
<i>Celosia argentea</i> (Cockscomb)	Bariba: Affônou, Ditamari: Tinonyawoti,	Fresh leaves or pre-cooked (in salted	Addition into tomato or onion-based sauce to get a vegetable	Very nutritious.

Scientific names (English names)	Local names ¹	Dietary utilization		
		Form of used	Processing methods	Beneficial effects
	Fon: Sôman, Nagot: Odjogodo.	boiling water)	soup.	
<i>Ceratotheca sesamoides</i> (False sesame)	Bariba: Kpééwori, Ditamari: Siwadouanwe, Fon: Agbô, Lokpa: Hounoume, Nagot: Idjabô.	Fresh leaves or dried leaf powders	Addition into boiling water containing kanmu or potash to get a slimy soup.	Very nutritious, manage digestives disorders.
<i>Corchorus olitorius</i> (Jute mallow)	Bariba: Yôyôkou, Ditamari: Tifanwounti, Fon: Ninnouwi, Lokpa: Ayôyôkou, Nagot: Owoyô.	Fresh leaves	Addition into boiling water containing kanmu or potash to get a slimy soup.	Very nutritious, restore appetite and manage digestives disorders.
<i>Hibiscus sabdariffa</i> (Roselle)	Bariba: Sinri, Ditamari: Tikouan n'ti, Fon: Sinku, Lokpa: Ankpaman, Nagot: Kpakpa.	Pre-cooked leaves (in boiling water containing kanmu or potash)	Addition into tomato-based sauce to get a vegetable soup.	Very nutritious, manage digestives disorders.
<i>Justicia tenella</i>	Bariba: Kourôkountônu, Ditamari: Tinoukounti, Lokpa: Tiletoussi, Nagot: Djagoudjagou.	Fresh leaves or pre- cooked (in boiling water containing kanmu or potash)	Addition into tomato-based sauce to get a vegetable soup.	Very nutritious
<i>Manihot esculenta</i> (Cassava)	Bariba: Kpakiwourousou, Ditamari: Tikôkônouwôti, Fon: Finyinman, Lokpa: Agbedehatou, Nagot: Ewékpaki.	Pre-cooked leaves (in boiling water containing kanmu or potash)	Addition into tomato-based sauce to get a vegetable soup.	Very nutritious.
<i>Moringa oleifera</i> (Drum-stick tree)	Bariba: Yorouyara, Ditamari: Mounpèkom, Fon: Kpatiman, Lokpa: Lôtaha, Nagot: Lagalanga.	Fresh leaves or dried leaf powders	Fresh leaves are used to make teas and as an ingredient of salad and vegetable soup. Dried leaf powders are used as diet supplements.	Very nutritious, stimulate appetite and manage digestives disorders.

Scientific names (English names)	Local names ¹	Dietary utilization		
		Form of used	Processing methods	Beneficial effects
<i>Ocimum gratissimum</i> (African basil)	Bariba: Danbakarou, Ditamari: Tinassiyinti, Fon: Tchiayo, Lokpa: Assôhou, Nagot: Simonba.	Fresh leaves, crushed or pre-cooked (in boiling water containing kanmu or potash)	Raw leaves are used to make teas and as a seasoning, addition of crushed or pre-cooked leaves into tomato or onion-based sauce to get a vegetable soup.	Very nutritious, manage digestives disorders.
<i>Sesamum radiatum</i> (Black benniseed)	Bariba: Dossi, Ditamari: Tiwadounati, Fon: Akanmanku, Lokpa: Touhoonôme, Nagot: Dossé, Goolowo.	Fresh leaves or dried leaf powders	Cooking in boiling water containing kanmu or potash to get a slimy soup.	Very nutritious, manage digestives disorders.
<i>Solanum macrocarpon</i> (African eggplant)	Bariba: Sambinouwourousou, Ditamari: Tikawounfanti, Fon: Gboman, Lokpa: Mètècou, Nagot: Katakoukpakou.	Pre-cooked leaves (in boiling water containing kanmu or potash)	Addition into tomato or onion-based sauce to get a vegetable soup.	Very nutritious, manage digestives disorders.
<i>Talinum triangulare</i> (Water leaf)	Bariba: Odôndôn, Ditamari: Yêmontouo, Fon: Aglassoeman, Lokpa: Kamplékankann'dê, Nagot: Odôndôn.	Fresh leaves or pre-cooked (in boiling water containing kanmu or potash) and dried leaves	Addition of pre-cooked leaves or dried leaves into tomato-based sauce to get a vegetable soup.	Very nutritious, manage digestives disorders.
<i>Vernonia amygdalina</i> (Bitter leaf)	Bariba: Touwan, Ditamari: Souwaka, Fon: Amanvivè, Lokpa: Souwaka, Nagot: Eéwoh.	Pre-cooked leaves (in boiling water containing kanmu or potash)	Addition into tomato-based sauce to get a vegetable soup.	Very nutritious, manage digestives disorders.
<i>Vigna unguiculata</i> (Cowpea)	Bariba: Suiwourousou, Ditamari: Titouti, Fon: Ayiman, Lokpa: Tchaasé, Nagot: Oson.	Pre-cooked leaves (in boiling water containing kanmu or potash)	Addition into tomato-based sauce to get a vegetable soup.	Very nutritious, stimulate appetite and manage digestives disorders.
<i>Vitex doniana</i> (Black plum)	Bariba: Kounonkou, Ditamari: Timantoun n'ti, Fon: Fonman, Nagot: Ewa.	Fresh leaves or pre-cooked (in boiling water containing	Fresh leaves are used to make teas, addition of pre-cooked form into tomato-based sauce	Very nutritious, manage digestives disorders.

Scientific names (English names)	Local names ¹	Dietary utilization		
		Form of used kanmu or potash)	Processing methods to get a vegetable soup.	Beneficial effects

Sources: survey in October 2015 and December 2016 and additional information from Dansi *et al.* (2008) and Achigan-Dako *et al.* (2010)

¹: choice of local languages is related to the main cultural groups encountered [Bariba (17.40 %), Ditamari (11.60 %), Fon (19.40 %), Lokpa (9.20 %) and Nagot (9.60 %)].

1 Table 3: Various types of sauces obtained from each leafy vegetable) among which
2 washing and cooking were reported as the most important steps for Amaranth (59% and 36%,
3 respectively), African basil (54% and 38%, respectively) and African eggplant (52% and
4 47%, respectively). Washing of African basil can be associated with trituration (10%) and this
5 combination is an important step. Sorting was also considered an important step for
6 processing of Amaranth (20%), African basil (13%) and African eggplant (15%). It is
7 essential to carefully perform each unit operation to avoid digestive disorders and vomiting
8 after consumption of sauces made with Amaranth (71% and 17%, respectively), African basil
9 (67% and 17%, respectively) and African eggplant (65% and 18%, respectively).

10 Most respondents (61%) preferred to purchase leafy vegetables in fresh and on site due
11 to processing concerns (e.g., noncompliance with good hygiene and manufacturing practices)
12 and exposure to dust and flies which are diseases vectors. The high price of processed leafy
13 vegetables (1.20%) and the possibility of a mix with other leafy vegetables species (0.20%)
14 also contributed to reasons why respondent prefer fresh forms. Few households stated that
15 they would purchase processed vegetables even if they were available in pre-cooked and
16 packaged (5.40%); pre-cooked, packaged and frozen (1.80 %); packaged leaf powder (1.60
17 %); fresh, packaged and refrigerated (0.60 %) or dried and packaged (0.60 %).

18 Multivariate analysis on consumption frequencies and processing methods

19 The variability within consumption of Amaranth frequencies was related to socio-
20 cultural groups inversely to processing methods which were linked to age groups and
21 geographic areas. People from Bariba, Biali, Dendi, Ditamari, Fulani, Gourmantche, Lokpa,
22 Pedah, and Yom socio-cultural groups consumed Amaranth two to five times per week.
23 Inversely, those belonging to Adja, Ahoussa, Ani, Fon, Goun, Idatcha, Itcha, Kabiere, Lama,
24 Mahi, Mina, Nagot, Waama, Xlwa and Yoruba socio-cultural groups consumed Amaranth
25 'once per week'. People having more than fifty years old and living in southern part of the

26 country applied pre-cooking to Amaranth while those aged from eighteen to thirty years old
27 and living in northern part used direct cooking.

28 People belonging to Ahoussa, Bariba, Fulani, Idatcha, Mahi, Nagot, Waama and Yom
29 socio-cultural groups consume African basil rarely ('once or twice per month'). Inversely,
30 person from Adja, Biali, Ditamari, Dendi, Goun, Fon, Itcha, Kabiere, Lokpa, Mina, Pedah
31 Xlwa and Yoruba socio-cultural groups consume it 'once per week' and 'two or three times
32 per week'. Direct cooking and crushing were used by people from Ani socio-cultural group,
33 having more than fifty years old and living in north-eastern areas. Pre-cooking was performed
34 by people from Lama socio-cultural group aged from eighteen to thirty years old and who
35 lived in southern and north-western areas.

36 People from Adja, Ahoussa, Ani, Dendi, Fon, Goun, Idatcha, Itcha, Kabiere, Lama,
37 Mina, Nagot, Xlwa and Yoruba socio-cultural groups consume African eggplant 'once per
38 week' contrary to those from Bariba, Biali, Ditamari, Fulani, Gourmantche, Lokpa, Mahi,
39 Pedah, Waama and Yom socio-cultural groups who consumed it 'two or three times per
40 week'. People living in southern areas applied pre-cooking, regardless of their age inversely
41 to those who lived in northern areas and who performed direct cooking.

42 *3.5. Leafy vegetables preservation*

43 Leafy vegetables are highly perishable in fresh form and various preservation methods are
44 used (53.80 %) to avoid yellowing and deterioration. Short-term preservation methods,
45 including storage in baskets, exposition to dew, storage in clay pots and blanching followed
46 by soaking or refrigeration, are used for Amaranth, African basil and African eggplant, at
47 ambient temperature, to extend shelf life at two or three days. Storage in baskets (24.20%)
48 was the most popular method used by all socio-cultural groups as it allowed the extension of
49 shelf life by an average of two days. This method consists of storing vegetables in baskets
50 placed in a cool and dry place followed by the sprinkling of water (two or three times per day)

51 to prevent dehydration. People from Bariba, Ditamari, Fon, Idatcha, Nagot and Yoruba socio-
52 cultural groups preserved vegetables by spreading them on the roof or on the tufts of grass at
53 nightfall (9.60%) and removing them in the morning to keep them away from heat and
54 sunlight. This process was repeated for 3 days on average before significant deterioration
55 occurred. Storage in clay pots (1.20 %), which maintained coolness and humidity, also
56 extended shelf life for up to two days. Blanching followed by soaking in water (11.20 %) was
57 also used by Ditamari, Fon and Goun socio-cultural groups to increase preservation. Ordinary
58 sun-drying (4.00 %) was used for by Bariba, Dendi, Ditamari, Lokpa, Nagot and Yom for
59 African basil and African eggplant. Some socio-cultural groups (Ditamari and Lokpa, 5.00 %)
60 sun-dried blanched leaves and leaf powders were stored in a dry place for three to six months.
61 Many factors hindered proper storage including exposure to heat, excessive drying, and high
62 moisture. Thus, proper leaf washing, avoiding exposition to heat , regular water sprinkling on
63 fresh leaves, storage of dried leaves and leaf powders in a dry place were main precautions
64 took by respondents to extend availability and shelf life of these leafy vegetables.

65 **4. DISCUSSION**

66 The information collected during this research indicates beneficial properties related to use of
67 traditional leafy vegetables in household food systems. These findings confirm the findings of
68 others who have reported on the dietary and medicinal benefits of leafy vegetables (e.g.,
69 Chadaré *et al.* 2008, Dansi *et al.* 2008, Achigan-Dako *et al.* 2010, Mahapatra *et al.* 2010,
70 Rahmatullah *et al.* 2010, Dougnon *et al.* 2012, Nana *et al.* 2012, Sossa-Vihotogbé *et al.* 2012,
71 and Muhammad and Shinkafi 2014). Amaranth, African basil and African eggplant were the
72 most consumed by respondent; findings consistent with Achigan-Dako *et al.* (2010) who also
73 determined that Amaranth, African basil and African eggplant were amongst the top five of
74 frequently used leafy vegetables in all phyto-geographical zones of Republic of Benin.
75 However, the consumption of Amaranth, African basil and African eggplant does differ

76 according to age, cultural and geographical groups. Consumption of Amaranth, African basil
77 and African eggplant, in southern part of the country, was related to ethno-botanical
78 knowledge of socio-cultural groups (Vodouhè *et al.*, 2012a). Similar observations were made
79 by Chadaré *et al.* (2008) and Sossa-Vihotogbé *et al.* (2012) on consumption of other
80 traditional green leafy vegetables in northern areas of the country. Likewise, previous reports
81 demonstrated that dietary habits depended on interactions between foods availability, cultural
82 identity, geospatial environment and consumers' spending power (Chadaré *et al.*, 2008; Dansi
83 *et al.*, 2008; Achigan-Dako *et al.*, 2010; Vodouhè *et al.*, 2012a; Dinssa *et al.*, 2016).

84 Various reasons were mentioned to explain the consumption of the studied vegetables,
85 including dietary habits, taste, need to diversify sauces, nutrients and therapeutic benefits.
86 Traditional leafy vegetables were reported as highly nutritive foods containing vital
87 components of balanced and healthy diets such as vitamins and their precursors, polyphenols,
88 minerals, proteins and fibres (Makobo *et al.*, 2010; Ogbadoyi *et al.*, 2011; Oluwalana *et al.*,
89 2011; Sossa-Vihotogbé *et al.*, 2013). Richness of Amaranth, African basil and African
90 eggplant in vitamins were reported as well-known by Beninese who considered it a key
91 selection criterion of these vegetables (Vodouhè *et al.*, 2012a). Therapeutic properties of the
92 studied vegetables are widely recognized in folk medicine; for instance Amaranth species
93 were recommended to children, lactating mothers and people who suffered of constipation,
94 fever, bleeding, anemia or kidney problems (Grubben *et al.*, 2004; Vodouhè *et al.*, 2012a).
95 Uses of African basil by pregnant women and lactating mothers to facilitate childbirth, clear
96 waste after childbirth, stimulate milk secretion, treat postpartum infections and wound healing
97 were reported by Dansi *et al.* (2008) within Nagot socio-cultural group. The knowledge that
98 respondent had on potential health benefits related to consumption of Amaranth, African basil
99 and African eggplant could explained the high consumptions of these vegetables regardless
100 age, cultural and geographical groups. However, some households mentioned lack of

101 knowledge, organoleptic attributes and totemic considerations to justify the non-consumption
102 of Amaranth, African basil and African eggplant. Similar reasons were recorded for the non-
103 consumption of some traditional leafy vegetables (Batawila *et al.*, 2007; Chadaré *et al.*, 2008;
104 Dansi *et al.*, 2008; Sossa-Vihotogbé *et al.*, 2012).

105 The diversity of production sources (home gardens, gardening sites and fields) are used
106 to gain access to plant resources throughout the year. Leafy vegetables were reported to be
107 harvested from native vegetation, as well as cultivated fields, home gardens, and local
108 markets (Batawila *et al.*, 2007). Our findings revealed that many criteria were used for
109 selection of Amaranth, African basil and African eggplant among which availability, taste and
110 therapeutic properties were of paramount interest. Literature indicated that availability, the
111 ability to process, nutritional properties, and taste were mains criteria for selection of
112 Amaranth, African basil and African eggplant.

113 According to Dinssa *et al.* (2016), consumption of traditional leafy vegetables is
114 increasing in Africa as a result of consumers' awareness on their nutritional properties and
115 willingness to pay premium prices for quality food. This study lends further evidence to this
116 finding and highlights the relationship between selection criteria and age, cultural and
117 geographical groups of respondent. Consumption frequencies (key indicators of dietary
118 habits) of Amaranth, African basil and African eggplant varied from once to thrice per week
119 and could be easier and useful way to fully enjoy potential health benefits associated to these
120 leafy vegetables (Uusiku *et al.*, 2010) although variations were observed according to
121 vegetables species and socio-cultural groups in this study.

122 The investigated households used Amaranth, African basil and African eggplant in
123 various forms for sauces preparation with a preference for fresh form rather than dried form.
124 This preference was reported in the northern part of the country (Chadaré *et al.*, 2008; Sossa-
125 Vihotogbé *et al.*, 2012). The processing methods varied according to vegetable species, types

126 of sauces, age and geographical groups. Direct cooking and pre-cooking were the most used
127 methods and pre-cooking required the use of additives (potash or bicarbonate of soda) to
128 accelerate the leaves' softening. Amaranth leaves were reported by respondent to be easily
129 softened in salted boiling water; in agreement with report from Grubben *et al.* (2004) on
130 softening of Amaranth leaves after five to ten minutes in salted boiling water. However,
131 processing of Amaranth, African basil and African eggplant required some key unit
132 operations (sorting, washing and cooking) which must be achieved carefully and properly to
133 avoid digestive disorders. This survey showed the use of some additional ingredients used to
134 enhance the taste of leafy vegetables sauces; consistently with findings of Sossa-Vihotogbé *et*
135 *al.* (2012). Recipes used to process these vegetables in food systems were fairly homogeneous
136 and not numerous and that could be a limiting factor to benefit of their health-promoting
137 attributes. Indeed, processing methods are well-known to decrease nutritional properties of
138 raw leaf whether by the diffusion of water-soluble components in pre-cooking water or by the
139 destruction of thermo-labile and/or oxidizable substances (Babalola *et al.*, 2010; Avallone *et*
140 *al.*, 2007). The detrimental effects increase with the increase of water volume used,
141 temperature and duration of exposure (Babalola *et al.*, 2010; Avallone *et al.*, 2007) although
142 heat and water treatments were useful to reduce the incidence of health hazards caused by
143 anti-nutritive factors (hydrocyanic acid, oxalic acid, alkaloids, saponins, etc.) contained in raw
144 leaves (Udousoro *et al.*, 2013; Fabbri and Crosby, 2016).

145 Therefore, there is a need of diversification in forms used in food systems since limited
146 dietary diversity was reported as a major challenge and source high prevalence of malnutrition
147 and undernourishment in rural farming communities of sub-Saharan Africa (Afari-Sefa *et al.*,
148 2012). Steaming was presented as the best approach to maintaining the nutritional quality of
149 Amaranth, African basil and African eggplant (Vodouhè *et al.*, 2012b). Otherwise, the easy
150 perishability of leafy vegetables posed major challenges for their storage and various methods

151 were used by investigated households to avoid their yellowing and deterioration. Sun-drying,
152 as a popular preservation method used to face scarcity periods, can increase shelf life and alter
153 nutritional quality depending on drying conditions (Batawila *et al.*, 2007). Oven-drying at 60
154 °C have been demonstrated to allow higher retention of nutritional and sensory attributes
155 while reducing contamination risk (Oluwalana *et al.*, 2011). Blanching before sun-drying has
156 been suggested as a valuable option for leafy vegetables preservation without compromising
157 their quality (Makobo *et al.*, 2010; Ogbadoyi *et al.*, 2011). The slowing-down ability of cold
158 storage in metabolic processes (respiration and enzymatic activities) responsible of post-
159 harvest losses offered promising prospects to extension of leafy vegetables shelf life and
160 preservation of their nutritional quality (Acho *et al.*, 2015). The high perishability of
161 traditional leafy vegetables was related to their small-scale production in Africa mainly
162 intended to home consumption and local open markets (Dinssa *et al.*, 2016). Therefore, there
163 is a need to develop value-added foods deriving from leafy vegetables extracts since
164 perishability is the limiting factor for development of value chain of fresh leafy vegetables.

165 5. CONCLUSION

166 This study was conducted in eleven districts of southern and northern areas of Republic of
167 Benin and allowed to gather information on households' dietary habits concerning leaves of
168 Amaranth, African basil and African eggplant. Results obtained revealed that a diversity of
169 leafy vegetables was used in the investigated areas for culinary and medicinal purposes with
170 Amaranth, African basil and African eggplant as the most consumed green leafy vegetables.
171 These leafy vegetables were mainly selected based on their availability (Amaranth),
172 therapeutic properties (African basil) and taste (African eggplant) to be used alone or in
173 association with other leafy vegetables for consumption. The importance of these selection
174 criteria were dependent on age, cultural and geographical groups. The main supply mode of
175 Amaranth, African basil and African eggplant was purchasing at market. The importance of

176 these leafy vegetables in dietary habits of socio-cultural groups encountered were reflected by
177 variability in weekly consumption frequencies, processing methods and preservation
178 techniques used. An overall observation of relation between consumption frequencies,
179 processing methods, age, cultural and geographical groups of each leafy vegetable revealed
180 that consumption frequencies varied with socio-cultural groups while processing methods
181 were related to age groups and geographic areas. Regarding information collected, lack of
182 diversification within processing methods and consumption forms (other than sauces) are the
183 major challenges of contribution of Amaranth, African basil and African eggplant in
184 nutritional security of households in the investigated areas. Diversification of consumption
185 forms could be addressed by incorporating the extracts obtained from these leafy vegetables
186 to commonly consumed foods and beverages in order to enhance their nutrient profile without
187 compromising their quality and consumers' acceptability.

188 **Consent Disclaimer:**

189 As per international standard respondents' written consent has been collected and preserved by the
190 author(s).

191 **6. REFERENCES**

- 192 Achigan-Dako, E. G., M. W. Pasquini, F. Assogba- Komlan, S. N'danikou, H. Yédomonhan, A.
193 Dansi, and B. Ambrose-Oji. 2010. Traditional vegetables in Benin. *Institut National des*
194 *Recherches Agricoles du Bénin*. Imprimeries du CENAP. p. 282.
- 195 Acho, F. C., L. T. Zoue, and S. L. Niamke. 2015. Effect of refrigeration storage on nutritive and
196 antioxidant properties of five leafy vegetables consumed in Southern Côte d'Ivoire.
197 *Pakistan Journal of Nutrition*. 14(7):401-408. <https://doi.org/10.3923/pjn.2015.401.408>.
- 198 Adewale, O. B., A. Onasanya, O. A. Fadaka, H. Iwere, S. O. Anadozie, A. O. Osukoya, and I. I.
199 Olayide. 2014. In vitro antioxidant effect of aqueous extract of *Solanum macrocarpon*
200 leaves in rat liver and brain. *Oxidants and Antioxidants in Medical Sciences*. 3(3):225-29.
201 <http://dx.doi.org/10.5455/oams.161214.or079>.

202 Afari-Sefa, V., A. Tenkouano, C. O. Ojiewo, J. D. H. Keatinge, and J. d'A. Hughes. 2012.
203 Vegetable breeding in Africa: constraints, complexity and contributions toward achieving
204 food and nutritional security. *Food Security*. 4:115-127.
205 <http://dx.doi.org/10.1007/s12571011-0158-8>.

206 Akpo-Djenontin, D. O. O., V. B. Anihouvi, V. P. Vissoh, F. Gbaguidi, and M. Soumanou. 2016.
207 Processing, storage methods and quality attributes of spices and aromatic herbs in the local
208 merchandising chain in Benin. *African Journal of Agricultural Research*. 11(37):3537-
209 3547. <https://doi.org/10.5897/AJAR2016.11262>.

210 Avallone, S., S. Brault, C. Mouquet, and S. Trèche. 2007. Home-processing of the dishes
211 constituting the main sources of micronutrients in the diet of preschool children in rural
212 Burkina Faso. *International Journal of Food Sciences and Nutrition* 58:108-115.
213 <https://doi.org/10.1080/09637480601143320>.

214 Babalola, O.O., O.S. Tugbobo, and A.S. Daramola, 2010. Effect of processing on the vitamin
215 C content of seven Nigerian green leafy vegetables. *Advance Journal of Food Science*
216 *and Technology*. 2(6):303-305.

217 Batawila, K., S. Akpavi, K. Wala, M. Kanda, R. Vodouhè, K. Akpagana. 2007. Diversité et
218 gestion des légumes de cueillette au Togo. *African Journal of Food, Agriculture, Nutrition*
219 *and Development*. 7(3):55-68.

220 Chadaré, F. J., J. D. Hounhouigan, A. R. Linnemann, M. J. R. Nout, and M. A. J. S. van Boekel.
221 2008. Indigenous knowledge and processing of *Adansonia digitata L.* food products in
222 Benin. *Ecology of Food and Nutrition*. 47:338-362.
223 <https://doi.org/10.1080/03670240802003850>.

224 Dansi, A., A. Adjatin, R. Vodouhè, K. Adéoti, H. Adoukonou-Sagbadja, V. Faladé, H.
225 Yedomonhan, A. Akoègninou, et K. Akpagana. 2008. Biodiversité des légume-feuilles
226 traditionnels consommés au Bénin. *Bibliothèque Nationale, République du Bénin*. p.182.

227 Dinssa, F. F., P. Hanson, T. Dubois, A. Tenkouano, T. Stoilova, J. d'A. Hughes, and J.D.H.
228 Keatinge. 2016. AVRDC - The World Vegetable Center's women-oriented improvement
229 and development strategy for traditional African vegetables in sub-Saharan Africa.
230 *European Journal of Horticultural Sciences*. 81(2):91-105.
231 <http://dx.doi.org/10.17660/eJHS.2016/81.2.3>.

232 Dougnon, T. V., H. S. Bankolé, R. S. Johnson, J. R. Klotoé, G. D. Fernand, and G. F. Assogba.
233 2012. Phytochemical screening, nutritional and toxicological analyses of leaves and fruits
234 of *Solanum macrocarpon* Linn (*Solanaceae*) in Cotonou (Benin). *Food and Nutrition*
235 *Sciences*. 3:1595-1603. <https://doi.org/10.4236/fns.2012.311208>.

236 Fabbri, A. D. T., and G. A. Crosby. 2016. A review of the impact of preparation and cooking on
237 the nutritional quality of vegetables and legumes. *International Journal of Gastronomy*
238 *and Food Science*. 3:2-11. <https://doi.org/10.1016/j.ijgfs.2015.11.001>.

239 Grubben, G. J., O. A. Denton, R. H. Messiaen, R. H. Lemmens, and L. P. Oyen. 2004. Plant
240 Resources of Tropical Africa 2: Vegetables. *PROTA Foundation/CTA*, Netherlands
241 (Wageningen: Backhuys Publishers).

242 Mahapatra, S. K., S. P. Chakraborty, and S. Roy. 2010. Aqueous extract of *Ocimum gratissimum*
243 *Linn* and ascorbic acid ameliorate nicotine-induced cellular damage in murine peritoneal
244 macrophage. *Asian Pacific Journal of Tropical Medicine*. 3(10):775-782.
245 [http://dx.doi.org/10.1016/S1995-7645\(10\)60186-1](http://dx.doi.org/10.1016/S1995-7645(10)60186-1).

246 Makobo, N., M. Shoko, and T. Mtaita. 2010. Nutrient content of Amaranth (*Amaranthus*
247 *cruentus*) under different processing and preservation stages. *World Journal of*
248 *Agricultural Science*. 6(6):639-643.

249 Muhammad, S., and M. A. Shinkafi. 2014. Ethnobotanical survey of some medicinal important
250 leafy vegetables in north western Nigeria. *Journal of Medicinal Plants Research*. 8(1):6-8.
251 <https://doi.org/10.5897/JMPR07.031>.

252 Nana, F. W., A. Hilou, J. F. Millogo, and O. G. Nacoulma. 2012. Phytochemical composition,
253 antioxidant and xanthine oxidase inhibitory activities of *Amaranthus cruentus L.* and
254 *Amaranthus hybridus L.* extracts. *Pharmaceuticals.* 5(12):613-628.
255 <http://dx.doi.org/10.3390/ph5060613>.

256 Ogbadoyi, E. O., A. Musa, J. A. Oladiran, M. I. Ezenwa, and F. H. Akanya. 2011. Effect of
257 processing methods on some nutrients, antinutrients and toxic substances in *Amaranthus*
258 *cruentus*. *International Journal of Applied Biology and Pharmaceutical Technology.*
259 2(2):487-502.

260 Oluwalana, I. B., J. A. Ayo, M. A. Idowu, and S. A. Malomo. 2011. Effect of drying methods on
261 the physicochemical properties of Waterleaf (*Talinum triangulare*). *International Journal*
262 *of Biology and Chemical Sciences.* 5(3):880-889.
263 <http://dx.doi.org/10.4314/ijbcs.v5i3.72167>.

264 Rahmatullah, M., M. E. Hasan, M. A. Islam, M. T. Islam, F. I. Jahan, S. Seraj, A. R. Chowdhury,
265 F. Jamal, M. S. Islam, Z. U. M. Emdad Ullah Miajee, R. Jahan and M. H. Chowdhury.
266 2010. A survey on medicinal plants used by the folk medicinal practitioners in three
267 villages of Panchagarh and Thakurgaon district, Bangladesh. *American Eurasian Journal*
268 *of Sustainable Agriculture.* 4:291-301.

269 Segnon, A. C., and E. G. Achigan-Dako. 2014. Comparative analysis of diversity and utilization
270 of edible plants in arid and semi-arid areas in Benin. *Journal of Ethnobiology and*
271 *Ethnomedicine.* 10(80):1-20. <https://doi.org/10.1186/1746-4269-10-80>.

272 Sossa-Vihotogbé, C. N. A., N. H. Akissoe, B. V. Anihouvi, G. L. Amadji, and J. D.
273 Hounhouigan. 2013. Effect of organic and mineral fertilization on nutritive value of three
274 leafy vegetables harvested at different periods. *International Journal of Biological and*
275 *Chemical Sciences.* 7(1):271-286. <http://dx.doi.org/10.4314/ijbcs.v7i1.23>.

- 276 Sossa-Vihotogbé, C. N., N. H. Akissoe, B. V. Anihouvi, B. C. Ahohuendo, A. Ahanchede, A.
277 Sanni, and J. D. Hounhouigan, 2012. Endogenous knowledge of four leafy vegetables used
278 by rural populations in Benin. *Ecology of Food and Nutrition*. 51:22-36.
279 <https://doi.org/10.1080/03670244.2012.635570>.
- 280 Tchiegang, C. and A. Kitiki. 2004. Données ethnonutritionnelles et caractéristiques physico-
281 chimiques des légumes-feuilles consommés dans la savane de l'Adamaoua (Cameroun).
282 *Tropicultura*. 22 (1):11-18.
- 283 Udousoro, I. I., R. U. Ekop, E. J. Udo. 2013. Effect of thermal processing on antinutrients in
284 common edible green leafy vegetables grown in Ikot Abasi, Nigeria. *Pakistan Journal of*
285 *Nutrition*. 12(2):162-67. <https://doi.org/10.3923/pjn.2013.162.167>.
- 286 Uusiku, P. N., A. Oelofsenc, K. G. Duodu, M. J. Bester, and M. Faber. 2010. Nutritional value of
287 leafy vegetables of sub-Saharan Africa and their potential contribution to human health: A
288 review. *Journal of Food Composition and Analysis*. 23:499-509.
289 <https://doi.org/10.1016/j.jfca.2010.05.002>.
- 290 Visweswari, G., R. Christopher, and W. Rajendra. 2013. Phytochemical screening of active
291 secondary metabolites present in *Withania somnifera* root: role in traditional medicine.
292 *International Journal of Pharmaceutical Sciences and Research*. 4(7):2770-2776.
293 <http://dx.doi.org/10.13040/IJPSR.0975-8232>. 4(7).2770-76.
- 294 Vodouhè, S., R. Tossou, et M. Soumanou. 2012a. Perception des consommateurs sur la qualité
295 nutritionnelle et sanitaire de quelques légumes feuilles locaux produits dans la zone côtière
296 du Sud Bénin. *Bulletin de la Recherche Agronomique du Bénin*. 1-11.
- 297 Vodouhè S, A, Dovoedo V. B., Anihouvi and M. M. Soumanou. 2012b. Influence of cooking
298 mode on the nutritional value of *Solanum macrocarpum*, *Amaranthus hybridus* and
299 *Ocimum gratissimum*, three traditional leafy vegetables acclimated in Benin. *International*

300 *Journal of Biological and Chemical Sciences.* 6(5):1926-37.
301 <http://dx.doi.org/10.4314/ijbcs.v6i5.3>.

302 Vorster, H.J., W. Jansen Van Rensburg, J. J. Van, B. Ziji, and V. C. Sonja. 2007. The importance
303 of traditional leafy vegetables in South Africa. *African Journal of Food, Agriculture,*
304 *Nutrition and Development.* 7(4):1-13.

305 Yabi I, and F. Afouda. 2012. Extreme rainfall years in Benin (West Africa). *Quaternary*
306 *International.* 262(7):39-43. <https://doi.org/10.1016/j.quaint.2010.12.010>.

307 ACKNOWLEDGEMENT

308 The study was funded by the International Development Research Centre (IDRC) and Global
309 Affairs Canada, through the Canadian International Food Security Research Fund (CIFSRF)
310 Project No. 107983. The author would like to thank Modoukpè I. DJIBRIL MOUSSA, Carole
311 N. A. SOSSA-VIHOTOGBE, Irenikatché P. B. AKPONIKPE and André J. DJENONTIN for
312 their contribution to this research.

313

315 Table 1: List of villages surveyed and their socio-cultural groups

Zones	Districts	Investigated areas	Cultural groups
South	Abomey-Calavi	Abomey-Calavi, Akassato, Godomey, Glo-djigbe, Hevie, Kpanroun, Ouedo, Togba, Zinvie	Fon
	Cotonou	Aibatin 2, Aidjedo, , Ayelawadje, Cadjehoun, Dandji, Fifadji, Gbedokpo, Gbgamey, Kouhounou, Saint Michel Sainte Rita, Sodjatime, Yenawa,	Fon
	Seme-kpodji	Agbangandan, Aholouyeme, Djeregbe, Ekpe, Seme-kpodji, Tohoue	Goun
	Bohicon	Gbangnikon, Hezonho, Houawe ouianssa, Hountonho, Sodohome, Sogba	Fon
	Djidja	Agbohoutogon, Bookou, Djidja aglomey, Djidja centre	Fon
North-east	Boukoumbe	Koudadagou, Koudogou, Koukoua, Kounacogou, koutagou	Ditamari
	Djougou	Kilir, Taefa, Timba	Dendi, Yom
	Ouake	Itchode, Tchalade, Tchalinga	Lokpa
North-west	N'Dali	Boko, Gomez Kaprou, Mareborou, Suanin, Wobakarou,	Bariba
	Parakou	Kabounari, Sokounon, Titirou, Tourou, Zongo II	Bariba, Dendi
	Tchaourou	Kaki koka, Kassouala, Kika, Kpassa, Tchatchou	Nagot

Table 2: Dietary uses of traditional leafy vegetables consumed in study areas

Scientific names (English names)	Local names ¹	Dietary utilization		
		Form of used	Processing methods	Beneficial effects
<i>Abelmoschus esculentus</i> (Okra)	Bariba: Kôbusa, Ditamari: Tinoufanti, Fon: Féviman, Lokpa: Maatou, Nagot: Ewéila.	Fresh leaves, dried leaves or dried leaf powders	Cooking in boiling water containing softening ingredient (potash or kanmu: alkaline rock salt) to get a slimy soup.	Very nutritious, stimulate appetite and manage digestives disorders.
<i>Adansonia digitata</i> (African baobab)	Bariba: Sônnawourousou, Ditamari: Titonankanti, Fon: Azizonman, Lokpa: Kôtôlassa, Nagot: Ewéotché.	Fresh leaves, dried leaves or dried leaf powders	Cooking in boiling water containing kanmu or potash to get a slimy soup.	Very nutritious, manage digestives disorders.
<i>Amaranthus cruentus</i> (Amaranth)	Bariba: Affônou, Ditamari: Aléfô, Fon: Fotètè, Lokpa: Aléfô, Nagot: Effô tètè.	Fresh leaves or pre-cooked (in salted boiling water)	Raw leaves are used as an ingredient of salad, addition of raw or pre-cooked form into tomato or onion-based sauce to get a vegetable soup.	Very nutritious, stimulate appetite and manage digestives disorders.
<i>Celosia argentea</i> (Cockscomb)	Bariba: Affônou, Ditamari: Tinonyawoti, Fon: Sôman, Nagot: Odjogodo.	Fresh leaves or pre-cooked (in salted boiling water)	Addition into tomato or onion-based sauce to get a vegetable soup.	Very nutritious.
<i>Ceratotheca sesamoides</i> (False sesame)	Bariba: Kpééwori, Ditamari: Siwadouanwe, Fon: Agbô, Lokpa: Hounoume, Nagot: Idjabô.	Fresh leaves or dried leaf powders	Addition into boiling water containing kanmu or potash to get a slimy soup.	Very nutritious, manage digestives disorders.
<i>Corchorus olitorius</i> (Jute mallow)	Bariba: Yôyôkou, Ditamari: Tifanwounti, Fon: Ninnouwi, Lokpa: Ayôyôkou, Nagot: Owoyô.	Fresh leaves	Addition into boiling water containing kanmu or potash to get a slimy soup.	Very nutritious, restore appetite and manage digestives disorders.
<i>Hibiscus sabdariffa</i> (Roselle)	Bariba: Sinri, Ditamari: Tikouan n'ti, Fon: Sinku, Lokpa: Ankpaman, Nagot: Kpakpa.	Pre-cooked leaves (in boiling water containing kanmu or potash)	Addition into tomato-based sauce to get a vegetable soup.	Very nutritious, manage digestives disorders.

Scientific names (English names)	Local names ¹	Dietary utilization		
		Form of used	Processing methods	Beneficial effects
<i>Justicia tenella</i>	Bariba: Kourôkountônu, Ditamari: Tinoukounti, Lokpa: Tiletoussi, Nagot: Djagoudjagou.	Fresh leaves or pre-cooked (in boiling water containing kanmu or potash)	Addition into tomato-based sauce to get a vegetable soup.	Very nutritious
<i>Manihot esculenta</i> (Cassava)	Bariba: Kpakiwourousou, Ditamari: Tikôkônouwôti, Fon: Finyinman, Lokpa: Agbedehatou, Nagot: Ewékpakî.	Pre-cooked leaves (in boiling water containing kanmu or potash)	Addition into tomato-based sauce to get a vegetable soup.	Very nutritious.
<i>Moringa oleifera</i> (Drum-stick tree)	Bariba: Yorouyara, Ditamari: Mounpèkom, Fon: Kpatiman, Lokpa: Lôtaha, Nagot: Lagalanga.	Fresh leaves or dried leaf powders	Fresh leaves are used to make teas and as an ingredient of salad and vegetable soup. Dried leaf powders are used as diet supplements.	Very nutritious, stimulate appetite and manage digestives disorders.
<i>Ocimum gratissimum</i> (African basil)	Bariba: Danbakarou, Ditamari: Tinassiyinti, Fon: Tchiayo, Lokpa: Assôhou, Nagot: Simonba.	Fresh leaves, crushed or pre-cooked (in boiling water containing kanmu or potash)	Raw leaves are used to make teas and as a seasoning, addition of crushed or pre-cooked leaves into tomato or onion-based sauce to get a vegetable soup.	Very nutritious, manage digestives disorders.
<i>Sesamum radiatum</i> (Black benniseed)	Bariba: Dossi, Ditamari: Tiwadounati, Fon: Akanmanku, Lokpa: Touhoonôme, Nagot: Dossé, Goolowo.	Fresh leaves or dried leaf powders	Cooking in boiling water containing kanmu or potash to get a slimy soup.	Very nutritious, manage digestives disorders.
<i>Solanum macrocarpon</i> (African eggplant)	Bariba: Sambinouwourousou, Ditamari: Tikawounfanti, Fon: Gboman, Lokpa: Mètècou, Nagot: Katakoukpakou.	Pre-cooked leaves (in boiling water containing kanmu or potash)	Addition into tomato or onion-based sauce to get a vegetable soup.	Very nutritious, manage digestives disorders.

Scientific names (English names)	Local names ¹	Dietary utilization		
		Form of used	Processing methods	Beneficial effects
<i>Talinum triangulare</i> (Water leaf)	Bariba: Odôndôn, Ditamari: Yêmontouo, Fon: Aglassoeman, Lokpa: Kamplékankann'dê, Nagot: Odôndôn.	Fresh leaves or pre-cooked (in boiling water containing kanmu or potash) and dried leaves	Addition of pre-cooked leaves or dried leaves into tomato-based sauce to get a vegetable soup.	Very nutritious, manage digestives disorders.
<i>Vernonia amygdalina</i> (Bitter leaf)	Bariba: Touwan, Ditamari: Souwaka, Fon: Amanvivè, Lokpa: Souwaka, Nagot: Eéwoh.	Pre-cooked leaves (in boiling water containing kanmu or potash)	Addition into tomato-based sauce to get a vegetable soup.	Very nutritious, manage digestives disorders.
<i>Vigna unguiculata</i> (Cowpea)	Bariba: Suiwourousou, Ditamari: Titouti, Fon: Ayiman, Lokpa: Tchaasé, Nagot: Oson.	Pre-cooked leaves (in boiling water containing kanmu or potash)	Addition into tomato-based sauce to get a vegetable soup.	Very nutritious, stimulate appetite and manage digestives disorders.
<i>Vitex doniana</i> (Black plum)	Bariba: Kounonkou, Ditamari: Timantoun n'ti, Fon: Fonman, Nagot: Ewa.	Fresh leaves or pre-cooked (in boiling water containing kanmu or potash)	Fresh leaves are used to make teas, addition of pre-cooked form into tomato-based sauce to get a vegetable soup.	Very nutritious, manage digestives disorders.

Sources: survey in October 2015 and December 2016 and additional information from Dansi *et al.* (2008) and Achigan-Dako *et al.* (2010)

¹: choice of local languages is related to the main cultural groups encountered [Bariba (17.40 %), Ditamari (11.60 %), Fon (19.40 %), Lokpa (9.20 %) and Nagot (9.60 %)].

Table 3: Various types of sauces obtained from each leafy vegetable

Leaves form	Species	Additive/Ingredient	Unit operations	Sauces
Fresh	Amaranth, African basil, African eggplant	Tomato, onion, vegetable oil, seasoning	Sorting/clipping/washing /adding to sauce/cooking	Simple sauce 1
Dried and powders from dried leaf	Amaranth, African basil, African eggplant	Tomato, onion, vegetable oil, seasoning	Sorting (for dried leaf)/washing (for dried leaf) / adding to sauce/cooking	Simple sauce 2
Fresh	Amaranth, African basil, African eggplant	Potash Tomato, onion, vegetable oil, seasoning	Sorting/clipping/washing /pre-cooking/draining/ adding to sauce/cooking	Simple sauce 3
Fresh and crushed	African basil	Onion, vegetable oil, seasoning	Sorting/washing/crushing /adding to sauce/ cooking	Simple sauce 4

UNDER PEER REVIEW

LIST OF FIGURES




Figure 1: Benin map showing the geographical location of the survey sites