

Case report

A case report on Twin Babies , One Born with Gastroschisis & the Other with Sirenomelia in Iraq

Abstract

Back ground: Both Sirenomelia & Gastroschisis are very rare congenital anomalies & it is extremely rare to find each of the two anomalies in both twins of one pregnancy ,here we reported the two anomalies in one twin pregnancy.

Case report: we reported twin pregnancy in G3p0A2 19 years old healthy woman who had irregular antenatal visits , ultrasound exam done 4 times during pregnancy , all shown monochorionic, diamniotic twins with polyhydromnia & Gastroschisis , abnormal spine and sacrococcygeal teratoma in the 1st twin, and oligohydromnia in the 2nd twin in which the anomaly was not clear & later on the baby born with Sirenomelia.

In all the references we reviewed, we couldn't find the two anomalies in one twin pregnancy and this case report was considered to be the 1st reported case globally.

Key words: Twin, Sirenomelia, Gastroschisis, Fallujah.

Introduction

Gastroschisis is a birth defect of the abdominal wall which can be small or large through which the intestine and some times other organs such as stomach & liver protrude outside the baby's body. It occurs early during pregnancy when the muscles that make up the baby's abdominal wall do not form correctly. Because the intestines are not covered by a protective sac & are exposed to the amniotic fluid , the intestines can become irritated causing them to shorten , twist or swell¹ . The incidence is 0.4-3/10.000 live¹ ,

It's etiology is unknown in most cases , some studies suggested some risk factors for its occurrence like young age (teenage) mothers, Alcohol & tobacco smoking during pregnancy. it can be diagnosed early during pregnancy by ultrasonography¹ .

Sirenomelia , also called Mermaid syndrome, is a rare congenital anomaly of uncertain etiology , characterized by fusion of the lower limbs & commonly associated with severe urogenital & gastrointestinal malformation, it should be suspected in antenatal cases presenting with severe oligohydromnias & IUGR² . The incidence is about 1.5-4.2/100.000 births , it is 100-150 times more likely in identical (monozygotic twins) than in singletons or fraternal twins³ .

In our hospital we previously reported 3 cases of unusual anterior abdominal wall defect in an article published at 2012 ¹ , And 2 cases of Sirenomelia also published at 2013 ² , but it was the 1st time to document the 2 conditions in 2 twins babies in the same pregnancy , & it is the 1st reported case globally .

Case presentation:

The mother was 19 years old, previously healthy, G3p0A2, of medium socio-economic status , non-consanguineous marriage. No family history of any kind of congenital anomalies from both paternal sides. She gave history of 2 previous miscarriages (at 6 & 8 wks successively) due to unknown cause about 1y before the last pregnancy . She had negative antenatal serology, not diabetic nor hypertensive, and gave no history of having smoked or drank alcohol which is not allowed in our culture , There was no history of exposure to toxins or drugs or any significant febrile episode in the antenatal period. She had irregular antenatal visits, ultrasound exam done 4 times during pregnancy, all shown monochorionic, diamniotic twins with polyhydromnia & Gastroschisis, abnormal spine and sacroccygeal teratoma in the 1st twin , and oligohydromnia in the 2nd twin in which the anomaly was not clear and the mother was told that the condition of the 2nd twin was suspicious , that is why it was difficult to take the decision for the termination of pregnancy .

EDD was 10/7/2019 .The last antenatal Us has done on 30/3/2019 in privet Obstetric Us clinic & it was decided that the 2nd twin looks normal (Figure 1& figure 2).

On 30/4/2019 , the mother was at her 32nd wk of pregnancy presented to our hospital with labor pain and was admitted to the labor room , the twins were delivered by spontaneous vaginal delivery , the 1st born alive, weighing 2200 gm , with fused lower extremities, absent genitalia and absent anal and urethral orifices with appearance of single foot containing 3 abnormal separated toes , the babie's face , trunk and upper limbs were apparently looking normal , not dyspneic or cyanosed , investigation including skeletal x.ray exam and abdominal and pelvic sonography were requested but the father refused any interference and discharged the baby on his own (Figure 3,4,5 &6) .



Figure 1 (2nd twin looks normal)




Figure 2(2nd twin looks normal)



Figure 3 (1st twin born with Sirenomelia)

,the 2nd with Gastroschisis



Figure 4(1st twin with fused lower limbs &absent genitalia)



Figure 5(Head ,face & trunk look normal)



Figure 6(appearance of single foot containing 3 abnormal separated toes)

The 2nd twin was male, weighing 2000gm, born dead (stillbirth) with a big defect (about 3x4cm) on the right para-umbilical area with the liver, stomach ,small & large bowel herniated outside his body , distorted spine with big sacrococcygeal swelling shown by the prenatal ultrasound as spina bifida & meningocele , and abnormal right knee with the leg & foot rotated upward in addition to congenital heart disease (Figure 7, 8 & 9) . In our patient (the 1st twin) , the associated anomalies were complete agenesis of urogenital system and imperforated anus, Ultrasonography and skeletal radiology to check for other abdominal or pelvic anomalies were not done because the baby's father refused any interference and discharged the baby from the hospital on his own responsibility.



Figure 7(liver, stomach & bowel herniated)



through abdominal wall defect)

Figure 8(Gastroschisis & deformed lower limbs)



Figure 9 (spina bifida& sacrococcygeal swelling)

Discussion and conclusion :

Sirenomelia is a rare congenital anomaly characterized by partial or complete fusion of lower limbs and usually associated with other severe anomalies. It is considered, by many authors, as severe form of caudal regression syndrome. The associated anomalies may include bilateral renal agenesis, complete or partial agenesis of genitourinary system, imperforate anus, absence or ambiguous external genitalia, single umbilical artery, lung hypoplasia and vertebral and cardiac anomalies⁽⁴⁾.

Etiology of sirenomelia is uncertain and various theories have been proposed to explain its origin. An embryonic insult to caudal mesoderm between 28-32 days of gestation and vascular hypo-perfusion has been proposed as possible factors. Others associated the condition with maternal diabetes mellitus, exposure to teratogens and genetic predisposition Sirenomelia carries a grim prognosis, with survival dependent on the nature of the visceral anomalies⁵.

Abdominal wall defects can be identified prenatally via ultrasound or measurement of alpha-fetoprotein levels (AFP) within the amniotic fluid, Alpha-fetoprotein is the fetal analog of albumin, and maternal serum AFP (MSAFP) reflects the level of AFP in the amniotic fluid⁽⁶⁾.

In gastroschisis, MSAFP is markedly abnormal; up to 9 times the median value^(2&6). Gastroschisis is a right paraumbilical defect involving all layers of the abdominal wall⁶. The small bowel always eviscerates through the defect and is, by definition, non-rotated and lacking secondary fixation to the posterior abdominal wall⁽⁷⁾. The loops of bowel are never covered by a membrane; hence, they are directly exposed to the amniotic fluid^(2,8,9,10).

In our patient in addition to the presence of big abdominal defect with herniation of most of the visera there was also spina bifida and meningocele in addition to the abnormal right knee with the leg & foot rotated upward.

From this and all our other previous reports and studies during the last 11 years, we conclude that the high prevalence of birth defects in Fallujah is impairing the population's health and its capacity to care for the surviving children especially in the absence of sufficient facilities for the prenatal and postnatal management of cases of birth defects and also the absence of clear serious future plans for improvement of

the health system or taking serious measures to clean the post wars contaminated environment .

Patient's consent :

Consent for publication was obtained by the patient's family

Conflict of interest :

The author has no conflict of interest .

References :

1. Alaani S,MD , paediatrician , Fallujah maternity & children hospital, Report of Three Cases of Unusual Anterior Abdominal Wall Defects in Fallujah General Hospital During the Year 2010; JIMA: Vol 44, 2012, DOI: <http://dx.doi.org/10.5915/44-1-10354>
2. Alaani S, MD , paediatrician , **Murie A,MD**, Clinical geneticist, Fallujah maternity & children hospital,Fallujah city ,Anbar province , Iraq, **Sirenomelia within a one-year period at Fallujah general hospital: two case reports , Science Journal of Clinical Medicine** 2013; 2(3) : 64-67, doi: 10.11648/j.sjcm.20130203.11
3. Onyeije, Chukwuma; Sherer, David; Handwerker, Sara; Shah, Leena (2008). "Prenatal Diagnosis of Sirenomelia with Bilateral Hydrocephalus: Report of a Previously Undocumented form of VACTERL-H Association". *American Journal of Perinatology*. **15** (3): 193–7. doi:10.1055/s-2007-993925. PMID 9572377 .
4. Gouri MSR, Kumari KMK, Rao SG, Devi U. Sirenomelia. *Ind J PatholMicrobiol* 2009; 52:579-80.
5. Van KJ, Cannie M, Robrechts C, et al. First trimester diagnosis of Sirenomelia.*PrenatDiagn* 2006; 26(8):684-88.
6. Holmgren G, Sigurd J. Prenatal diagnosis of two cases of gastroschisis following α -fetoprotein (AFP) screening. *Acta Obstet Gynecol Scand*. 1984;63:325-8. <http://doi.org/b3fsp5> .
7. Redford DH, McNay MB, Whittle MJ. Gastroschisis and exomphalos: precise diagnosis by midpregnancy ultrasound. *Br J Obstet Gynaecol*. 1985;92:54-9. <http://doi.org/cvk69h> .
8. Bair JH, Russ PD, Pretorius DH, et al. Fetal and gastroschisis: a review of 24cases. *AJR Am J Roentgenol*. 1986;147:1047-51. [PubMed PMID: 2945411](http://pubmed.ncbi.nlm.nih.gov/2945411/) .
- 9 .Emami CN, Youssef F, Baird RJ, Laberge JM, Skarsgard ED, Puligandla PS. A risk-stratified comparison of fascial versus flap closure techniques on the early outcomes of infants with gastroschisis. *J Pediatr Surg*. 2015;50(1):102-6. <http://doi.org/f6wfxc>. [[Links](#)]
- 10 . Chesley PM, Ledbetter DJ, Meehan JJ, Oron AP, Javid PJ. Contemporary trends in the use of primary repair for gastroschisis in surgical infants. *Am J Surg*. 2015;209(5):901-5. <http://doi.org/f7bj56>. [[Links](#)]