

**Dynamics of sediment organic matter along Bandama River in the department of Sinématiali, northern Côte d'Ivoire**

**ABSTRACT**

This study examines the distribution of organic matter in areas affected by frequent floods along the east bank of the Bandama River in the department of Sinématiali. The sites sampled are defined by two zones, one near the stream and one far from the stream. Samples collected were analyzed, including for texture with aggregation analysis by the Robinson pipette, and standard sediment analysis methods for measuring organic carbon (CO), nitrogen (N), and organic matter (MO). Statistical analyzes were carried out to assess the differences between the physico-chemical parameters of the different sampling areas. Results show that sediment from the various study sites has a sando-limonous to limono-clay texture. Total organic matter levels are higher in surface sediments that contain the lowest proportions of clay. Rates range from 31.98gkg<sup>-1</sup> to 38.98gkg<sup>-1</sup>. In depth, the rates obtained are very low and range from 6.3gkg<sup>-1</sup> to 8.19 3gkg<sup>-1</sup>. The low rates recorded in depth are reported to be related to leaching caused by periodic flooding. These results show that successive floods have a direct effect on the dynamics of the physico-chemical properties of the sediments along the shore.

**Keywords:** sediment, shore, organic matter, particle size, flooding

**1. INTRODUCTION**

During the past decade, the number of studies on the effects of climate change on the environment has increased steadily. One of the potential impacts of climate change on the water cycle is an increase in the number and intensity of floods [1,2,3]. At present, it is important to quantify the impact of climate change on the river environment in order to better understand how these environments will evolve in the coming decades. Riparian ecosystems vary widely depending on river dynamics [4]. The constant supply of sediment transported by successive floods has an impact on the physico-chemical properties of soils. The latter can vary greatly in vertical distribution and spatial distribution, which are affected by various hydrological processes [5,6]. Sediment is a relatively heterogeneous matrix consisting of water, inorganic and organic materials and anthropogenic compounds. It can be described by its composition and structure [7] (Power and Chapman. Surface sediments are among the most important storage and dynamic reservoirs of organic matter [8]. Each year, nearly 0,4 Gt of terrestrial organic carbon is transported to coastal environments [9]. MO consists of a large number of chemical species that may be of size, chemical composition, and complex and varied physical forms [10]. These different parameters will depend primarily on the origin but also on the environment in which the MO will be located [11]. MO is a complex substance consisting primarily of carbon and hydrogen and in variable proportions of oxygen, nitrogen, sulfur and phosphorus [11]. Nitrogen and organic carbon play very important roles in MO biogeochemistry [12] particularly in microbiological activity [13] and metal complexation processes [14]. The study of sedimentary organic matter in current environments could be used as a basis for paleoclimatic, and paleoenvironmental interpretations [15]. Previous work indicates little information on the distribution of these physico-chemical parameters in sediments along Bandama River shore. As a result, this study will attempt to highlight the impact of recurring flooding on sediment organic matter in this area. The aim is to assess the levels of organic matter in sediments away from the stream and compare them with those in the surrounding environment in order to highlight the impact of the floods on the river environment. No studies were conducted on sediment organic matter along the rivers of this region. The main purpose of this study is to address this shortcoming. To carry out this work, a multidisciplinary approach has been taken. It is based in particular on sedimentological and pedological studies.

56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71

## 2. MATERIAL AND METHODS

### 2.1 Location and sampling

Four sampling points along the shore were predefined (figure 1) and were sampled from surface sediments at a depth of 10 to 150 cm. The various analyzes carried out in this study were carried out on the fine length part (< 2 mm), which we separated at the laboratory a few weeks later after sampling.

These samples were analyzed to determine:

- The particle size (sandy, clay and lemon fraction), pH, total organic carbon, total nitrogen and organic matter.
- The analysis of each of these parameters involved specific methods as described below.




Figure 1: Location map

72  
73  
74  
75  
76  
77  
78  
79  
80

## 2.2 Physico-chemical characterization of sediment

### 2.2.1 Physical Parameters of Sediments

- Granulometry

81  
82 It was determined by Robinson's Pipette method [16].

83 The application of this analysis allowed:

- 84 - to know the substances (MO and nitrogen) associated with the particle size contained in the
- 85 sediment. It is used to determine whether they are in the fine, medium or coarse fractions;
- 86 - to reconstitute the conditions for transport and deposition of particles.

87  
88 

- **Texture**

89 We used the triangular diagram of fine soils proposed by [17]. This type of diagram is particularly  
90 suitable for sediments because sediments can then be characterized according to the respective  
91 content of these three particle size fractions (clays, silt and sand) [18,19].

92  
93 **2.2.2 Chemical sediment parameters**

94  
95 On the same samples we also determined CO, MO, nitrogen and PH levels.

96  
97 **pH**

98 pH of water was determined by measuring H<sub>3</sub>O<sup>+</sup> ion activity using a pHmeter [20].  
99

100 **Organic carbon**

101 Organic carbon of sediments is determined by Anne's method [21]. The carbon of the organic matter is  
102 oxidized by potassium bichromate in sulfuric medium.

- 103  
104 • Organic matter

105  
106 The assessment of the organic matter content was made by quantifying its major constituent element,  
107 organic carbon, which represents almost 50% of this element [22]. The content of the MO was  
108 assessed based on the following conventional relationship: MO= C×1,72 [23].

109  
110 

- **Nitrogen**

111 Nitrogen in sediments was determined by the Kjeldahl method [20]. The principle of this method is to  
112 transform the nitrogen of organic compounds from a finely crushed sediment sample into ammoniacal  
113 nitrogen under the action of concentrated sulfuric acid, which, when boiled, behaves as an oxidant.  
114 Organic substances are decomposed: carbon comes out as carbon dioxide, hydrogen gives water,  
115 and nitrogen is transformed into ammonia nitrogen. The latter is fixed immediately by sulfuric acid in  
116 the form of ammonium sulfate.

- 117  
118 • **Carbon-to-Nitrogen Ratio (C/N)**

119  
120 The C/N ratio is an indicator of sediment biological activity that provides information on the degree of  
121 organic matter evolution, biological activity, mineralization. The smaller the biological activity, the more  
122 difficulties encountered in mineralization. This reflects conditions of anaerobic, excessive acidity. The  
123 study of the C/N report is an approach to the problem of the origin, nature and evolution of organic  
124 matter [24].


125  
126  
127  
128 **2.2.3 Statistical analysis**

129  
130 For data analysis, we used variance analysis of different variables using SAS 9.4 software. The  
131 significance test is Fischer's distribution or "F test" at the 5% probability threshold. Correlation tests  
132 (Pearson) were also performed between variables (MO, pH, C.O., N, texture). These tests allowed  
133 comparisons of different parameters according to the horizontal gradient (channel distance) and  
134 vertical (depth). Finally, a primary component (PCA) analysis was conducted to verify that there is a  
135 link between the different physico-chemical parameters, the layers and the positions close to or far  
136 from the stream. We converted the units of % to gkg<sup>-1</sup> (international unit) for carbon, nitrogen and  
137 organic matter (MO).


138 **3. RESULTS**

139  
140 **3.1 Sediment Granulometry and Texture**

141  
142 The clay fraction increases with depth in the profile near the stream bed (Figure 2). The remote profile  
143 of the source has similar characteristics. Limon percentages decrease with depth near the channel  
144 while sand percentages increase. The sandy fraction is greater when you move away from the  
145 channel. Sediments are coarser by moving away from the stream (Figures 2 and 3). Sediments near  
146 the river are characterized by higher percentages in silt. The results of our analyzes reveal a limono-  
147 clay texture near the channel and a sandy-clay texture as one moves away from the channel (Table  
148 1).  
149



150  
151 **Figure 2** : particle size composition  
152 F10 = pit near the river F40 = remote pit  
153  
154



155  
156 **Figure 3** : Granulometry relative to profile positioning  
157

158  
159  
160  
161  
162  
163

Table 1: Granulometry and sediment texture of layers according to positions

Layer	F10 <sub>1</sub>	F10 <sub>2</sub>	F10 <sub>3</sub>	F40 <sub>1</sub>	F40 <sub>2</sub>	F40 <sub>3</sub>
-------	------------------	------------------	------------------	------------------	------------------	------------------

Coarse sand (p.c.)	5,55	5,25	24,95	16,95	17,55	14,75
Fine sand (p.c.)	17,2	13,15	7,65	37,7	34,1	30,1
Coarse Limon (p.c.)	42,65	44,2	24,2	16,35	12,55	16,15
Fine Limon (p.c.)	13,5	9	11	10,5	8,25	7,75
Clay (p.c.)	19,25	26	31	17	26,5	29,25
Texture	LS	LA	LA	SL	SA	SA

164

165

### 166 3.2 Chemical parameters

167

#### 168 3.2.1 pH

169

170 The pH of the sediments relative to their stream position. Sediments have low acidity pH ranging from  
 171 5.72 to 5.79 (Figure 4). This low acidity could be explained by the absence of abundant surface litter in  
 172 this area. Decomposition of litter and humus releases acidifying products such as fluvial and humic  
 173 acids that acidify surface sediments.

174


175

#### 176 Change in pH in different layers

177

178 A change in the pH value from the surface to the depth is noted (Figure 5). Biomass is often almost  
 179 non-existent in depth and therefore very little input from acidifying products. This could explain the  
 180 high pH value in the depth layers.

181




182

183 **Figure 4 : Change in pH by position**

184

185



186  
187 **Figure 5** : Variation in layers


188  
189  
190 **3.2.2 Organic Matter**

191  
192  
193 **Change of MO in pits to position**


194  
195 Organic matter does not vary significantly from pit position (Figure 6). These values indicate relatively  
196 lower rates in areas far from the stream. Concentrations ranging from 21.01 to 18.21 (gkg<sup>-1</sup>) indicate  
197 medium-rich organic sediment [25].

198  
199  
200 **Change in MO in layers**

201  
202 Analyzes show a sharp decrease in organic matter content when going deep (Figures 6 and 7). The  
203 MO rate increases from 38.98gkg<sup>-1</sup> on the surface to 8.19 gkg<sup>-1</sup> deep when close to the stream.  
204 Further away from the watercourse, surface rates are 31.73gkg<sup>-1</sup> and depth rates are 6.3gkg<sup>-1</sup>.



206  
207  
208 **Figure 6:** Change in organic matter content by position



210  
211  
212


**Figure 7:** Change in MO content in layers by position

213  
214  
215

### Organic Carbon

The results of our investigations show a decrease in organic carbon in all pit areas at the base of the profile (Figure 8). However, it is noted that this organic carbon is slightly elevated at the surface layer level. For remote pits it would be the lack of plants on the ground and therefore the absence of litter. Generally, reduced organic matter intake in soil directly influences low organic carbon levels in the different profile horizons [26].

220  
221



222  
223  
224  
225

**Figure 8:** Change in CO in transects and layers relative to profile position  
F10= Close Fosse F40= Fosse distant

226

### 2.4. The Nitrogen

227

228


229

230

231

232

Analyzes show that nitrogen has approximately the same evolution in the profiles as organic carbon and organic matter (Figure 9). This suggests a correlation between these three elements.



233  
234  
235


**Figure 9:** Changes in nitrogen levels in layers

236  
237

### 3.2.5. Report C/N

238  
239  
240  
241  
242  
243  
244  
245

Analytical results indicate a decrease in C/N ratio with depth near the stream (Figure 10). This ratio ranges from 20.81 to 7.25. Biological activity in sediments is therefore reduced and organic matter decomposition is slowed. In addition, this ratio increases slightly with depth for the remote layers of the stream. Values range from 20.20 to 24.66. The slight increase in C/N in the deep layers reflects faster degradation of nitrogen compounds.



246  
247  
248

**Figure 10:** Change in C/N ratio to stream position

249  
250



251 **3.3. Statistical analyzes**

252

253 **3.3.1 Analysis of variance and significance testing**

254

255 **Results** show that there is no significant difference between areas far from the stream and areas close  
 256 to the clay. There is also no significant variation in sand levels in the different layers in the two sectors  
 257 after ANOVA (Table 2). The results show that there is no significant change in the carbon, nitrogen,  
 258 organic matter and pH levels in sediment from the 5.p.c. threshold position according to the Fischer  
 259 test. Whether the pit is distant or near the stream, the levels of chemical elements in the sediment are  
 260 substantially the same. However, significant variations are recorded by the vertical gradient (Table 3).

261

262 Table 2: Comparison of physical soil characteristics between positions and layers

263

	<b>% clay</b>	<b>% limon</b>	<b>% sand</b>	<b>% physical elements</b>
<b>Position</b>				
F10 (Near)	23,29 a	39,28 a	35,61 a	98,18 a
F40 (far away)	21,08 a	29,27 b	48,38 a	98,73 a
<b>Pr &gt; F</b>	0,8025	0,0180	0,9134	0,8834
<b>layer</b>				
C1	17,38 b	41,03 a	40,05 a	98,45 a
C2	22,75 a	33,40 a	42,33 a	98,48 a
C3	26,44 a	28,39 a	43,60 a	98,43 a
<b>Pr &gt; F</b>	0,0135	0,1702	0,9972	0,9985
<b>Moyenne</b>	0,22	0,34	0,42	0,98
<b>C.V. (p.c.)</b>	5,80	12,67	12,34	2,73

264 **Nb:** Means followed by the same letters in a column are not significantly  
 265 different from the 5 p.c. threshold

266

267

268 Table 3: Comparison of chemical characteristics between positions and layers

269

	<b>Carbon (gkg<sup>-1</sup>)</b>	<b>N (gkg<sup>-1</sup>)</b>	<b>C/N</b>	<b>OM (gkg<sup>-1</sup>)</b>	<b>pH</b>
<b>Position</b>					
F10 (Near)	12.19 a	0.65 a	18.60 a	21.01 a	5.72 a
F40 (far away)	10.56 a	0.55 a	18.34 a	18.21 a	5.79 a
<b>Pr &gt; F</b>	0.6724	0.1407	0.1709	0.6697	0.7342
<b>Layer</b>					
C1	19.93 a	1.00 a	19.84 a	34.35 a	5.67 b
C2	9.99 b	0.53 b	19.15 a	17.23 b	5.59 b
C3	4.20 b	0.27 c	16.41 a	7.25 b	6.02 a
<b>Pr &gt; F</b>	0.0119	0.0079	0.6409	0.0119	0.0388
<b>Moyenne</b>	11.38	0.60	18.47	19.61	5.76
<b>C.V. (p.c.)</b>	19.10	14.66	27.83	19.15	2.77

270 **Nb:** Means followed by the same letters in a column are not significantly  
 271 different from the 5 p.c. threshold

272

273

274 **3.3.2 Pearson Correlation**

275

276 The Pearson correlation test showed a very good correlation between carbon and organic matter with  
 277 a correlation coefficient  $r = 1$  and a probability  $P < 0.0001$ . Also, the Pearson correlation test showed  
 278 good correlations between nitrogen and organic matter with a correlation coefficient  $r = 0.989$  and a  
 279 probability  $P = 0.0001$ , between carbon and nitrogen with a correlation coefficient  $r = 0.989$  and a


280 probability  $P = 0.0001$ . The silt was positively correlated with nitrogen with a correlation coefficient  $r =$   
 281  $0.816$  and a probability  $P = 0.048$ . Clay was negatively correlated with carbon ( $r = -0.897$ ;  $P = 0.015$ ),  
 282 nitrogen ( $r = -0.893$ ;  $P = 0.016$ ) and organic matter ( $r = -0.897$ ;  $P = 0.015$ ). The silt was negatively  
 283 correlated with sand ( $r = -0.844$ ;  $P = 0.035$ ) (Figure 11).  
 284

	limon	sand	total	carbon	N	C/N	OM	pH
Clay	-0,491	-0,053	-0,311	-0,897	-0,893	-0,521	-0,897	0,605
	0,323	0,921	0,549	0,015	0,016	0,290	0,015	0,204
limon	1.00000	-0,844	-0,451	0,759	0,816	0,270	0,759	-0,589
		0,035	0,369	0,080	0,048	0,605	0,080	0,219
sand		1.00000	0,728	-0,324	-0,387	-0,011	-0,324	0,305
			0,101	0,530	0,448	0,984	0,530	0,556
total			1.00000	-0,118	-0,084	-0,316	-0,118	0,100
				0,824	0,874	0,542	0,824	0,850
carbon				1.00000	0,989	0,587	1,000	-0,656
					0,000	0,220	< 0,0001	0,157
N					1.00000	0,507	0,989	-0,657
						0,305	0,000	0,156
C/N						1.00000	0,587	-0,649
							0,220	0,163
OM							1.00000	-0,656
								0,157

285 Values in blue are different from 0 to a level of  $\alpha=0.05$  meaning  
 286 Figure 11: Correlation matrix (Pearson) of sediment characteristics following positions and layers  
 287

### 288 3.3.3 Principal Component Analysis (PCA)

289 The CPA showed that 83.76 p.c. information is reported by the F1 and F2 axes. For variables, clay,  
 290 silt, carbon, nitrogen, organic matter, and pH were well correlated with F1 factor, respectively, with  
 291 squared cosinus of 0.685, 0.711, 0.944, 0.945, 0.944, and 0.600 and the sand and total of the physical  
 292 elements that formed the F2 factor respectively with squared cosines of 0.720 and 0.747. For  
 293 individuals, F10mC1, F10mC3, F40mC1 and F40mC3 constituted the bulk of the information reported  
 294 by the F1 axis. Finally, the PCA showed a link between silt, carbon, nitrogen, organic matter and  
 295 F40mC1, F10mC1 individuals; also a link between clay, pH and the individual F10mC3; finally, a link  
 296 between sand, total physical elements and the F40mC2 horizon (Figure 12).  
 297  
 298  
 299  
 300  
 301  
 302  
 303



304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322

Figure 12: Primary Component Analysis (PCA) on F1 and F2 axes for physico-chemical characteristics of sediments according to position and layer

323  
324  
325

#### 4. DISCUSSION

326  
327  
328

##### 4.1 Variation in the physico-chemical properties of the sediment by the horizontal gradient

329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340

Results show relatively low levels of organic matter in the study area. These relatively low levels appear to be characteristic of this medium. They are the response of the disturbances related to successive floods. The accumulation of organic matter is difficult due to the flood and decay phenomena, which is an obstacle to the formation of dense vegetation cover. Successive or periodic river floods are causing changes in riparian ecosystems [27]. These results are identical to the work done by [28], who argue that alluvial soils are also characterized by low concentrations of organic matter in situ due to the absence or near-absence of surface litter. Lower M.O. values have been recorded in sediments away from the stream. These results are inconsistent with those of [29]. **These studies show that the highest concentrations of M.O. are found in areas further away from the river (5, 10, 20, 30 m).** The results of our investigations could be explained in two non-contradictory ways: first, the high sedimentation rate prevents oxidative degradation of this material at the interface, where

341 degradation processes are generally most active [30]. However, the sedimentary organic matter,  
342 partially altered by its river transit, is particularly resistant to bacterial attacks [31]. The organic matter  
343 content depends on alluvial sedimentation [32]. The organic matter transported by the river is  
344 deposited near the river. This could explain higher levels of organic matter in areas near the stream.  
345 In fact, organic matter deposited in an aquatic environment may have an indigenous origin, and to a  
346 more variable degree, an allochthonous origin [33]. A significant fraction of this organic matter is  
347 chemically and biologically degraded in the water column. A more or less significant amount (10-60%)  
348 occurs at the sediment surface [34] where it will undergo further chemical and biological  
349 transformations. A final, most stable fraction will be buried [35]. Numerous studies have shown the  
350 different impacts of water front flooding [36,37,38]. Floods and decouples can have beneficial or  
351 adverse effects on riparian ecosystems [39,40,41]. The C/N ratio is generally between 10 and 20 or  
352 higher. Values between 10 and 20 are typical in the organic matter corresponding to soil plants  
353 humified [42]. Values above 18 characterize sediments where terrestrial plant debris has accumulated  
354 [43, 24]. A low C/N ratio clearly indicates a significant source of organic matter of detritic origin [44].  
355

#### 356 **4.2 Variation in the physico-chemical properties of sediment by vertical gradient**

357

358 At the vertical gradient, our results show that the highest concentrations of MO, CO, and N are mainly  
359 in the surface layers. Our results are consistent with those of [26]. There is generally a strong  
360 correlation between organic carbon and surface layers. Most of the time the concentration of organic  
361 carbon decreases with depth. CO has very low concentrations in depth and its highest concentrations  
362 are within the first 20 centimeters of soil [45]. The high rate in this area is due to the biochemical  
363 exchanges taking place there, but this may vary depending on the conditions of the environment. The  
364 slight increase in C/N in the deep layers reflects faster degradation of nitrogen compounds, which  
365 overlaps the results of [36, 46, 24]. The distribution of organic matter in sediment depends on many  
366 and various factors. The content of a sample depends on the inputs themselves, the degree of  
367 evolution of the inputs and their dilution by the minerals [28]. In depth, we record the highest pH  
368 values. The increase in pH results in the dissolution of organic matter [47]. This may partly explain the  
369 low levels of deep organic matter. Nitrogen and organic carbon have a strong similarity in the  
370 distribution of levels within pedols. The values of these two variables decrease with depth. This trend  
371 towards a decrease in N and C.O. In fact, in the deeper layers, various studies have been carried out  
372 [48]. Organic Carbon (CO) and Organic Nitrogen (AO) are reported to have similar behavior in soils,  
373 sediments and aquatic environments [12].  
374

#### 375 **4.3 Variation of organic matter content by particle size fractions**

376

377 Results show that CO is higher in surface areas than in depths containing the highest proportions of  
378 clay. The vast majority of authors agree that the levels of clay and C.O. are positively correlated. [49],  
379 organic carbon concentration in sediments is related to the abundance of different granulometric  
380 fractions. There is a significant correlation between organic carbon and particle size distribution. The  
381 work of [50, 51] indicate that high organic carbon is often associated with a high proportion of clays in  
382 sediments. Indeed, the proportion of clay is an important factor in the stabilization of the O.C. in the  
383 soil because of the formation of the argilo-humic complex and the physical protection it provides to the  
384 O.C. linking to the inside of the aggregates. Furthermore, the stability of aggregates caused by an  
385 increase in clay levels would reduce the risk of erosion, which may affect organic carbon reserves  
386 [50,51]. There are close relationships between sediment mineralogical composition, including clay  
387 fraction and organic matter preservation [52], organic carbon concentration is higher in fine matrix  
388 sediments than those with coarse matrix [53, 48, 56, 32]. However, several studies have shown that  
389 there may be significant variability in concentrations in O.C. in the entire profile, in particular for  
390 riparian soils [48, 45]. This diversity of opinion makes any categorical conclusion difficult.  
391

### 392 **5. CONCLUSION AND RECOMMANDATION**

393  
394  
395

396 Results show that sediment from the various study sites has a sando-limonous to limono-clay texture.  
397 The observed textural variability in the area is due to the diversity of moveable deposits. This area is  
398 characterized by large and extremely rapid sedimentary rearrangements that influence sediment  
399 particle size distribution. Organic content of surface sediments on the east bank of the Bandama River  
400 in the locality of Sinématiali ranges from 31.98gkg<sup>-1</sup> to 38.98gkg<sup>-1</sup>. These relatively average rates  
401 would be related to the mineralization of organic matter, which occurs primarily within the first  
402 centimeters of the sediment, due to reducing conditions and stream-related inputs. We record a slight  
403 decrease in organic matter as we move away from the stream. But these are not significant. However,  
404 this indicates that the constant supply of alluvium transported by successive floods and reflecting  
405 current hydroclimatic conditions contributes to the organic enrichment of the near stream area. In  
406 depth, the rates obtained are very low and range from 6.3gkg<sup>-1</sup> to 8.19 3gkg<sup>-1</sup>. These low rates are  
407 due to leaching caused by periodic flooding. The MO content is generally higher in sediments of  
408 surfaces that contain the lowest proportions of clay. Organic carbon and nitrogen follow almost the  
409 same pattern as MO. These results show that successive floods have a direct effect on the dynamics  
410 of the physico-chemical properties of the sediments along the shore. An imbalance in organic matter in  
411 sediments can have a long-term impact on the vitality of the ecosystem in general.

412 **Based on the results of this study, the authorities should plan shoreline restoration programs to**  
413 **maintain the vitality and diversity of the riparian environment.**

414

415

## 416 6. REFERENCES

417

418 1. Saint-Laurent D, ST-Laurent J, Lavoie L, Ghaleb B. Use geopedological methods for the  
419 evaluation of sedimentation rates on river floodplains. CA TENA. 2008a; 73: 321-337.

420 2. Saint-Laurent D, Lavoie L, ST-Laurent J. Réurrence des inondations et édification des  
421 plaines alluviales des bassins du Centre-Sud du Québec (Canada). Revue des Sciences de  
422 l'Eau. 2009b; 22 (1) : 51-68.

423 3. Pfister L., Kwaduijk J., Musy A., Bronstert A., Hoffmann L. Climate change, land use change  
424 and runoff prediction in the Rhine-Meuse basins, River. Res. Applic. 2008a; 2004; 20: 229-  
425 241.

426 4. Boyer, C., D. Chaumont, Chartier et A. G. Roy. "Impact of climate change on the hydrology of  
427 St. Lawrence tributaries". Journal of Hydrology. 2010 ; 384(1-2): 65-83.

428 5. Clinton S.M., Power M., Swenson R.. Impacts of inundation regime, floodplain vegetation, and  
429 burrowing animals on the incorporation of carbon into floodplain soils, CALFED Science  
430 Fellows Program, Project Number: R/SF-1, 2006, 2003:1-9.

431 6. Watkins S.C., Quinn G.P., Gawne B.. Changes in inorganic-matter dynamics and physic  
432 chemistry, associated with riparian vegetation loss and river regulation in floodplain wetlands  
433 of the Murray River, Australia, Marine and Fresh water Research. 2010; 61:1207-1217.

434 7. Power EA., Chapman PM. Assessing sediment quality, In Burton, GAJ (Eds). Sediment  
435 Toxicity Assessment, Lewis, MI, USA. 1992 :1-18.

436 8. Cyril M. Origine et devenir de la matière organique des sédiments de mangroves de Guyane  
437 française. - Précurseurs, Environnements de dépôt, Processus de décomposition et Relation  
438 avec les métaux lourds –. Géologie appliquée. Université d'Orléans. 2003 : 272.

439 9. Siegenthaler, U., Sarmiento, JL. Atmospheric carbon dioxide and the ocean. Nature 1993;  
440 365: 119-125.

441 10. Filella, M.. Freshwaters : which NOM matters ? *Environmental Chemistry Letters*. 2009; 7 (1) :  
442 21-35.

443 11. Labanowski, J. *Matière organique naturelle et anthropique : vers une meilleure*  
444 *compréhension de sa réactivité et de sa caractérisation*. Thèse, Université de Limoges. 2004 :  
445 199.

446 12. Kalbitz K, Geyer S. Different effects of peat degradation on dissolved organic carbon and  
447 nitrogen. *Org. Geochem*. 2002 ; 33(3): 319-326.

448 13. Zhu WX, Carreiro MM. Variations of soluble organic nitrogen and microbial nitrogen in  
449 deciduous forest soils along an urban–rural gradient. *Soil Biol. Biochem*. 2004; 36: 279-288.

450 14. Croué JP, Benedetti MF, Violleau D, Leenheer JA. Characterization and copper binding of  
451 humic and nonhumic organic matter isolated from the South Platte river : Evidence for the  
452 presence of nitrogenous binding site. *Environ. Sci. Technol*. 2003; 37(2): 328-336.

453 15. Meyers PA., Lallier-Vergès E. Lacustrine sedimentary organic matter records of Late  
454 Quaternary paleoclimates. *J. Paleolimnol*. 1999; 21: 345-372.

- 455 16. AFNOR. Qualité des sols : Méthodes d'analyse, Recueil de normes françaises. Afnor, 1ère  
456 édition. 1987 : 136.
- 457 17. Shepard F. Nomenclature based on sand-silt-clay ratios. *Journal of sedimentary Petrology*.  
458 1954 ; 24 : 151-158.
- 459 18. Bonnet C. Développement de bio-essais sur sédiments et applications de l'étude, en  
460 laboratoire, de la toxicité de sédiments dulçaquicoles contaminés. Thèse de l'Université de  
461 Metz. 2000 : 326.
- 462 19. Mamadou A. S. Hydrologie et géochimie des transports fluviaux dissous et particuliers dans  
463 le bassin versant du Milo (République de Guinée), Thèse de doctorat, Université de Toulouse  
464 (INPT). 2018 : 183 .
- 465 20. Rofes G. Etude des sédiments. méthodes de prélèvement et d'analyses pratiquées au  
466 laboratoire de sédimentologie. Etude N° 47, *Ministère de l'Agriculture, GTGREF*. 1980 : 50 p.
- 467 21. Aubert G. Méthodes d'analyses des sols. *Centre National de Documentation Pédagogique*,  
468 Marseille. 1978 : 189.
- 469 22. Thurman EM, Malcolm RL. Preparative isolation of aquatic humic substances. *Environ. Sci.*  
470 *Technol.* 1981 ;15(4) : 463-466.
- 471 23. Baize D. Guide des analyses en pédologie. Ed. INRA, Paris. 2000 : 257.
- 472 24. Delmas D. Minéralisation de la matière organique et échanges ioniques à l'interface eau-  
473 sédiment de l'étang de Berre, Méditerranée. *Oceanol Acta*. 1980 ; 3(3) : 347-356.
- 474 25. Lele N. Potentiel d'amélioration de la fertilité des sols sableux et acides de Kinshasa (RDC)  
475 par l'usage du charbon des bois (biochar), de la biomasse végétale et des engrais minéraux  
476 Thèse Université de Liège. 2016 : 196.
- 477 26. Drouin A, Saint-Laurent D, Lavoie L, Ouellet C. High-Precision Digital Elevation Model to  
478 evaluate the spatial distribution of Soil organic carbon inactive floodplains. *Wetlands*. 2011;  
479 31: 1151-1164.
- 480 27. Saint-Laurent D. Reconstruction of flood events and links with climatic factors: a case study of  
481 the Saint-François Basin. *JETS*. 2007 ; 31: 5-24.
- 482 28. Daniel J, Dupont J, Christian Jouannic C. Etude de la relation entre le carbone organique et  
483 l'azote dans les sédiments de la Baie d' Ambaro. ORSTOM Nosy-Bé, Doc. Multigr.  
484 1970; 16:11.
- 485 29. Bridgham D, Megonigal JP, Kedler JK, Bliss NB, Trettin C. The carbon balance of North  
486 American wetlands, *Wetlands*. 2006; 26(4): 889-916.
- 487 30. Hartman M, Muller PJ., Suess E, Van der Weijden CH., Chemistry of late quaternary  
488 sediments and their interstitial waters from the N-W African continental margin, "*Meteor*"  
489 *Forschungsergeh., Reihe C*. 1976; 24: 1-67.
- 490 31. Bordovskiy O. K. Accumulation and transformation of organic substances in marine  
491 sediments. 2. Source of organic matter in marine basins, *Mar. Geol.* . 1965 a ;3:5-31.
- 492 32. Cierjacks A, Kleinschmit B, Kowarik L, Graf M, Lang F. Organic matter distribution in  
493 floodplains can be predicted using spatial and vegetation structure data, *River. Res. Applic*;  
494 2010 ; 1-10.
- 495 33. Bordovskiy OK. Accumulation and transformation of organic substances in marine sediments.  
496 4. Transformation of organic matter in bottom sediments and its early diagenesis, *Mar. Geol.*  
497 1965 b ; 3 83-114.
- 498 34. Ouertani N., Hamouda R. et Belayouni H. Etude de la matière organique dans les sédiments  
499 superficiels d'un système aquatique confiné anoxique évoluant en milieu urbain : cas du Lac  
500 sud de Tunis. *Géo-Eco-Trop*, 2006 ; 30(2): 21-34
- 501 35. BELIN S. Distribution microscopique de la matière organique disséminée dans les roches  
502 mères : technique d'étude, interprétation des conditions de dépôt et de diagenèse. Thèse de  
503 doctorat de l'Université d'Orsay. 1992 : 371.
- 504 36. Kaplan 1R, Rittenberg SC. Basin, sedimentation and diagenesis, in: *The Sea 3*, edited by M.  
505 N. Hill, Interscience Publish, New York, London. 1963: 583-619.
- 506 37. NESTLER JM., LONG KS. Development of hydrological indices to aid cumulative impact  
507 analysis of riverine wetlands, *Regulated Rivers: Research & Management*. 1997; 13: 317-334.
- 508 38. MONIRUL M, MIRZA Q, WARRICK RA., ERICKSEN NJ. The implications of climate change  
509 on floods on the Ganges, Brahmaputra and Meghna Rivers in Bangladesh, *Climatic Chang*.  
510 2003 ; 57: 287-318.
- 511 39. SHIELDS FD, KNIGHT SS, COOPER CM. Cyclic perturbation of lowland river channels and  
512 ecological response, *Regul. Rivers: Res. Mgmt*. 2000; 16: 307-325.

- 513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549
40. Lavoie L, Saint-Laurent D, St-Laurent J. Analyse pédologique et sédimentologique des sols alluviaux et paléosols des terrasses d'inondation. *Canadian Journal of Soil Science*. 2006 ; 86 : 1-14.
  41. FRAZIER P, PAGE K. A reach-scale remote sensing technique to relate wetland inundation to river flow, *River. Res. Applic.* 2008; (25) : 836-849.
  42. Debyser J. Contribution à l'étude géochimique des vases marines. *Inst. Fr. du Pétrole - Division Sédimentologie - Réf. 6005 Edit. Technip PARIS*. 1961:249 p
  43. Waksman SA.. On the distribution of organic matter in the sea bottom and the chemical nature and origin of marine humus, *Soil. Sei.* 1933; 36: 125-147.
  44. Cifuentes LA, Sharp JH, Fogel ML. Stable carbon and nitrogen isotope biogeochemistry in the Delaware estuary. *Limnology and Oceanography*. 1988; .33: 1102-1115.
  45. Su ZY, Xiong YM, Zhu JY, Ye YC, Ye M. Soil organic carbon content and distribution in a small landscape of Dongguan, South China, *Pedosphere*. 2006; 16(1): 10-17.
  46. Bordovskiy OK. Accumulation and transformation of organic substances in marine sediments. 2. Source of organic matter in marine basins, *Mar. Geol.* 1965a ; (3) : 5-31.
  47. Chaignon V. Biodisponibilité du cuivre dans la rhizosphère de différentes plantes cultivées. Cas de sols viticoles contaminés par des fongicides. Ecole doctorale, Sciences de l'Environnement : Système Terre, Aix-Marseille III. 2001 : 183.
  48. Schilling KE, Palmer JA, Bettis lii EA, Jacobson P, Schultz RC, Isenhardt TM. Vertical distribution of total carbon, nitrogen and phosphorus in riparian soils of Walnut Creek, southern Iowa, *Catena*. 2009; 77: 266-273.
  49. Nadeu E, De Vente J, Martinez-Mena M, Boix-Fayos C. Exploring partic1e size distribution and organic carbon pools mobilized by different erosion processes at the catchment scale, *J Soil Sediment*. 2010; Vol. II: 667-678.
  50. Krull ES., Baldock IA., Skjemstad JO. Importance of mechanisms and processes of the stabilization of soil organic matter for modelling carbon turnover. *Functional Plant Biology*. 2003; 30: 207-222.
  51. Jindaluang W, Kheoruenromne L, Suddhiprakarn Bhupinder PS., Balwant S. Influence of soil texture and mineralogy on organic matter content and composition in physically separated fractions soils . of Thailand. *Geoderma*. 2013.195/196 : 207-219.
  52. Keil, RG, Montlucon, DB, PrahI, FR, Hedges JI. Sorptive preservation of labile organic matter in marine sediments. *Nature*. 1994; 370 : 549-552.
  53. GREENER B, STROMQUIST L. Overbank sedimentation rates of fine grained sediments. A study of the recent deposition in the lower river Fyrisan. *Geografiska Annaler*. 1987 ; 69A: 139-146.