

# 1 Plankton Composition in relation to Water Quality in the coastal waters of Nigeria

2 \*Ajibare A.O.<sup>1</sup>, Ayeku P.O.<sup>2</sup>, Akinola J.O.<sup>3</sup> and Adewale A.H.<sup>2</sup>

3 <sup>1</sup>Department of Biological Sciences, KolaDaisi University Ibadan, Nigeria.

4 <sup>2</sup>Department of Biological Sciences, Wesley University Ondo, Nigeria.

5 <sup>3</sup>Department of Fisheries and Aquaculture Technology, Federal University of Technology Akure,  
6 Nigeria.

7 \*Corresponding Author: [mrajifem@yahoo.com](mailto:mrajifem@yahoo.com),

## 8 Abstract

9 Coastal waters of Ondo State, Nigeria have diverse assemblage of fish, yet there is dearth of  
10 information on its plankton composition. This study investigates plankton components in relation  
11 to physico-chemical characteristics of the coastal waters bordering Olotu, Ayetoro and Bijimi in  
12 providing baseline information that can be used for planning and implementation of policies for  
13 monitoring, impacts assessment and conservation. Surface water samples were collected on  
14 monthly basis from March to June 2015 to analyze physico-chemical parameters while plankton  
15 net of 55µm mesh size was used for collection of plankton using standard methods prescribed by  
16 APHA. The light and dark bottle method was used to determine primary productivity. Shannon-  
17 wiener, Margalef and Equitability Indices were used for diversity. Values of the physico-  
18 chemical parameters observed ranged as follows: temperature, 27.47±2.06-29.27±0.31°C;  
19 turbidity, 43.43±0.91-65.33±2.52NTU; pH, 5.54±0.31-6.12±0.30; BOD, 2.20±0.29-  
20 5.43±0.54mg/l; COD, 6.08±2.71-6.66±1.52mg/l; dissolved oxygen, 6.39±0.39-7.78±0.19mg/l  
21 and salinity, 2.03±0.06-3.77±0.04mg/l. Fifteen species of phytoplankton and three  
22 developmental stages of zooplankton were recorded. Phytoplankton accounted for 83.3% as  
23 against 16.7% zooplankton. Diatoms (93.3%) and dinoflagellates (6.7%) represented  
24 phytoplankton whereas 66.7% of zooplankton belonged to the phylum Arthropoda. Primary  
25 productivity ranged between 132.194±13.48m<sup>-3</sup>hr<sup>-1</sup> and 134.48±15.27m<sup>-3</sup>hr<sup>-1</sup>. Some dominant  
26 species recorded were *Coscinodiscus*, *Biddulphia*, *Copepod*, *Skeletonema* and *Ditylum*. pH and  
27 Temperature were major determinant of the composition, diversity and abundance of plankton.  
28 The observed plankton group indicates the suitability of the creeks as habitat and breeding  
29 ground for diverse aquatic species. The water quality falls within acceptable range hence the  
30 environment can be classified as healthy ecosystem.

31 **Key Words:** Phytoplankton, zooplankton, physico-chemical, Estuarine, algae and pollution.

## 32 INTRODUCTION

33 Planktons are major contributors of biomass and are crucial to the productivity and sustainability  
34 of the aquatic ecosystem (Harris and Vinobaba, 2012). Phytoplankton constitute the base of the  
35 aquatic food chain, producing organic and inorganic substances through carbon dioxide and  
36 photosynthesis. However, their composition, distribution and abundance are largely influenced  
37 by various environmental factors from physical, biological and chemical changes. These factors  
38 include; pollution, urbanization, industrialization, anthropogenic activities, climate change etc.

39 Biological approaches to assessing the water quality (interacting physical and chemical factors)  
40 in aquatic ecosystems cannot be over-emphasized, considering its influence on the levels of  
41 trophic structure, primary productivity and total biomass in the aquatic food web (Offem *et al.*,  
42 2011). The coastal waters of Ondo State, Nigeria which is richly blessed with favourable  
43 ecological and climatological conditions that favours optimum growth, reproduction and  
44 productivity of aquatic species (Daramola *et al.*, 2009). These conditions support fish farming

45 operations, thus improving the socio-economic status and livelihood of the coastal dwellers and  
46 country. However, much concern has been given to the assessment of eco-environmental quality  
47 due to the increase in agricultural, anthropogenic and industrial activities that exist in and around  
48 the area (Benson *et.al*, 2007; Olawusi-Peters *et al.*, 2017).

49 Several authors have carried out extensive analysis on the trophic status of various coastal water  
50 bodies in Nigeria (Imoobe, 2011; Cako *et al*, 2013) and reported changes in mean temperature,  
51 nutrient availability and hydrology to be the most crucial variables that determine the abundance,  
52 distribution of plankton and productivity of the entire ecosystem (Offem *et al.*, 2011). Planktons  
53 therefore become ideal for biomonitoring of the ecological changes in coastal ecosystems  
54 considering their position in the aquatic food chain. They reflect the composite influence of  
55 different parameters of water quality in the waterbody (Gharib *et al.*, 2011).

56 Comprehensive researches have been done in the coastal waters of Ondo State: Akegbejo-  
57 Samson (1995); Asaolu (1998); Adeparusi (2003); Adebowale *et al.*, (2008); Abdus-Salam *et al.*,  
58 (2010); Bayode *et al.*, (2011); Olawusi-Peters *et al.*, (2014; 2017). However, the composition,  
59 diversity and abundance of plankton in the region are yet to be investigated. Thus, this study  
60 provides baseline information that bridges the existing gap in research on the diversity and  
61 abundance of plankton as it relates to the quality and productivity of the coastal waters of Ondo  
62 State, Nigeria.

## 63 **MATERIALS AND METHOD**

### 64 **Study Area**

65 This study was conducted in three communities within the coastal area of Ondo state (Ilaje Local  
66 Government Area), Nigeria. The study area falls within Latitudes  $06^{\circ}$  &  $06^{\circ} 30^1$  North and  
67 Longitudes  $004^{\circ} 45^1$  and  $005^{\circ} 45^1$  East of the Greenwich Meridian. Olotu, Ayetoro and Bijimi  
68 were purposively selected based on the human population, catch volume, diversities of the fishes  
69 and possible anthropogenic inputs in the areas. The area is positioned within the equatorial  
70 evergreen swamp forest sharing boundaries with Okitipupa Local Government Area in the North;  
71 the Atlantic Ocean in the South; Ijebu Waterside Local Government Area (Ogun State) in the  
72 West and Delta State in the East (Bayode *et al.*, 2011). The environment has two seasons; the dry  
73 and the wet seasons and it experiences consistently high temperatures (about  $32^{\circ}\text{C}$ ) all year  
74 round. Since temperature varies only slightly, rainfall distribution, over space and time, becomes  
75 a single crucial factor (Adeparusi *et al.*, 2003). Ilaje LGA consists of over five hundred  
76 settlements spreading over  $3,000 \text{ km}^2$  with emerging communities dispersed within the coast and  
77 an increasing population size of 2.2% annually (Adebowale *et al.*, 2008).

### 78 **Collection of water samples and determination of physico-chemical parameters**

79 Water samples for the physico-chemical analysis were collected on monthly basis from each  
80 station at sub-surface level, using 250 ml sampling bottles and transported in ice-chest to the  
81 laboratory for analysis according to APHA (1998). Samples were taken at the three stations on  
82 the same day and at the same sampling points for ease of reference. Temperature, pH, Turbidity,  
83 Salinity, Dissolved Oxygen, Chemical Oxygen Demand (COD) and Biological Oxygen Demand

84 (BOD) of the water from each sampling station was determined using the Hanna multi-parameter  
85 Model HI 9828.

### 86 **Sampling of Plankton**

87 The plankton samples were collected using sampling net of 55µm-mesh size net tied to the boat  
88 towed at low speed for 10 minutes. The sample were then preserved in 4% formalin and kept  
89 refrigerated prior to analysis. In the laboratory, three replicates of each sample concentrate were  
90 observed under microscope Olympus model using standard guides for identification (UNESCO,  
91 1978 and Yamaguchi and Gould 2007). The different components of the plankton samples were  
92 determined and classified by their relative abundance.

### 93 **Determination of Primary Productivity**

94 The light and dark bottle method was used in the determination of primary productivity using  
95 three 125ml oxygen bottles with each designated as initial, light and dark oxygen bottle (Boyd,  
96 1979). The dark bottles were painted with black paint and covered with aluminum foil to prevent  
97 light penetration. All the bottles were filled with water from each depth. The initial water bottle  
98 was immediately fixed for dissolved oxygen using managanous sulphate and alkaline solutions.  
99 The bottles (light and dark) were incubated for six hours. and the dissolved oxygen concentration  
100 was determined for the initial, light and dark bottles. Thus, the primary productivity was  
101 calculated using conversion formula according to Smith *et al.*, (1996).

$$\text{Gross primary productivity} = \frac{(LB - DB)}{T \text{ (hrs)}} \times 0.375 \times 1000 \text{mg (m}^{-3}\text{hr}^{-1}\text{)}$$

102 Where: LB = DO of light bottles

103 DB = DO of dark bottles

104 T = Incubation period

105 0.375 = ratio of weight of carbon to oxygen

106 1000 = litres in m<sup>3</sup>

### 107 **Statistical Analysis**

108 Data were subjected to Multi-Variate Analysis of Variance (MANOVA) to evaluate statistical  
109 variation across the stations (P=0.05) using Statistical Package for Social Sciences (SPSS) version  
110 16.0. Standard deviations were estimated. Descriptive analysis was also used to present tables and  
111 figures. Principal Component Analysis (PCA) was used to determine the relationship between the  
112 plankton and the physico-chemical parameters using PAST software.

113 The Plankton data were subjected to the following ecological indices:

Index	Formula	Reference
Margalef's Diversity Index	$d = \frac{S - 1}{\ln N}$	Margalef (1968)
Shannon-Wiener Index	$H' = (-P_1 \ln P_1) + (-P_2 \ln P_2) + \dots$	Shannon and Wiener (1963)
Pielou's Equitability Index	$J = \frac{H'}{\ln(S)}$	(Pielou, 1966)
Number of Occurrence Index (NOI)	$NOI = \frac{n}{N} \times 100$	Ogbeibu (2005)

114 Where:

115 S = number of observed species;

116 N = total number of individuals of all species in the catch;

117 H' = Shannon-Wiener index;

118 P = total proportion of each species in sample;

119 n = number of individuals of each species in the catch.

## 120 RESULTS

### 121 Physico-Chemical Parameters of Coastal Waters of Ondo State

122 The physico-chemical parameters of water in the three stations is presented in Table 1. The table  
 123 revealed that temperature, Dissolved Oxygen (DO), salinity, BOD, COD showed no significant  
 124 difference across the stations, while turbidity and pH exhibited locational variation at 95%  
 125 confidence limit. The table shows that the highest mean Temperature (28.48±0.91 °C) was  
 126 recorded in Bijimi while the lowest temperature (27.88±1.26 °C) was recorded in Ayetoro. The  
 127 lowest (51.48±7.76 NTU) and highest (53.86±8.47 NTU) mean turbidity was recorded at Bijimi  
 128 and Ayetoro respectively, while the highest mean (5.86±0.23) and lowest mean pH (5.66±0.20)  
 129 was recorded at Olotu and Ayetoro respectively. Moreover, the lowest (7.30±0.46 mg/l) and  
 130 highest mean DO (7.42±0.31 mg/l) was recorded in Bijimi and Olotu respectively, while the  
 131 lowest (3.34±1.63mg/l) and highest BOD (3.74±1.15mg/l) was recorded at Ayetoro and Bijimi  
 132 respectively. The table further shows that the Salinity was highest (3.31±0.65mg/l) and lowest  
 133 (3.03±0.70mg/l) at Bijimi and Ayetoro respectively, while COD was highest (6.66±1.52 mg/l)  
 134 and lowest (6.08±2.71 mg/l) at Bijimi and Ayetoro respectively.

135 **Table 1: Physico-chemical parameters of coastal waters of Ondo State**

Parameter	Stations		
	Olotu	Ayetoro	Bijimi
Temperature (°C)	27.91±1.13 <sup>a</sup>	27.88±1.26 <sup>a</sup>	28.48±0.91 <sup>a</sup>

Turbidity (NTU)	52.04±6.94 <sup>ab</sup>	53.86±8.47 <sup>b</sup>	51.48±7.76 <sup>a</sup>
pH	5.86 ±0.23 <sup>b</sup>	5.66 ±0.20 <sup>a</sup>	5.80 ±0.25 <sup>ab</sup>
DO (mg/l)	7.42±0.31 <sup>a</sup>	7.35±0.64 <sup>a</sup>	7.30±0.46 <sup>a</sup>
BOD (mg/l)	3.50±1.48 <sup>a</sup>	3.34±1.63 <sup>a</sup>	3.74±1.15 <sup>a</sup>
Salinity (mg/l)	3.16±0.67 <sup>a</sup>	3.31±0.65 <sup>a</sup>	3.03±0.70 <sup>a</sup>
COD (mg/l)	6.61±2.57 <sup>a</sup>	6.08±2.71 <sup>a</sup>	6.66±1.52 <sup>a</sup>

### 136 Plankton Composition and Abundance in Coastal Waters of Ondo State

137 The observed composition of plankton is presented in Table 2 which shows that 18 species from  
 138 15 families were recorded throughout the study. The phytoplankton community composed of 14  
 139 diatoms and 1 dinoflagellate while the zooplankton community was made up of fish embryo,  
 140 copepod naupli and odonata nymphs.

141 The abundance of plankton as shown in Table 3 reveals that *Coscinodiscus spp* had the most  
 142 abundance in the three stations with 19.05% (n=8) in Olotu, 17.65% (n=6) in Ayetoro and  
 143 15.38% (n=4) in Bijimi. The table also reveals that *Biddulphia mobiliensis* 9.52% (n=4), 5.88%  
 144 (n=2), 7.69% (n=2); *Copepod naupli* 4.76% (n=2), 5.88% (n=2), 7.69% (n=2); *Fish embryo*  
 145 4.76% (n=2), 5.88% (n=2), 7.69% (n=2);  
 146 (=2); *Biddulphia aurita* 7.14% (n=3), 8.82% (n=3), 3.85% (n=1); and *Skeletonema costatum*  
 147 9.52% (n=4), 11.76% (n=4), 3.85% (n=1) were found in Olotu, Ayetoro and Bijimi accordingly.

148 The ecological indices including Margalef's diversity index (*d*), Pielou's measure of evenness  
 149 (*J'*) Shannon-Wiener index (*H'*), and the number of species (*S*) caught across the three stations  
 150 are presented in Table 4. The number of species caught throughout the period of study ranged  
 151 between 12 and 14 with the highest recorded at Bijimi and lowest in Ayetoro. The Margalef's  
 152 diversity index (*d*) of the plankton ranged from 7.18 (Ayetoro) to 8.48 (Bijimi), while the  
 153 Shannon-Wiener index (*H<sub>s</sub>*) ranged from 2.36 (Ayetoro) to 2.50 (Olotu). The Pielou's measure of  
 154 evenness (*J'*) range from of 2.18 (Olotu) to 2.22 (Bijimi). The table also reveals that the primary  
 155 productivity of the three stations were not significantly different from one another with the  
 156 highest value (134.48±15.27) recorded at Olotu while the lowest value (132.19±13.48) was  
 157 recorded at Ayetoro.

158 **Table 2: Composition of plankton in coastal waters of Ondo state**

Species	Family	Type of Plankton
<i>Coscinodiscus sp</i>	Coscinodiscaceae	Phytoplankton (Diatom)
<i>Biddulphia mobiliensis</i>	Biddulphiales	Phytoplankton (Diatom)
<i>Biddulphia aurita</i>	Biddulphiales	Phytoplankton (Diatom)
<i>Skeletonema costatum</i>	Skeletonemataceae	Phytoplankton (Diatom)
<i>Ditylum brightwelli</i>	Lithodesmiaceae	Phytoplankton (Diatom)
<i>Odontella sp</i>	Eupodiscaceae	Phytoplankton (Diatom)

<i>Thalassiora sp</i>	Thalassiosiraceae	Phytoplankton (Diatom)
<i>Rhizosolenia</i>	Rhizosoleniaceae	Phytoplankton (Diatom)
<i>Pseudo-nitzschia</i>	Bacillariaceae	Phytoplankton (Diatom)
<i>Brachionus falcatus</i>	Branchionidae	Phytoplankton (Diatom)
<i>Brachionus quadridentatus</i>	Lepadellidae	Phytoplankton (Diatom)
<i>Lepadella patella</i>	Lithodesmiaceae	Phytoplankton (Diatom)
<i>Asplanchna brightwelli</i>	Lithodesmiaceae	Phytoplankton (Diatom)
<i>Filinia oregonensis</i>	Trichoshaeridae	Phytoplankton (Diatom)
<i>Ceratium hirundinella sp</i>	Ceratiaceae	Phytoplankton (Dinoflagellate)
Copepod nauplii	Copepoda	Zooplankton
Fish embryo	Osteichthyes	Zooplankton
<i>Odonata nymphs</i>	-	Zooplankton

---

159

160

161 **Table 3: Number of Occurrence Index (% and n) of plankton in coastal waters of Ondo**  
 162 **state**


SPECIES	Olotu		Ayetoro		Bijimi	
	NOI (%)	n	NOI (%)	N	NOI (%)	N
Copepod nauplii	4.76	2	5.88	2	7.69	2
<i>Coscinodiscus sp</i>	19.05	8	17.65	6	15.38	4
<i>Biddulphia mobiliensis sp</i>	9.52	4	5.88	2	7.69	2
Fish embryo	4.76	2	5.88	2	7.69	2
<i>Biddulphia aurita sp</i>	7.14	3	8.82	3	3.85	1
<i>Skeletonema costatum</i>	9.52	4	11.76	4	3.85	1
<i>Ditylum brightwelli</i>	0	0	5.88	2	7.69	2
<i>Odontella sp</i>	0	0	11.76	4	0.00	0
<i>Ceratium hirundinella sp</i>	0	0	2.94	1	0.00	0
<i>Thalassiora sp</i>	0	0	11.76	4	0.00	0
<i>Rhizosolenia</i>	2.38	1	2.94	1	0.00	0
<i>Pseudo-nitzschia</i>	7.14	3	8.82	3	0.00	0
<i>Brachionus falcatus</i>	4.76	2	0.00	0	7.69	2
<i>Brachionus quadridentatus</i>	11.9	5	0.00	0	11.54	3
<i>Lepadella patella</i>	4.76	2	0.00	0	7.69	2
<i>Asplanchna brightwelli</i>	4.76	2	0.00	0	3.85	1
<i>Filinia oponienses</i>	4.76	2	0.00	0	3.85	1
<i>Odonata nymphs</i>	4.76	2	0.00	0	11.54	3

163 **Table 4: Biodiversity Indices of plankton and Primary productivity in coastal waters of**  
 164 **Ondo state**  
 165

Biodiversity Index	Olotu	Ayetoro	Bijimi
Shannon-Wiener (H')	2.50 <sup>b</sup>	2.36 <sup>a</sup>	2.47 <sup>ab</sup>
Margalef's Index (d)	8.02 <sup>b</sup>	7.18 <sup>a</sup>	8.48 <sup>c</sup>
Pielou's measure of Evenness (J)	0.66 <sup>a</sup>	0.64 <sup>a</sup>	0.69 <sup>a</sup>
Number of species (S)	14.00 <sup>a</sup>	13.00 <sup>a</sup>	12.00 <sup>a</sup>
Primary productivity	133.91±9.79 <sup>a</sup>	132.19±13.48 <sup>a</sup>	134.48±15.27 <sup>a</sup>

166

167 The initial component matrix indicates that most physico-chemical parameters showed high  
 168 values in the first principal component (PC1) which means that 38% of the total variance loads  
 169 heavily on BOD (0.976), COD (0.957), salinity (0.971), turbidity (0.871) and DO (0.676). The  
 170 second principal component (PC2) which accounts for 30% of the total variance, exhibited  
 171 elevated loadings for Shannon-Wiener Index (0.679), Margalef's Index (0.293), Evenness Index  
 172 (0.558), pH (0.877) and Temperature (0.744) as shown in Figure 1 and Table 5.  
 173



174  
 175 **Figure 1: Relationship between Plankton and Water quality of the coastal waters of Ondo**  
 176 **State Nigeria**

177 **Table 5: Principal Components Analysis of Plankton and Physico-Chemical properties of**  
 178 **the coastal waters of Ondo state, Nigeria**

	Component	
	PC1	PC2
Shannon Wiener Index	-0.201	0.679
Margalef's Diversity Index	0.09	-0.293
Evenness Index	0.037	-0.558
Species (Taxa)	-0.069	0.888
Temperature	-0.258	-0.744
Turbidity	0.871	-0.197
pH	0.038	0.877
DO	0.676	0.468
BOD	0.976	-0.12
Salinity	-0.971	-0.036
COD	0.957	-0.046
% Variance	38	30
Eigen value	4.15	3.24



## 179 DISCUSSION

180 Temperature which is the most important physical variable affecting the metabolic rate of living  
181 organisms in the aquatic environment was within the optimal water temperatures of 18°C –  
182 38°C, for optimum production and growth of planktons and fish in water bodies (Begum *et al.*,  
183 2003). The result was also similar to previous findings (Adebowale *et al.*, 2008; Ajibare 2014;  
184 Bolarinwa *et al.*, 2016 and Olawusi-Peters and Akinola 2017) in the Coastal waters of Ondo  
185 State. The water pH and turbidity values were significantly different across the stations and  
186 could be associated with the presence of suspended particles due to sewage discharges, acid  
187 runoffs and other human-mediated activities such as dredging and other activities in and around  
188 the study area. Moreover, the pH was in consonance with the report of Onyema and Ojo (2008),  
189 Nkwoji, (2010); Abowei (2010) and Ajibare (2014) in the Niger Delta region.

190 The salinity of this study reveals a brackish environment with no significant difference across the  
191 stations. This supports the distribution, abundance, growth and diversity of planktons and aquatic  
192 organisms that requires stable salinity for survival. The DO concentrations obtained across the  
193 stations supports the relationship between BOD and COD. High DO and Low BOD values are  
194 mainly due to higher algal productivity, along with increased solubility of oxygen at low  
195 temperatures (Ajibare, 2014) as observed in this study. The observed COD were higher when  
196 compared to BOD, signifying that the study area was highly affected by organic matter which  
197 comes from dead and decayed plant and animals in the ecosystem. These also clearly indicates  
198 that the areas receive high amount of sewage waste containing high level of organic matter that  
199 supports the growth of planktons with adverse effect on aquatic ecosystem.

200 The composition, species richness, abundance and diversity indices revealed 18 different species  
201 across the stations which ranged from 12 to 14 per station. This shows that the study area was  
202 relatively rich in the assemblage of plankton. The Pielou's measure of evenness showed that the  
203 three stations had low dominance of a single-species because the values were closer to one (1)  
204 than zero (0). The Shannon-Weiner index ( $H'$ ) also suggests a generally high diversity and  
205 similarity among the plankton communities in the study area and this is buttressed by the non-  
206 variation of the indices across the stations. However, the differences in the level of  
207 anthropogenic activities/pollution in the communities may be responsible for the slight variation  
208 obtained in the Margalef's diversity as it was also observed by Olawusi-Peters and Ajibare  
209 (2014) who worked on the species richness and abundance of the coastal waters of Ondo State.

210 The percentage composition/Number of Occurrence Index (%) of plankton's families at the three  
211 different stations showed that Coscinodiscaceae (*Coscinodiscus sp*) dominated the plankton  
212 groups, followed by Skeletonemataceae (*Skeletonema costatum*), Biddulphiales (*Biddulphia*  
213 *mobiliensis* and *Biddulphia aurita*), Lepadellidae (*Brachionus quadridentatus*), Osteichthyes  
214 (Fish embryo), Bacillariaceae (*Pseudo-nitzschia*), and the least represented family was  
215 Ceratiaceae (*Ceratium hirundinella sp*). The dominance of family Coscinodiscaceae across the  
216 stations was similar to the findings of Varadharajan and Soundarapandian, (2015). Diatoms were  
217 dominant in the plankton collection and could be as a result of their ability to tolerate wide  
218 geographical and climatic conditions (Balogun and Ajani, 2015). In addition, diatoms are  
219 considered euryhaline and eurythermal species as they grow and are widely distributed under  
220 marine conditions (Emmanuel and Onyema, 2007). The trend of dominance in plankton  
221 composition across the stations followed the order Bijimi<Ayetoro<Olotu. The high diversity

222 and population of plankton in Olotu could be adduced to high tidal influx, organic production  
223 and nutrient availability in the area.

224 According to Adirondack Ecologists (2010), the species abundance and composition of  
225 phytoplankton have significant effects on both the water quality and clarity of aquatic  
226 ecosystems. The eighteen species and fifteen families of phytoplankton caught in this study  
227 revealed the area to be an area of high primary productivity, where the rate of photosynthesis is  
228 relatively high. This is in line with the report of Offem *et al.*, (2011) who stated that  
229 phytoplankton carry out most of the primary productivity that takes place in the marine/brackish  
230 environment eventhough other marine plants (e.g. algae) contribute to primary productivity. This  
231 research also compares favourably with the reports of Balogun and Ajani (2015) who worked on  
232 the coastal waters of Lagos state and tidal creeks of south-west Nigeria.

233 The analysis of the relationship between plankton and physicochemical properties reveals that  
234 BOD, COD, salinity, turbidity and DO were highly correlated and had 38% influence on the  
235 entire aquatic ecosystem. Similarly, diversity indices (Shannon-Wiener Index, Margalef's Index  
236 and Evenness Index) correlated significantly with pH and Temperature to have 30% influence on  
237 the health of the waterbody. This shows that the composition, distribution and diversity of  
238 plankton are influenced by the physicochemical parameters of the aquatic ecosystem.

## 239 **CONCLUSION**

240 This study revealed that the physico-chemical parameters fall within acceptable range and it  
241 directly influences the occurrence, growth, diversity and distribution of plankton in the study  
242 area. The composition of plankton indicates suitability of the environment as habitat and  
243 breeding ground for diverse aquatic species. The study also revealed that BOD, COD, salinity,  
244 turbidity, DO, pH and temperature were major determinant of the composition, abundance and  
245 diversity of plankton. The variation in the plankton groups reflects the locational dynamics and  
246 the impact of human activities or/and pollution on the water quality. Thus, proper and continuous  
247 monitoring of the coastal waters of Ondo State is recommended in order to safeguard the entire  
248 ecosystem from collapse as well as contribute to the survival and growth of aquatic organisms.

## 249 **REFERENCES**

- 250 Abdus-Salam, N., Adekola F.A. and Apata A.O. (2010): A Physicochemical Assessment of  
251 Water quality of oil producing areas of Ilaje, Nigeria. *Advances in Natural and Applied*  
252 *Sciences*, 4(3): 333-344.
- 253 Abowei, J.F.N. (2010). Salinity, Dissolved Oxygen, pH and Surface Water Temperature  
254 Conditions in Nkoro River, Niger Delta, Nigeria. *Advance Journal of Food Science and*  
255 *Technology*, 2(1): 30 -40.
- 256 Adebowale, K.O., Agunbiade, F.O. and Olu-Owolabi, B.I. (2008). Impacts of natural and  
257 anthropogenic multiple source of pollution on the environmental conditions of Ondo  
258 State coastal water, Nigeria. *Journal of Environment, Agriculture and Food Chemistry*.  
259 7(4): 2798-2810.
- 260 Adeparusi, E.O., Ajibefun, A.I. and Akeremale, E.O. (2003). Smoke-Curing of Fish by Artisanal  
261 Fisher Folks in Ilaje, Ondo State, Nigeria. *UNAAB. ASSET Ser. A.*; 3(4):101-109.

- 262 Adirondack Ecologists, L.L.C. (2010). Schroon Lake Phytoplankton Monitoring Project.  
263 Schroon Lake Association, New York, USA.
- 264 Ajibare, A.O. (2014). Assessment of physico-chemical parameters of waters in Ilaje Local  
265 Government Area of Ondo State, Nigeria. *International Journal of Fisheries and Aquatic*  
266 *Studies*. 1(5): 84-92.
- 267 Akegbejo-Samson, Y. (1995). Ecology of the Fisheries Resources of Coastal Wetlands of Ondo  
268 State and Its Management Implications. Ph.D. Thesis, Federal University of Technology,  
269 Akure, 297.
- 270 APHA. (1998). Standard methods for the examination of water and waste water. 20 edition,  
271 American Public Health Association, Washington D.C. 1-47 pp.
- 272 Asaolu, S.S. (1998). Chemical pollution studies of coastal waters of Ondo State. *Doctor of*  
273 *Philosophy (Ph. D) Thesis*. Federal University of Technology, Akure, Nigeria. Pp: 15-17.
- 274 Balogun, K. J. and Ajani, E. K. (2015). Spatial and temporal variations of Phytoplankton  
275 pigments, Nutrients and Primary productivity in water column of Badagry Creek,  
276 Nigeria. *American Journal of Research Communication*. 3(7): 157-172
- 277 Bayode, O.J., Adewunmi, E.A. and Odunwole, S. (2011). Environmental Implications of Oil  
278 Exploration and Exploitation in the Coastal region of Ondo State, Nigeria: A regional  
279 planning appraisal. *Journal of Geographical and Regional Planning*. 4(3):110-121.
- 280 Begum, M., Hossain, M.Y., Wahab, M.A. and Kohinoor, A.H.M. (2003). Effects of iso-  
281 phosphorus fertilizers on water quality and biological productivity in fish pond. *Journal*  
282 *of Aquaculture in Tropics*. 18: 1-12
- 283 Benson, N. U., Essien, J. P. and Bassey, D. E. (2007). Hydrobiological constraints of trace  
284 metals in surface water, coastal sediment and water lily of Calabar River, Nigeria.  
285 *African Journal of Biotechnology*. 6 (20): 2358-2362.
- 286 Bolarinwa, J. B., Fasakin, E. A. and Fagbenro, A. O. (2016). Physicochemical analysis of the  
287 coastal waters of Ondo State, Nigeria. *International Journal of Research in Agriculture*  
288 *and Forestry*. 3(11): 13-20.
- 289 Boyd, C. E. (1979) Water quality in warm water fish pond. Auburn University Agricultural  
290 Experimental Station, Auburn, Alabama, USA
- 291 Boyd, C.E. (1982). Water Quality Management for Pond Fish Culture. 1st Edn., Elsevier  
292 Scientific Publishing Company, Amsterdam, Oxford, New York, Pages: 318.
- 293 Cako, V., Baci, S. and Shena, M. (2013). Water Turbidity as One of the Trophic State Indices in  
294 Butrinti Lake. *Journal of Water Resource and Protection*. 5: 1144-1148
- 295 Daramola, J.O., Adekunle, M.F., Olaniyi, M.O. and Alayaki, F.M. (2009). Diagnostic survey  
296 report of Ondo State agricultural production. Institute of Food Security, Environmental  
297 Resources and Agricultural Research, University of Agriculture, Abeokuta. P: 1-171
- 298 Emmanuel, B E, and Onyema, I.C. (2007): The Plankton and Fishes of a Tropical Creek in South  
299 – western Nigeria. *Turkish Journal of Fisheries and Aquatic Sciences*. 7 (2):105 – 113
- 300 Gharib, S., Radwan, Ahmed. And Abdel-Haim, A. (2011). Phytoplankton and environmental  
301 variables as a water quality indicator for the beaches at Matrouh, south-eastern  
302 Mediterranean Sea, Egypt: An assessment. *Oceanologia*. 53(3): 820-838
- 303 Harris, J.M. and Vinobaba, P. (2012). Impact of Water Quality on Species Composition and  
304 Seasonal Fluctuation of Planktons of Batticaloa Lagoon, Sri Lanka. *Journal of Ecosystem*  
305 *Ecography* 2:4-12

306 Imoobe T.O.T. (2011). Diversity and Seasonal Variation of Zooplankton in Okhuo River, a  
307 Tropical Forest River in Edo State, Nigeria, *Centrepoint Journal (Science Edition)*. 17(1):  
308 37-51.

309 Margalef, R. (1968). *Perspective in Ecological Theory*, University of Chicago Press. Pp: 112.

310 Nkwoji J.A., Yakubu A., Ajani G.F., Balogun K.J., Renner K.O. and Igbo J.K. (2010): Seasonal  
311 variations in water chemistry and benthic macro invertebrates of a South Western  
312 Lagoon, Lagos, Nigeria. *Journal of Amplified Science*. 6(3):85-92.

313 Offem, B.O., Ayotunde, E.O., Ikpi, G.U., Ada F.B. and Ochang S.N (2011): Plankton-Based  
314 Assessment of the Trophic State of Three Tropical Lakes. *Journal of Environmental*  
315 *Protection*. 2: 304-315.

316 Olawusi Peters, O. and Akinola, J. O. (2017). Assessment of Heavy metals concentration in  
317 some coastal waters of Ondo state, Nigeria. Proceedings of the 5<sup>th</sup> annual Conference  
318 School of Sciences, Federal University of Technology, Akure, Ondo State, Nigeria.  
319 SOS/CS/002; 175-184.

320 Olawusi-Peters, O. O., Akinola, J. O. and Jelili, A. O. (2017). Assessment of Heavy Metal  
321 Pollution in Water, Shrimps and Sediments of Some Selected Water Bodies in Ondo  
322 State. *Journal of Researches in Agricultural Sciences*. 5(2): 55-66

323 Olawusi-Peters, O.O. and Ajibare, A.O (2014). Species richness, diversity and abundance of  
324 some Decapod Crustaceans In coastal waters of Ondo State, South West, Nigeria.  
325 *International Journal of Fauna and Biological Studies*. 1 (5): 44-51.

326 Olawusi-Peters, O.O., Ajibare A.O. and Bello-Olusoji, O.A. (2014). Length-weight relationship  
327 and condition factor of shrimps in coastal waters of Ondo state, South West, Nigeria.  
328 *International Journal of Fisheries and Aquatic Studies*. 1(3):137-142.

329 Onyema, I.C. and Ojo, A.A. (2008). The zooplankton and phytoplankton biomass in a tropical  
330 creek, in relation to water quality indices. 5 (4): 75-82,

331 Pielou, E.C. (1966): Species diversity and pattern diversity in the study of ecological succession.  
332 *Journal of Theoretical Biology*. 10:370–383.

333 Shannon CE, Weaver W. (1963). *The mathematical theory of communication*. University Illinois  
334 Press, Urbana, IL.

335 Smith D, Johnson K.B and Kendall-Hunt (1996). *A Guide to Marine Coastal Plankton*, rev.

336 UNESCO. (1978). *Phytoplankton manual*: ISBN 92-3-101572-9

337 Varadharajan, D. and Soundarapandian, P. (2015). Biodiversity and Abundance of  
338 Phytoplankton from Muthupettai Mangrove Region, South East Coast of India. *Journal of*  
339 *Aquaculture Research and Development* 6:383-389

340 Yamaguchi, E. and Gould, A. (2007). *Phytoplankton Identification Guide*. The University of  
341 Georgia Marine Education Center and Aquarium.

342