

# Original Research Article

## The Relationship of Megazoobenthos Association With Seagrass Ecosystem Conditions In Ujung Genteng Waters, Sukabumi

### ABSTRACT

Seagrass beds in Ujung Genteng waters are one of the shallow marine ecosystems that have relatively high biodiversity. Seagrass ecosystem as one of the coastal ecosystems in these waters has a diversity of marine life that is associated with it. The purpose of this study was to establish the association pattern and the correlation of megazoobenthos to seagrass ecosystem conditions. The method used was a survey method by conducting a survey direct observation of the stations that have been determined. Three seagrass species were identified, namely: *Thalassia hemprichii*, *Cymodoceae rotundata* and *Enhalus acoroides*. The highest density of seagrass was found in station 2 with the type of seagrass *Thalassia hemprichii* as much as 207 ind/m<sup>2</sup>. Fifteen megazoobenthos species were identified representing 3 phyla. Species Megazoobenthos which is most commonly found in Ujung Genteng waters is *Ophiothrix fragilis*, *Ophiocoma erinaceus*, and *Diadema setosum*. The megazoobenthos Diversity Index at the observation station is included in the medium category that is 2.33 - 2.66 and the evenness index is 0.63 - 0.84. Correlation of seagrass density with megazoobenthos obtained by 0.92 shows that there is a very strong relationship between seagrass density with megazoobenthos. The result shows that there are 4 patterns of close association and 2 patterns of very close association. *Ophiocoma erinaceus* of the ophiuroidea class has a pattern of very close positive associations with seagrass *Cymodoceae rotundata* and *Dendrodoris sp* of the order nudibranchia has a pattern of very close negative associations with seagrass *Cymodoceae rotundata*.

**Keywords:** *Cymodoceae rotundata*, *Ophiocoma erinaceus*, Diversity Index, Association, and Correlation .

### 1. INTRODUCTION

Ujung Genteng is a coastal area, south of West Java, that is included in the administrative area of Sukabumi Regency. The waters of the tile end support the life of marine organisms such as megazoobenthos and seagrass. According to Badan Pengendalian Lingkungan Hidup Daerah Provinsi Jawa Barat [5], there is a seagrass ecosystem in the waters of the tip of the tile which is dominated by *Thalassia sp*.

Seagrass beds are complex shallow-water ecosystems, having high biological productivity. It is therefore an important marine resource both ecologically and economically [2]. The bioecological functions of seagrass beds include nurseries, spawning areas, foraging areas, and areas to seek protection for various types of marine biota such as fish, crustacea, mollusks, echinoderm, and etc [2]. Waters that have seagrass ecosystems have a higher diversity and abundance of associated organisms

when compared to waters where there are no seagrass ecosystems. It is for this reason that seagrass ecosystem management is needed to maintain the productivity of fisheries resources and the preservation of natural resources in coastal areas [15].

Megazoobenthos that is animal organism that is more than 1 cm in size that live on or in the seabed, including sticking, crawling and melting biota [11]. Megazoobenthos is an organism whose survival depends on seagrass ecosystems. Megazoobenthos which will be examined in this study is divided into three categories such as echinoderms, mollusks, and crustaceans. The high megazoobenthos diversity is influenced by the quality and condition of the seagrass ecosystem itself, meaning that the better the condition and quality of the seagrass ecosystem, the higher the diversity of the megazoobenthos [15]. Considering the important bioecological function and the observation of megazoobenthos organisms in seagrass beds and the study of its existence is still not much, this study still needs to be done.

## 2. MATERIALS AND METHODS

The study was conducted in May 2019 - June 2019. Data were taken 3 times every 2 weeks during the study. The research location is at Ujung Genteng Beach, Sukabumi. The research method used was a survey method. The sampling used in research by *purposive sampling*. Sampling and sample observation were carried out at 3 predetermined locations. Station 1 has medium seagrass cover conditions, station 2 has high seagrass cover conditions, and station 3 has very high seagrass cover conditions (Figure 1).




Fig. 1. Research location map

### 2.1 Research Procedure

Observation of seagrass samples was carried out by stretching three-line transects in a position perpendicular to the coastline towards the edge  $\pm 50$  meters, on each line, three plots were laid. In each observation plot, a 1 x 1 meter squared transect is divided into 25 sections with a size of 20 cm x 20 cm each. The distance between the transect squares ranges from 25 meters or adjusted to the existing seagrass area. Megazoobenthos observations were carried out together with recording the type and number of seagrass stands. The identification of megazoobenthos is done by matching the sample of

megazoobenthos to the picture contained in the megazoobenthos identification book (Marine Biology Book). To support field data, physical parameters are also measured and chemical measurements of physical parameters are temperature, transparency, depth, and current. While the chemical parameters are DO, pH, salinity, and nutrient content in the substrate. Measurement of temperature, DO, pH, and salinity using the potentiometric method, measurement of depth and transparency using visual methods, measurement of current using secondary data accessed from <https://www.ecmwf.int> and measurement of nutrients in sediments carried out by taking 250 grams of sediment then brought to the laboratory and measured by the spectrophotometric method.

## 2.3 Data Analysis

Seagrass biota is analyzed by calculating Species Density (Di), Relative Density (RDi), Species Frequency (Fi), Relative Frequency (RFi), Coverage Species (Ci), Relative Closure (RCi) and Importance Value Index (IVI) [6]. Whereas the megazoobenthos biota was analyzed by calculating the Abundance Index (K), Diversity Index (H'), and the Evenness Index [14]. Megazoobenthos abundance relationship with seagrass density calculation using the correlation coefficient *Pearson* (*Pearson's Product Moment of Correlation*). And the association between species of seagrasses and between species of megazoobenthos uses the calculation *presence-absence* or contingency table and then proceed with the Chi-square test  $\chi^2$ .

## 3. RESULTS AND DISCUSSION

### 3.1 General Condition of Research Location

Treatment station at the study site has different regional characteristics. Station 1 is at the coordinate point 7° 21 '33' 'LS - 106° 24' 10 " East. Station 1 has a seagrass cover condition with sediment texture included in the Sand category with a composition of 92% sand, 3% dust, and 5% clay. Station 2 is at coordinates 7° 22 '4' 'LS - 106° 24' 1 " BT. Station 2 has a very high seagrass cover condition with sedimentary textures including the Sand category with a composition of 90% sand, 4% dust, and 6% clay. Station 3 is at the coordinate point 7° 14 '20' 'LS - 106° 31' 7 " East. Station 3 has a high seagrass cover condition with sediment textures including the Sand category with a composition of 92% sand, 3% dust, and 5% clay.

#### 3.1.1 Physical and Chemical Parameters Of The Waters

Based on Table 1 the temperature obtained in Ujung Genteng waters is 28.5- 31.6°C. The temperature needed for seagrass to photosynthesis around 28-35°C. As for growing, seagrasses require optimal temperatures between 28-30°C [10].

**Table 1. Physicochemical Parameters**

No	Physicochemical Parameters	Research Station		
		I	II	III
1	Temperature (°C)	28.5-31.6	28.8-31.2	28.3-31.4
2	pH	6, 8-7.5	6.4-7.4	6.8-7.5
3	Salinity (‰)	31-33	31-33	31-33
4	DO (mg / L)	6.3-7.5	7- 8.8	6.5-8.2
5	Depth (cm)	0-100	0-100	0-100
6	Transparency (%)	100%	100%	100%

The results of pH measurements obtained in Ujung Genteng waters range from 6.4 to 7.5. The pH value in the Ujung Genteng waters can still be tolerated for seagrasses and megazoobenthos organisms. According to Kepmen LH No. 51 [10] concerning the quality standard of acidity (pH) in seagrass ecosystems ranges from 7 - 8.5. Water conditions that are very acidic or very basic will be detrimental to marine organisms.

Seagrass species have different tolerance to salinity. However, in general, the range of salinity that can be tolerated by seagrass is 10–40‰ and the optimum value is 35‰ [17]. The tolerable range of megazoobenthos, on the other hand, is 15-35‰. Salinity measurement results obtained in Ujung Genteng waters are 31-33‰. The salinity value is classified as good for growing seagrass species and living species megazoobenthos.

The results of DO measurements in Ujung Genteng waters range from 6.3-8.8 (mg / L), the condition is quite good for growing seagrass species. According to the Decree of the Kepmen LH No 51 [10] the optimum seagrass species grow in waters with DO values > 5 (mg/L). The dissolved oxygen conditions above can also support the life of megazoobenthos and other marine organisms. Almost all marine organisms like the condition of dissolved oxygen concentration > 5 mg/L [8].

The depth of the water ranges from 0-100 cm and the brightness is constant at each observation, which is 100% shows the penetration of sunlight into the bottom of the water.

Current data are taken consist of May 2019 to June 2019. The maximum current speed in May is 0.219 m / s and the minimum current speed is 0.072 m / s. The maximum current speed in June is 0.174 m / s and the lowest current value is 0.07 m / s. The results of the current velocity obtained in Ujung Genteng waters are of moderate current because the average current velocity in the east season is still below 0.5 m/s [16].

**Table 2. Analysis of Nutrient Content and Substrate Type**

Station	C-org (%)	N-total (%)	C / N	P <sub>2</sub> O <sub>5</sub> (ppm)	Texture (%)			Type Substrate
					S	Si	C	
I	0.52	0.07	7	10.42	92	3	5	Sand
II	0.94	0.10	10	9.19	90	4	6	Sand
III	0.81	0.11	7	12.01	91	3	5	Sand

Based on table 2, the highest Nutrient content is at station 2 and the lowest is at station 1. The high nutrient content at station 2 causes this station to have a high seagrass rate. Nutrients are important elements of seagrass growth and metabolism. The nutrient content is used as a benchmark for aquatic fertility because of the more optimal the nutrient content of water the more abundant the growth of seagrass [13].

### 3.2 Seagrass

#### 3.2.1 Composition of Seagrass Species

In Ujung Genteng waters, 3 seagrass species were found for all stations. At station 1 found 2 species of seagrasses, *T. hemprichii* and *C. rotundata*. At stations 2 and 3 found 3 species of seagrasses, *T. hemprichii*, *C. rotundata* and *E. acoroides*.

**Table 3. Seagrass Species in Ujung Genteng Waters**

No	Seagrass Type	Station		
		1	2	3
1	<i>Thalassia hemprichii</i>	√	√	√
2	<i>Cymodoceae rotundata</i>	√	√	√
3	<i>Enhalus acoroides</i>	-	√	√

### 3.2.2 Density of Seagrass

In all stations *T. hemprichii* had the highest density value than other types of seagrass this is caused because *T. hemprichii* has a good adaptation to environmental conditions. The highest density of seagrass is at station 2, namely *T. hemprichii* with a density of 207 ind/m<sup>2</sup> and the lowest at station 1 is *E. acoroides* with a density of 0 ind/ m<sup>2</sup> which means that the species is not found in station 1 (Tabel 4).

### 3.2.3 Frequency of Seagrass

Based Tabel 4, at station 1 *T. hemprichii* has a frequency value of 0.9 and *C. rotundata* has a frequency value of 0.8 which means that both species have fairly even distribution in the observation area of station 1. Whereas for the *E. acoroides* species the frequency value is 0 this indicates that this species was not found at all at station 1. At stations 2 and 3 *T. hemprichii* and *C. rotundata* have a frequency value of 1 this shows that in the observation plot area these two species are always found and can be said to be spreading seagrass species *T. hemprichii* and *C. rotundata* is evenly distributed at stations 2 and 3. For *E. acoroides* species the frequency values obtained are 0.4 at station 2 and 0.3 at station 3 this shows that the *E. acoroides* species are found at stations 2 and 3 but their distribution is uneven or it could be said in the area of the observation plot that only a few plots of *E. acoroides* were found.

### 3.2.4 Coverage of seagrass

Based on Tabel 4, station 1 has a total seagrass cover of 45% which is classified as a low seagrass coverage category, Station 2 has a total seagrass cover of 95% classified as a seagrass coverage category is very dense, while Station 3 has a total seagrass cover of 75% classified as a dense category. Seagrass coverage in the low category has a cover range of 26-50%, dense seagrass coverage has a cover range of 51-75%, and seagrass coverage in the very dense category has a cover range of 76-100% [9].

### 3.2.5 Important Value Index

*T. hemprichii* had the highest importance value index value of the species *C. rotundata* and *E. acoroides* that is equal to 1.8 and belong to the category of being, even though belonging to the category of being a role *T. hemprichii* effect on other seagrass species. This is evidenced by the existence of *T. hemprichii* which is almost found in all stations and has a high seagrass density (Tabel 4).

**Table 4. Seagrass Density, Frequency, Coverage, and Important Value Index**


Stasiun	Seagrass Species	Seagrass density (ind/m <sup>2</sup> )	Seagrass Frequency	Seagrass Coverage (%)	Important Value Index
Station I	<i>Thalassia hemprichii</i>	43	0.9	30%	1.8
	<i>Cymodoceae rotundata</i>	19	0.8	15%	
	<i>Enhalus acoroides</i>	0	0	0%	
Station II	<i>Thalassia hemprichii</i>	207	1	60%	1.1
	<i>Cymodoceae rotundata</i>	81	1	30%	
	<i>Enhalus acoroides</i>	3	0.4	5%	
Station III	<i>Thalassia hemprichii</i>	140	1	48%	0.1
	<i>Cymodoceae rotundata</i>	77	1	25%	
	<i>Enhalus acoroides</i>	1	0.3	2%	

### 3.3 Megazoobenthos

Megazoobenthos found at the study site contained 15 species of 3 phyla, Echinodermata, mollusca, and arthropoda. There are 6 species of phylum echinoderms, namely *Diadema*

*setosum*, *Tripneustes depressus*, *Tripneustes gratilla*, *Ophiothrix fragilis*, *Ophiocoma erinaceus*, and *Holothuria atra*. There are 3 species of crustacean species, *Squilla mantis*, *Calappa hepatica* and *Scopimera globosa*. There are 6 species of phylum mollusk, namely *Dendrodoris sp*, *Turbo sp*, *Chlamys sp*, *Anadara antiquate*, *Cypraea ventriculus*, and *Cypraea annulus*.

*O. erinaceus* and *O. fragilis* are the most abundant megazoobenthos found at the research location. As many as 46% of *O. erinaceus* was found at the study site and as many as 20% *O. fragilis* was found at the study site (Figure 6).



**Fig. 6. Megazoobenthos Species in Ujung Genteng Waters**

### 3.3.1 Abundance, Diversity, and Evenness Indeks Megazoobenthos

Table 5. Abundance, Diversity, and Evenness Indeks Megazoobenthos

Station	Abundance (K)	Diversity (H')	Evenness (E)
I	4 ind/m <sup>2</sup>	2.66	0.84
II	9 ind/m <sup>2</sup>	2.33	0.63
III	7 ind/m <sup>2</sup>	2.41	0.70

The Station 2 had the most abundant species at 9 individuals/m<sup>2</sup>. This is because the density of seagrass at station 2 is very high density so that food sources for several types of megazoobenthos are abundant at station 2, megazoobenthos species that have a high number of individuals at station 2 namely from the species *O. fragilis* and *O. erinaceus* (Table 5).

Station 1 has a diversity index value of 2.66, Station 2 has a diversity index value of 2.33, and Station 3 has a diversity index value of 2.41. The Diversity index of each station includes the category of moderate diversity, this is according to Odum [14] statement that megazoobenthos has a moderate diversity index of  $1 < H' < 3.00$  (Table 4), diversity that is showing the distribution of the number of individuals per moderate species, moderate community stability and moderate productivity. The highest diversity is found in station 1 and the lowest diversity is found in station 2. Even though the number of species of station 2 is greater but the abundance of individuals in each species at station 1 is better than station 2. According to Minarni et. al [12] states that the high and low diversity index values can be caused by various factors, including the number of types or

individuals obtained and the presence of several species found in quantities that are more abundant than other types. According to Odum [14] species diversity is not only synonymous with many types, but the nature of the community is determined by the number of species and the abundance of individuals of each type.

The evenness index (E) (Table 5) ranges from 0-1. According to Odum [14] If the value is close to 0, it means that the uniformity is low because there is a type that dominates, and if it is close to 1, a high evenness indicates that no species dominates. At station 1 it has an evenness index of 0.84, station 2 has an evenness index of 0.63, and station 3 has an evenness index of 0.70. In Ujung Genteng waters have a high population evenness because of the value of  $E > 0.6$ . A high evenness index indicates that there is a stable community. This is due to the distribution of individual megazoobenthos which are still the same between stations. The highest evenness index was found at station 1 because this was not as much at the station 1 species found in other stations other than that the number of individuals found was not much different from other individuals. The lowest evenness index is found at station 2, it is caused at station 2 which species are found more and the number of individuals found between species is uneven. One of the species with the highest number of individuals found at station 2 is *O. erinaceus*. The high number of *O. erinaceus* individuals causes the uniformity index at Station 2 to be the lowest among other stations.

### 3.4 Relationship of Seagrass Vegetation with Megazoobenthos

Correlation test results obtained a correlation value of 0.92. The correlation value includes the category of very strong relationship, according to Zaid [18] correlation with the value of 0.00 - 0.199 including a very low relationship, 0.20 - 0.399 including a low relationship, 0.40 - 0.599 including a moderate relationship, 0.60 - 0.799 including a strong relationship, and 0.80 - 1,000, including a very strong relationship. The relationship between the density of seagrass vegetation and the abundance of megazoobenthos obtained shows a directly proportional relationship. The higher the density of seagrass, the higher the abundance of living megazoobenthos, this occurs because the density of seagrass vegetation affects the increased production of litter and available nutrients. Seagrass also has many bioecological functions that are very important as a spawning area and care for various types of marine organisms such as megazoobenthos [15].

### 3.5 Associations of Megazoobenthos with Seagrass

The association pattern of megazoobenthos species with seagrass species was found as many as 38 pairs of species with positive associations and 7 pairs of negative associations (Table 6).

**Table 6. Patterns of association megazoobenthos species with seagrass species**

Species Seagrass	Species Megazoobenthos														
	Ds	Td	Tg	Of	Oe	Ha	Sm	Ch	Sg	De	Tu	Ch	Aa	Cv	Ca
Th	+	+	+	+	+	+	+	+	-*	+	+	+	+	+	+
Cr	+	+	+	+	+**	+	+	+	+	-**	+	+	+	+	+
Ea	-	+	+	+	+*	+	-	+	-	-	-	+*	+	+	+*

\* = close associations \*\* = very close associations

Seagrass : Th : *Thalassia hemprichii*, Cr : *Cymodocea rotundata*, Ea : *Enhalus acoroides*

Megazoobenthos : Ds : *Diadema setosum*, Td : *Tripneustes depressus*, Tg : *Tripneustes gratilla*, Of : *Ophiolithrix fragilis*, Oe : *Ophiocoma erinaceus*, Ha : *Holothuria atra*, Sm : *Squilla mantis*, Ch : *Calappa hepatica*, Sg : *Scopimera globosa*, De : *Dendrodoris sp*, Tu : *Turbo sp*, Ch : *Chlamys sp*, Aa : *Anadara antiquate*, Cv : *Cypraea ventriculus*, dan Ca : *Cypraea annulus*

**Table 7. Patterns of association megazoobenthos species with seagrass species**

No	Species Megazoobenthos	Species Lamun								
		<i>Thalassia hemprichii</i>			<i>Cymodoceae rotundata</i>			<i>Enhalus acoroides</i>		
		a	E (a)	$\chi^2$ count	a	E (a)	$\chi^2$ count	a	E (a)	$\chi^2$ count
1	<i>Diadema setosum</i>	27	26.0	1.56	26	25.6	0.16	6	8.00	1.07
2	<i>Tripneustes depressus</i>	12	11.6	0.54	11	11	0.02	6	3.56	2.80
3	<i>Tripneustes gratilla</i>	11	10.6	0.49	11	10	1.02	4	3.26	0.28
4	<i>Ophiothrix fragilis</i>	58	57.8	0.09	57	55.6	1.96	18	17.78	0.02
5	<i>Ophiocoma erinaceus</i>	64	63.6	0.45	65	61.1	18.04**	23	19.56	4.66*
6	<i>Holothuria atra</i>	18	17.3	0.89	18	16.7	1.85	7	5.33	0.95
7	<i>Squilla mantis</i>	7	6.7	0.29	7	6.5	0.61	1	2.07	0.87
8	<i>Calappa hepatica</i>	14	13.5	0.89	14	13.0	0.52	8	4.15	6.14*
9	<i>Scopimera globosa</i>	4	4.8	3.97*	5	4.6	0.43	0	0.00	0.00
10	<i>Dendrodoris sp</i>	16	15.4	0.77	10	14.9	28.22**	0	0.00	0.00
11	<i>Turbo sp</i>	18	17.3	0.89	18	16.7	1.85	2	5.33	3.81
12	<i>Chlamys sp</i>	9	8.7	0.39	9	8.3	0.81	4	2.67	1.07
13	<i>Anadara antiquate</i>	10	9.6	0.44	10	9.3	0.91	3	2.96	0.00
14	<i>Cypraea ventriculus</i>	16	15.4	0.77	16,0	14.8	1.60	7	4.74	1.91
15	<i>Cypraea annulus</i>	11	5.8	0.25	6	5.6	0.52	4	1.78	4.26*

\*  $\chi^2$  table with a standard error of 5% = 3.84

Megazoobenthos in the phylum Echinoderms occurs positive associations between the species of megazoobenthos with seagrass species. Species of the Echinoderms that are positively associated with seagrasses are sea urchins (*Diadema setosum*, *Tripneustes depressus*, *Tripneustes gratilla*). This confirms the idea suggested by Azkab [4] that sea urchins are one of the organisms known to be active in eating seagrasses, both those observed in nature and research conducted in the laboratory. However, there is one species of sea urchins that are negatively associated with seagrasses, namely *Diadema setosum* species with species *E. acoroides*. This condition is caused by *E. acoroides* having low density and also an uneven distribution of *E. acoroides* can only be found at station 2 and station 3 which causes the association of *Diadema setosum* to *E. acoroides* is negative. In addition, other species that positively associate with the Echinoderms phylum are the snaking star species (*Ophiothrix fragilis* and *Ophiocoma erinaceus*) as well as the sea cucumber species (*Holothuria atra*). This is in accordance with the statement of Dissanayake & Stefansson [7] that active sea cucumbers live in seagrass ecosystems, sea cucumbers are deposit-eating organisms and will gather in seagrass that have high densities. This has been proven to be found in sea cucumbers only at station 2 and station 3 with high seagrass density.

The positive association of crustacean classes to seagrasses occurs in the species *Squilla mantis*, *Calappa hepatica* and *Scopimera globosa*. Aswandy [3] states, some active crustaceans live in seagrass ecosystems such as crustaceans, crustaceans, which prey on other small animals that live attached (epizoa) to leaves or other parts of seagrass.

Megazoobenthos in mollusk phylum occurs positive and negative associations with seagrass species, megazoobenthos type of mollusk phyla that are actively associated with seagrass species, namely from the gastropod and Bivalvia classes. Gastropods that are actively associated with seagrass species are *Dendrodoris sp*, *Turbo sp*, *Cypraea ventriculus*, and *Cypraea annulus*. This is proven by Azkab statement [4] that a food chain event occurred between a gastropod and a seagrass, seagrass litter was used as a food source. Bivalves that are actively associated with seagrass species are *Chlamys sp*, and *Anadara antiquate*. Seagrasses and bivalves are related, one of which has the same substrate type characteristics that serve as habitat. In addition, the association between seagrass and bivalves has a strong connection in the food cycle. Evidenced by the



statement of Allifah & Rosmawati [1] litter on seagrass will settle to the bottom of the waters which are then broken down by microorganisms that become bivalve food.

The calculation result  $\chi^2$  shows (Table 7) that there are 4 patterns of close association and 2 patterns of association are very tight consisting of 1 pattern of associations between species is strongly negative and 3 patterns of association of species are closely positive, 1 pattern of association is very close negative and 1 pattern of association is very close positive. The pattern of negative inter-species associations is very close, that is, the species of the nudibranchia *Dendrodoris sp* against seagrass species *C. rotundata*. It is suspected that *Dendrodoris sp* does not like the presence of seagrass species *C. rotundata* as evidenced by the presence of this species only at station 1 which has a cover medium seagrass and species density is *C. rotundata* not so abundant. The pattern of positive inter-species associations that are very close to *Ophiocoma erinaceus* against seagrass species is *C. rotundata* evidenced by the presence of *Ophiocoma erinaceus* and *C. rotundata* which are quite high and are found in all stations. *Ophiocoma erinaceus* is a class of snaking stars that is negative phototaxis, *Ophiocoma erinaceus* takes shelter in the species of seagrass *C. rotundata* which has a high density to avoid direct sunlight. In addition, *Ophiocoma erinaceus* utilizes litter from the leaves of seagrass *C. rotundata* as a food source.

#### 4. CONCLUSION

The relationship of seagrass density with megazoobenthos is directly proportional, which means the density of seagrass has an impact on the abundance of megazoobenthos in Ujung Genteng waters. The pattern of association between megazoobenthos and seagrass species is positive and negative. *Ophiocoma erinaceus* of the ophiuroidea class has a pattern of very close positive associations with seagrass *Cymodoceae rotundata* and *Dendrodoris sp* of the order nudibranchia has a pattern of very close negative associations with seagrass *Cymodoceae rotundata*.

#### REFERENCES

1. Allifah AN, Rosmawati T. Relationship between Seagrass Density and Bivalvia Density on the Coast of Ori Coast, District of Haruku Island. Journal of Biology Science & Education. 2018; 70 (1): 81-96.
2. Arifin, Jamaluddin J. Study of Condition and Potential of the Seagrass Ecosystem as a Marine Biota Care Area. Indonesian Journal of Aquatic Sciences and Fisheries. 2005; 12 (2): 73-79.
3. Aswandy I. Crustaceans as Consumers in Seagrasses. Oceana Journal. 2008; 33 (1): 1-9
4. Azkab, MH. The Role of Seagrass for Animal Associations. Oceana Journal. 2014; 39 (2): 49-54.
5. Badan Pengendalian Lingkungan Hidup Daerah Provinsi Jawa Barat. Laut Dan Pesisir. Status Lingkungan Hidup. Bandung; 2008.
6. Brower JE, Zar JH. Field and Laboratory Methods for General Ecology. WM Brown Company Publ. Dubuque Iowa; 1989.
7. Dissanayake DCT, Stefansson G. Habitat preferences of sea cucumbers: *Holothuria atra* and *Holothuria edulis* in the coastal waters of Sri Lanka. Journal of the Marine Biological Association of the United Kingdom. 2012; 92 (3): 581-590.

8. Effendi, H. Water Quality Study for Water Resources and management Environmental. Kanisius. Yogyakarta; 2003.
9. Hutomo M, Nonjti A. Padang Seagrass Monitoring Guide. COREMAP LIPI: Jakarta; 2014.
10. Keputusan Menteri Lingkungan Hidup No : 51 Tahun 2004 Tentang Baku Mutu Air Laut. Menteri Negara Lingkungan Hidup.
11. Meyer KS, Bergmann M, Soltwede TI. Interannual variation in the epibenthic megafauna at the shallowest station of the HAUSGARTEN observatory (790N, 60E). Germany; 2013
12. Minarni, Jahidin, Darlian L. Abundance of Gastropods in Seagrass Habitat in the Waters of Tongali Village, Siompu District. Ampibi Journal. 2016; 1 (2): 17-21.
13. Nabilla S, Hartati R, Nuraini RAT. Nutrient Relationship to Sediment and Seagrass Closure in Jepara Waters. Tropical Marine Journal. 2019; 22 (1): 42-48.
14. Odum PE. Fundamentals of Ecology. Yogyakarta; 1993. Gajah Mada University Press.
15. Riniatsih I, Munasik. The Diversity of Megabentos Associating in the Padang Seagrass Ecosystem Wailiti Waters, Maumere Sikka District, East Nusa Tenggara. Tropical Marine Journal. 2017; 20 (2): 55–59
16. Setyawan WB, Aditya P. Comparison of Ocean and South Coastal Oceanographic Characteristics of Java Island: Tides, Currents, and Waves. Proceedings of the III National Ocean and Fisheries Seminar 2017. Trunojoyo University Madura.
17. Yusuf, M., Y. Koniyo., C. Panigoro. 2013. Seagrass Diversity in Waters Around Dudepo Island, Anggrek District, North Gorontalo Regency. Scientific Journal of Fisheries and Maritime Affairs 1 (1): 18-25.
18. Zaid MA. Correlation And Regression Analysis. Seisric. Turkey; 2015