

LATE HAEMORRHAGIC DISEASE OF NEWBORN: CAN IT BE PREVENTED BY CHANGING PROPHYLAXIS POLICY?

Abstract

Background: Vitamin K deficiency can cause severe haemorrhage in the newborn and is an important cause of infant morbidity and mortality. HDN can be classified according to the time of presentation after birth into early (0–24 hours), classical (1–14 days) and late (2–12 weeks) HDN. Late HDN, which presents after the first week of life, mainly manifests as intracranial haemorrhage, depending upon the site and amount of bleeding, it either results in mortality or life long sequelae in the form of cerebral palsy and scar epilepsy with or without cognitive impairment. **Objective:** To determine the frequency of ICH about vitamin K deficiency and outcome in infants aged 2 to 24 weeks. **Materials and methods:** From 1 September 2017 to 30 September 2019 we retrieved the retrospective data of 8 patients with late HDN admitted to Bapuji Child Health Institute and Chigateri Government General Hospital, Davangere. **Results:** Six of eight cases with late HDN had an intracranial haemorrhage, of whom 5 patients died (62%), one ended up with neurological sequelae (12%) and 2 cases had an extracranial bleed. Out of these 8 cases, 5 had not received vitamin K at birth. **Conclusion:** For neonates on strict breastfeeding, despite some with vitamin K prophylaxis, some patients still may suffer from intracranial and extracranial bleeding due to late HDN. Therefore, a change in strategy in the form of making the paediatricians and Anganwadi workers working in subcenters to give vitamin K, who have been vaccinating the babies after birth, would increase the vitamin K coverage.

Introduction

Vitamin K deficiency can cause severe haemorrhage in the newborn and is an important cause of infant morbidity and mortality. HDN can be classified according to the time of presentation after birth into early (0–24 hours), classical (1–14 days) and late (2–12 weeks) HDN.^(1,2)

Newborn infants are at risk for HDN for the following reasons:^(2,3)

- (1) reduced bioavailability because of poor placental transfer of vitamin K and the relatively short half-life of the K1 liver stores;
- (2) reduced vitamin K content in breast milk compared with fortified cow's milk-based formula; and
- (3) reduced production of vitamin K because of immature or altered gut flora.

Because dietary intake is an infant's main source of vitamin K, exclusively breastfed infants have a higher risk for HDN than formula-fed infants. In infants, the plasma concentrations of all vitamin K dependent clotting factors are 40-60% of the adult values and slowly rise during infancy but can take up to 90 days to completely normalize even with adequate vitamin K stores. Late HDN, which presents after the first week of life, mainly manifests as intracranial haemorrhage, depending upon the site and amount of bleeding, it either results in mortality or life long sequelae in the form of cerebral palsy and scar epilepsy with or without cognitive impairment.

47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87

Hemorrhagic Disease of Newborn (HDN) was first described by Townsend in 1894. In a definitive study published in 1944, prophylactic vitamin K given at birth was shown to reduce HDN associated death by greater than fivefold in the first 2 weeks of life.⁽²⁾ With evidence mounting over the next two decades, the American Academy of Pediatrics⁽⁴⁾ stated in 1961 recommending that a single dose of vitamin K be given to all neonates shortly after birth, either 0.5-1 mg intramuscular(IM) or 1-2 mg oral.

In the era of prophylaxis, HDN has become rare, with most reported cases being classical or late-onset and occurring in infants who either did not receive adequate vitamin K prophylaxis at birth and are exclusively breastfed or who had an undiagnosed malabsorptive or hepatobiliary disorder. Early HDN is mainly because of the effect of maternal medications and can be effectively prevented by vitamin K at birth; when no prophylaxis is given, the rate of early HDN is 6-12%.⁽⁵⁾ Without prophylaxis, the incidence of classical HDN is as high as 1.7% of live births,⁽⁶⁾ whereas the incidence of late HDN ranges from 4.4 to 7.2 per 100,000 live births.⁽⁷⁾ When IM vitamin K prophylaxis is given at birth, the rate of late HDN ranges from 0.24 to 3.2 cases per 100,000 live births.⁽⁸⁻¹²⁾ ICH occurs frequently in cases of late HDN and can lead to significant morbidity and mortality. In a pooled analysis of 131 cases, 63% of late HDN presented with ICH, with 14% mortality and 40% long-term neurological morbidity among surviving infants.⁽¹³⁾

Materials and Methods

From 1 September 2017 to 30 September 2019, we retrieved the retrospective data of 8 patients with late HDN admitted to Bapuji Child Health Institute and Chigateri Government General Hospital, Davangere.

Patients with prematurity and perinatal asphyxia were all excluded from the study. The details regarding pregnancy and delivery such as the timing of presentation, place of birth (hospital or home), signs and symptoms, bleeding sites, underlying illness of the baby, laboratory results, management, outcomes, route of vitamin K administration at birth, types of feeding were all recorded.

The diagnostic criteria of late HDN were established by the following criteria:

- (a) bleeding in an infant after seven days of life;
- (b) normal levels of fibrinogen and platelet counts;
- (c) returning to normal levels of prothrombin time (PT) and activated partial thromboplastin time (aPTT) after vitamin K administrations which were both elevated before vitamin K administration.

88
89
90





Figure 1:



91
92
93
94

Figure 2: CT brain showing multiple foci of acute haemorrhages in bilateral frontal, left parietal and left temporal lobes with interventricular extension along with diffuse cerebral oedema noted




Figure 3:

Laboratory results included complete blood count, clotting profile, liver function tests and urine analysis. Cranial computerised tomography and/ or magnetic resonance imaging was performed to whom suspected with ICH. All cases were evaluated for possible complications for ICH. During the follow-up period; psychomotor assessments and neurological examinations were done in all patients.

Results

Table 1: Demographic status, prophylaxis and outcome of patients included in the study

Sex	Age	Feeding	Vitamin K prophylaxis	Delivery place	Additional morbidity	Outcome
F	6 months	EBF	No	Home	No	Died
F	1 month	EBF	No	Ambulance	No	Died
M	2 months	EBF	No	Ambulance	No	Quadriparesis
M	1 month, 15 days	EBF	Don't know	PHC	No	Died
M	1 month, 20 days	EBF	No	Home	No	DAMA

M	3 months	EBF	No	PHC	Biliary atresia	Died
F	2 months	EBF	Yes	District hospital	Biliary atresia	No sequelae
M	2 months	EBF	Yes	District hospital	Cholestasis	No sequelae

108
109
110

Table 2: Natural course of disease in 1-year analysis

Total duration	1 year
No of cases in 1-year analysis	8 (100.00%)
No of cases expired	5 (62.50%)
Neurological sequelae	1 (12.50%)
Recovered	2 (25.00%)
Not received Vitamin K at birth	5 (62.50%)

111

Discussion

112

113

114

115

116

117

118

119

The frequency of ICH due to late HDN in infants aged two to 24 weeks of age was six times more than reported in a study by Visser et al. in 2011.⁽⁹⁾ The rate of late HDN ranges from 4.4 to 7.2 cases per 100 000 births, based on reports from Europe and Asia.^(4-6, 10-13) The higher frequency of late HDN found by this study suggests the need to confirm these data and to check the efficiency of the prophylaxis programme.

120

121

122

123

124

125

126

127

128

129

European studies prove that intramuscular vitamin K1 prophylaxis (1 mg) is highly effective in preventing late HDN and demonstrated higher incidence of late HDN in babies who received oral vitamin K compared with those with parenteral vitamin K at birth.^(12, 14-17) Most patients were presented at the fifth week, with a male-to-female ratio of 1.8:1, and this was comparable with other studies.⁽¹⁵⁻¹⁸⁾ The high percentage of exclusive breastfeeding reported in our patients raised the possibility that exclusive breastfeeding could be a predisposing factor for HDN.^(19,20) This raises the problem of low concentration of vitamin K in human breast milk. The predisposition to vitamin K deficiency bleeding following exclusive breastfeeding is emerging as a matter of concern, especially in developing countries where exclusive breastfeeding is vigorously advocated to promote optimal health in the infant.

130

131

132

133

134

135

136

137

138

139

140

141

Other studies reported that 88% and 97% of their studied patients with late HDN were exclusively breastfed.⁽⁹⁻¹³⁾ On the one hand, several studies revealed that vitamin K deficiency is seen following prolonged antibiotic therapy and during severe diarrhoeal diseases, especially infectious ones and despite prophylaxis.^(21,22) On the other hand, antibiotic use and infections before the onset of the bleeding were not detailed in the reports that included a large series of patients with HDN.^(23,24) Despite the national guidelines, the ground reality is that vitamin K is not given to all babies and late HDN continues to be a problem in India. The possible reasons could be lack of availability of injections, lack of trained adequate staff in health facilities, accountability and still 20% rate of home deliveries (NFHS -4 ,2015 -16).

142 **Conclusion**

143

144 There should be a serious discussion as to how to improve the coverage of vitamin K at
145 birth to prevent late HDN. With the rate of institutional delivery is 78.9%, the rate of BCG
146 vaccination is 92%(NFHS -4 ,2015 -16).

147

148 For neonates on strict breastfeeding, despite some with vitamin K prophylaxis, some
149 patients still may suffer from intracranial and extracranial bleeding due to late HDN. Therefore, a
150 change in strategy in the form of making the pediatricians and Anganwadi workers working in
151 subcenters to give vitamin K ,who have been vaccinating the babies after birth ,would increase
152 the vitamin K coverage of vitamin K prophylaxis specially among those born at home and those
153 born at institutes but for the same reason missed the dose.

154

155 **References**

156

157 1. Williams MD, Chalmers EA, Gibson BE. The investigation and management of neonatal
158 haemostasis and thrombosis. *Br J Haematol.* 2002;119(2):295-309.

159 2. Townsend CW. The haemorrhagic disease of the newborn. *Arch Paediatr.* 1894;11:559-
160 565.

161 3. Lehmann J. Vitamin K as a prophylactic in 13,000 infants. *Lancet.*1944;243(6294):493-
162 494.

163 4. American Academy of Pediatrics, Committee on Nutrition. Vitamin K compounds and
164 the water-soluble analogues: use in therapy and prophylaxis in pediatrics. *Pediatrics.*
165 1961;28:501-507.

166 5. Sutor AH, von Kries R, Cornelissen EA, McNinch AW, Andrew M. Vitamin K
167 deficiency bleeding (VKDB) in infancy: on behalf of the ISTH Pediatric/Perinatal
168 Subcommittee. *Thromb Haemost.* 1999;81(3):456-461.

169 6. American Academy of Pediatrics, Committee on Fetus and Newborn.Controversies
170 concerning vitamin K and the newborn. *Pediatrics.*2003;112(1):191-192.

171 7. Centers for Disease Control and Prevention (CDC). Notes from the field: late vitamin K
172 deficiency bleeding in infants whose parentsdeclined vitamin K prophylaxisTennessee,
173 2013. *MMWR Morb Mortal Wkly Rep.* 2013;62(45):901-902.

174 8. von Kries R, Hachmeister A, Göbel U. Oral mixed micellar vitamin K for prevention of
175 late vitamin K deficiency bleeding. *Arch Dis Child Fetal Neonatal Ed.* 2003;88(2):F109-
176 F112.

177 9. Ijland MM, Pereira RR, Cornelissen EA. Incidence of late vitamin K deficiency bleeding
178 in newborns in the Netherlands in 2005: evaluation of the current guideline. *Eur J Pediatr.*
179 2008;167(2):165-169.

180 10. Takahashi D, Shirahata A, Itoh S, Takahashi Y, Nishiguchi T,Matsuda Y. Vitamin K
181 prophylaxis and late vitamin K deficiency bleeding in infants: fifth nationwide survey in
182 Japan. *Pediatr Int.*2011;53(6):897-901.

183 11. Busfield A, Samuel R, McNinch A, Tripp JH. Vitamin K deficiency bleeding after NICE
184 guidance and withdrawal of Konakion Neonatal: British Paediatric Surveillance Unit
185 study, 2006-2008.*Arch Dis Child.* 2013;98(1):41-47.

- 186 12. Laubscher B, Bänziger O, Schubiger G. Prevention of vitamin K deficiency bleeding
187 with three oral mixed micellar phylloquinone doses: results of a 6-year (2005-2011)
188 surveillance in Switzerland. *Eur J Pediatr*. 2013;172(3):357-360.
- 189 13. Loughnan PM, McDougall PN. Epidemiology of late onset haemorrhagic disease: a
190 pooled data analysis. *J Paediatr Child Health*. 1993;29(3):177-181.
- 191 14. Busfield A, McNinch A, Tripp J. Neonatal vitamin K prophylaxis in Great Britain and
192 Ireland: the impact of perceived risk and product licensing on effectiveness. *Arch Dis*
193 *Child*. 2007;92:754-758.
- 194 15. Sutor AH, Kunzer W. Time interval between vitamin K administration and effective
195 hemostasis. In: Suzuki S, Hathaway WE, Bonnar J, Sutor AH, eds. *Perinatal thrombosis*
196 *and haemostasis*. Berlin, Heidelberg, New York, London, Tokyo, Hong Kong: Springer;
197 1991.
- 198 16. Wariyar U, Hilton S, Pagan J, Tin W, Hey E. Six years' experience of prophylactic oral
199 vitamin K. *Arch Dis Child Fetal Neonatal* 2000; 82: 64-8.
- 200 17. Yncecik F, Hergeuner MO, Sangun E O, E Ozcan K, Altunbapak P. Intracranial
201 haemorrhage related with late haemorrhagic disease of the new born. *Erciyes Med J*
202 2007; 29:v31-4.
- 203 18. Pooni PA, Singh D, Singh H, Jain BK. Intracranial haemorrhage in late hemorrhagic
204 disease of the newborn. *Indian Pediatr* 2003; 40: 243-8.
- 205 19. Al Ghwass MM, Ahmed D. Prevalence and predictors of 6-month exclusive
206 breastfeeding in a rural area in Egypt. *Breastfeed Med* 2011; 6: 191-6.
- 207 20. Lane PA, Hathaway WME. Vitamin K in infancy. *J Pediatr* 1985; 106: 351-9.
- 208 21. Tasker RC. Head and Spinal Cord Trauma. In: Nichols DG, editor. *Roger's textbook of*
209 *pediatric intensive care*, 4th ed. Baltimore, MD: Lippincott Williams & Wilkins, 2008:
210 888-911.
- 211 22. Yoshikawa H, Yamazaki S, Watanabe T, Abe T. Vitamin K deficiency in severely
212 disabled children. *J Child Neurol* 2003; 18: 93-7.
- 213 23. Aydinli N, Citak A, Caliskan M, Karabocuoglu M, Baysal S, Ozmen M. Vitamin K
214 deficiency: late onset intracranial haemorrhage. *Eur J Paediatr Neurol* 1998; 2: 199-203.
- 215 24. Pirinccioglu AG, Gurkan F, Bosna Mk, Acemoglu H, Davutoglu M. Intracranial
216 hemorrhage: clinical and demographic features of patients with late hemorrhagic disease.
217 *Pediatr Int* 2011; 53: 68-71.
- 218