

Genetic variability in chickpea (*Cicer arietinum* L.) under heat stress condition

Sanjay Kumar^{*}, Suresh B.G., Anand Kumar¹ and G.R. Lavanya

Department of Genetics and Plant Breeding, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj, U.P.-211007; ¹Bihar Agricultural University, Sabour (Bhagalpur)-813210, India.

ABSTRACT

The present experiment was carried out at field experimentation centre of the Genetics and Plant Breeding, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (Uttar Pradesh) to study genetic variability, correlation and path analysis in fifty germplasm of chickpea during *rabi*, 2017-18. The maximum phenotypic coefficient of variation (PCV) and genotypic coefficient of variation (GCV) were noticed for 100-seed weight, biological yield per plant, grain yield per plant and number of primary branches per plant. High heritability were recorded by 100 seed weight, biological yield per plant, primary branches per plant, grain yield per plant, effective pods per plant, total number of pods per plant, secondary branches per plant, plant height, days to 50% flowering and days to maturity. High heritability coupled with high genetic advance as percent of mean was observed for 100 seed weight, biological yield per plant, primary branches per plant, grain yield per plant, effective pods per plant, total number of pods per plant, secondary branches per plant and plant height which suggested that these characters can be considered as favorable attributes for the improvement through selection. Path coefficient analysis for grain yield per plant revealed that biological yield, harvest index, secondary branches, canopy temperature at vegetative stage, effective pods per plant had given the highest contribution on yield per plant. So the utmost importance should be given to these characters during the selection for yield improvement in late sown chickpea.

Keywords: *Cicer arietinum* L., genetic variability, correlation, path analysis and grain yield

1. INTRODUCTION

Chickpea (*Cicer arietinum* L.) $2n = 2x = 16$, is the third most important food legume belongs to family *Fabaceae*. It is globally, occupying an area of 14.56 m ha with a production of 14.78 mt (FAO STAT, 2019). India is the largest producer of chickpea in the world sharing about 72% of area and production covering about 10.56 million ha area with annual production of 11.17 million tones grain. The present yield level is 1077 kg/ha which is far below the potential yield (5000 kg/ha) of the crop. In spite of India being the largest chickpea producing country, a deficit exists in domestic production and demand, which is met through imports. Uttar Pradesh is a traditional state for chickpea cultivation covering an area of 501.0 thousand ha with production of 578.66 thousand tones and productivity of 1155 kg/ha (Project Coordinator's Report, AICRP on Chickpea, 2018-19). About 90% of chickpea production occurs on residual soil moisture under rainfed conditions (Kumar and Van Rheenan 2000), where terminal heat stresses are major limitations to higher productivity (Johansen *et al.* 1997). Chickpea productivity is constrained by several biotic and abiotic stresses (Gaur *et al.* 2008) and temperature is one of the most important determinants of crop growth over a range of environments (Summerfield *et al.* 1990) and may limit chickpea yield (Basu *et al.* 2009). Chickpea is a cool season food legume and causes severe yield losses when exposed to high temperatures about 35⁰C during reproduction. Heat stress is increasingly becoming a major constraint for chickpea production in India because it is expected to increase the area under late sown conditions due to increased crop intensity or late rainy season crop maturity, and to increase overall temperatures due to climate change (Gaur *et al.*, 2008). Chickpea reproductive stages (flowering and podding) are vulnerable to external environmental changes and heat stress (Krishnamurthy *et al.*, 2011). Frequent decreases in the yields of chickpea seed were observed when plants were exposed to high (> 35⁰C) temperatures at flowering and pod development stages (Wang *et al.*, 2006). Approximately 11.7 million ha of rice area in India is estimated to currently remain fallow in central and north-eastern India after late rice harvest during the winter season (Subbarao *et al.*, 2001). These lands can offer expansion in the cultivation of chickpea provided genotypes are made available that is capable of standing heat stress. Due to climate change and global warming, heat stress is projected to be an increasingly important constraint in the near future. Through 2050, with higher levels of warming in northern parts of India, a temperature rise of at least 2⁰C, particularly night temperatures, is

expected. The predicted climate change will reduce the grain yield in chickpea, particularly at high temperatures. For example, chickpea yield declined by as much as 301 kg/ha per 1⁰C in India (Karla *et al.*, 2010). There is an urgent need to search for various sources of heat tolerance from the gene bank. In chickpea cultivars, heat tolerance is highly needed to achieve higher yields in all growing conditions that expose chickpea to high temperatures, especially during the reproductive stage. So, heat tolerance varieties are needed for improving chickpea yields in late sown conditions.

Yield is a complex character in general and more complex in chickpea, as it is depending upon a number of components. Heat tolerance is greatly needed in chickpea cultivars for realizing higher yields in all growing conditions that expose chickpea to high temperature, particularly at the reproductive stage. The genetic variability presents in the base population for desired characters play an important role in development of desirable plant type. Less information is present in the cultivated chickpea lines grown under heat stress conditions. Therefore, the present investigation was carried out to assess the genetic variability, association of different traits towards yield and selection of high yielding genotypes with better architecture under heat stress conditions.

2. MATERIALS AND METHOD

The experimental material comprised of fifty germplasm of chickpea were sown on 15th December 2017 to coincide heat stress with pollination in *rabi*, 2017-18 at field experimentation centre of the Genetics and Plant Breeding, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (Uttar Pradesh). Chickpea reproductive stages (flowering and podding) are vulnerable to external environmental changes and heat stress. Frequent decreases in the yields of chickpea seed were observed when plants were exposed to high (> 35⁰C) temperatures at flowering and pod development stages (Wang *et al.*, 2006). The meteorological data recorded during crop growing period, (Rabi 2017-18 at SHUATS, Prayagraj (Uttar Pradesh) are shown in Fig.-1. The experiment was laid in randomized complete block design with three replications during *rabi*, 2017-18 with inclusion of the recommended packages and practices needed for a healthy crop. Data for 14 quantitative traits were recorded viz; days to 50 % flowering, days to maturity, Chlorophyll content, Canopy temperature, plant height (cm), number of primary branches per plant, number of secondary branches per plant, number of total pods per plant, Effective pods/plant, biological yield per plant, harvest index, 100-seed weight and seed yield per plant. The days to 50% flowering and days to maturity were accounted on a plot basis

and rest of the characters was documented from random sample of five plants in each plot. Each genotype was planted in a single row of four meters length with a spacing of 30 cm between rows and 10 cm between plants. The recommended agronomic practices and crop protection measures were followed during the crop growth period. Biometrical methods were followed to estimate genotypic and phenotypic coefficient of variation (Burton 1952), heritability in broad sense (Lush 1940), genetic advance (Johnson *et al.*, 1955) and correlation and path coefficient analysis (Singh and Chaudhry, 1979).

3. RESULTS AND DISCUSSION

The analysis of variance revealed significant differences among the genotypes (Table 1) for all the **studied characters** which provide an opportunity for selecting suitable genotypes with better performance for the traits. A wide range of variability was noticed for different agronomic and economic traits. The estimates of phenotypic coefficient of variation (PCV) in general, were higher than the estimates of genotypic coefficient of variation (GCV) for all the characters, which suggested that the apparent variation is not only due to the genotypes but also due to the influence of environment (Table 2). The maximum genotypic coefficient of variation was observed in seed index followed by grain yield per plant, biological yield per plant and primary branches per plant. This is an indicative of less amenability of these traits to environmental fluctuations and hence, greater emphasis should be given to these characters, while breeding cultivars from the present material. High GCV for number of pods per plant and 100-seed weight were also earlier reported by Jeena *et al.* (2005), Younis *et al.* (2008), Alwani *et al.* (2010) and Babbar *et al.* (2012). The high PCV were recorded by 100-seed weight, grain yield per plant, biological yield per plant, effective pods per plant, primary branches per plant, total number of pods per plant and secondary branches per plant. The magnitude of PCV ranged from 4.44 (days to maturity) to 30.08 (100 seed weight). The characters with high phenotypic coefficient of variation indicated more influence of environmental factors. Therefore, caution has to be exercised during the selection program because the environmental variations are unpredictable in nature and may mislead the results. The heritability for most of the characters ranged from 42.74 (canopy temperature at pod filling stage) to 98.08 (100-seed weight). The heritability estimates were recorded high for 100 seed weight, biological yield per plant, primary branches per plant, grain yield per plant, effective pods per plant, total number of pods per plant, secondary branches per plant, plant height which, days to 50% flowering and days to maturity suggested that the characters are least influenced by the environmental factors and also indicates the dependency of

phenotypic expression which reflects the genotypic ability of cultivars to transmit the genes to their offspring's. Similar results were also reported by Bicer and Sarkar (2008) and Younis *et al.* (2008). The expected genetic advance was high for grain yield per plot while number of pods per plant showed moderate heritability. High estimates of heritability does not always mean high genetic advance. Johnson *et al.* (1955) suggested that heritability estimates and the genetic advance as percent of mean together would provide a better judgment rather than heritability alone in predicting the resultant effect of selection. High heritability coupled with high genetic advance over mean was observed for 100 seed weight, biological yield per plant, primary branches per plant, grain yield per plant, effective pods per plant, total number of pods per plant, secondary branches per plant and plant height which suggested that these characters can be considered as favorable attributes for the improvement through selection and this may be due to additive gene action and thus, could be improved upon by adapting selection without progeny testing. Similar results have also been reported by Yadav *et al.* (2003). The study of inter-relationship among various characters in the form of correlation is, in fact, one of very important aspects in selection programme for the breeder to make an effective selection based on the correlated and uncorrelated response.

Knowledge of nature and magnitude of associations among different characters are important on three counts. Indirect selection is important when desirable characters have low heritability measure in one sex only. The efficiency of indirect selection is measured as a correlated response (Falconer, 1960). Knowledge of correlation is required when selection is to be made on several characters at a time through some simultaneous selection model (Pandey *et al.*, 2017). Even if, the objective is to make selection on a single trait, the knowledge of correlation is essential to avoid the undesirable correlated changes in other characters. In general, magnitude of genotypic correlation was higher than their corresponding phenotypic correlation coefficients in most of the characters suggesting that a strong inherent association exists for the traits studied and phenotypic selection may be rewarding. Similar results were also reported by Pathak *et al.* (1986). Higher magnitude of genotypic correlation helps in selection for genetically controlled characters and give a better response for seed yield improvement than that would be expected on the basis of phenotypic association alone (Robinson *et al.*, 1951).

Grain yield per plant exhibited positive significant correlations with biological yield per plant, 100-seed weight, secondary branches/plant, total no. of pods/plant and chlorophyll index respectively at genotypic and phenotypic level. However, primary branches/plant exhibited significant and positive association at genotypic level only while effective

Pods/plant and harvest index at phenotypic level. Negative and significant correlations were observed with days to 50% flowering, days to maturity at both genotypic and phenotypic level while canopy temperature at pod filling stage, canopy temperature at vegetative stage showed significant and negative correlation at genotypic level only (Table 3). Kuldeep *et al.* (2014) suggested that yield could be raised by selecting for more number of pods per plant and plant stand, which is evident in the present study. Similar findings were reported by Telebi *et al.* (2007), Hahid *et al.* (2010) and Ali *et al.* (2011).

Path coefficient analysis (Table 4) for grain yield per plant revealed that biological yield, harvest index, secondary branches, canopy temperature at vegetative stage, effective pods per plant had highest positive direct effect towards seed yield. These results are in agreement with the earlier reports of Priti *et al.* (2003). It means a slight increase in any one of the above traits may directly contribute towards seed yield. Similar results were reported by Talebi *et al.* (2007) and Babbar *et al.* (2012).

Breeding strategies for improvement of yield potential in chickpea genotypes under heat stress would aim on selection of plants having high biological yield per plant, 100-seed weight, secondary branches/plant, total no. of pods/plant, chlorophyll index and using these associated characters may be useful to the breeder to formulate appropriate breeding plans for selection of the genotype which tolerate high temperature condition.

4. CONCLUSION

The above findings revealed that under heat biological yield, harvest index, secondary branches, canopy temperature at vegetative stage, effective pods per plant had showed the maximum contribution towards seed yield. On the basis of seed yield and its attributing traits Sabour chana-1, BRC-1084-127, BRC-1047-33 and ICCV15112 were identified as promising heat tolerant genotypes. The identified promising heat tolerant genotypes may be used as donor in chickpea hybridization programme. Breeding strategies for improvement of yield potential in chickpea genotypes under heat stress would aim on selection of plants having higher number of effective pods, biological yield, harvest index and 100-seed weight and using these associated characters may be useful to the breeder to formulate appropriate breeding plans for selection of the genotype which tolerate high temperature conditions.

Conference Disclaimer

Some part of this manuscript was previously presented and uploaded in the following conference : Pulses for Nutritional Security and Agricultural Sustainability, At BAU, Sabour Bhagalpur, India in January 2018: Web Link of the proceeding :

5. REFERENCES

- Ali Q, Ahsan M, Khaliq I, Elahi M, Shahbaz M, Ahmed W and Naees M** Estimation of genetic association of yield and quality traits in chickpea (*Cicer arietinum* L.). International Research Journal of Plant Science 2011, **2(6)**:166-169.
- Alwani H, Moulla M and Chouhan W** Genotype environment interaction and genetic parameter in Chickpea (*Cicer arietinum* L.) landraces. Journal of Agricultural Science 2010, **2(1)**: 153-157.
- Babbar A, Prakash V, Prakash T and Iquabal MA** Genetic variability of chickpea (*Cicer arietinum* L.) under late sown condition. Legume Research 2012, **35(1)**: 1-7.
- Basu PS, Ali M and Chaturvedi SK** Terminal heat stress adversely affects chickpea productivity in northern India- Strategies to improve thermo tolerance in the crop under climate change on Agriculture, 23-25 February, New Delhi, India, 2009 Pp. 189-193.
- Bicer BT and Sakar D** Heritability and path analysis of some economical characteristics in lentil. Journal of Central Europe Agriculture 2008, **9(1)**: 191-196.
- Burton GW** Quantitative inheritance of grasses. Proc. 6th Int. Grassland congress 1952, **1**: 277-283.
- FAOSTAT** Food and Agriculture Organization of the United Nations , Rome, Italy. 2019 <http://faoapps.fao.org/site/567/default.aspx>.
- Falconer DS** Introduction to Quantitative Genetics. Oliver and Boyd, Edinburgh 1960, pp.365.
- Gaur PM, Kumar J, Gowda CLL, Pande S, Siddique KHM, Khan TN, Warkentin TD, Chaturvedi SK, Than AM and Ketema D** Breeding chickpea for early phenology: perspectives, progress and prospects. In: Kharkwal MC (ed) Food Legumes for Nutritional Security and Sustainable Agriculture, Vol. 2, New Delhi, India: Indian Society of genetics and Plant Breeding, 2008, pp. 39-48.
- Hahid S, Malik R, Bakhsh A, Asif MA, Iqbal U and Iqbal SM** Assessment of genetic variability and interrelationship among some agronomic traits in chickpea. International Journal of Agricultural Biology 2010, **12(1)**: 81-85.
- IIPR Project Coordinator's Report** AICRP on Chickpea, 2018-19, by Dr. G.P. dixit, Kanpur,.

- Jeena AS, Arora PP and Ojha OP** Variability and correlation studies for yield and its components in chickpea. *Legume Research* 2005, **28**(2):146-148.
- Johnson HW, Robinson HE and Comstock RE.** Estimates of genetic and environmental variability in soyabean. *Agronomy Journal*, 1955, **47**:314-318.
- Kalra N, Chakraborty D, Sharma A, Rai HK, Jolly M, Chander S, Kumar PR, Bhadraray S, Barman D, Mittal RB, Lal M and Sehgal M.** Effect of temperature on yield of some winter crops in northwest India. *Current Science* 2008, **94**:82-88.
- Krishnamurthy L, Gaur PM, Basu PS, Chaturvedi SK, Tripathi S, Vadez V and Rathore A.** Large genetic variation for heat tolerance in the reference collection of chickpea (*Cicer arietinum* L.) germplasm. *Plant Genetics Research* 2011, **9**(1): 59–69.
- Kuldeep RK, Pandey S, Babbar A and Mishra DK.** Genetic variability, character association and path coefficient analysis in chickpea grown under heat stress condition. *Electronic Journal of Plant Breeding* 2014, **5**(4):812-819.
- Kumar J and Van Rheenen HA** A major gene for time of flowering in chickpea. *J. Heredity* 2000, **91**: 67-68.
- Lush J L** Intra-sire correlation and regression of offspring in rams as a method of estimating heritability of characters. *Proceedings of the America Philosophical Society on animal Product* 1940, **33**: 292-301.
- Priti G, Semwal BD and Gupta D** Correlation and path analysis in black gram (*Vigna mungo* L. Hepper). *Progressive Agriculture* 2003, **3**(12): 63- 65.
- Summerfield RJ, Virmani SM, Roberts EH and Ellis RH** Adaption of chickpea to agro climatic constraints. In: Van Rheenen, H.A., Saxena, M.C.(Eds.), *Chickpea in the Nineties*, Proceeding of the second International workshop on chickpea Improvement, 4-8th December 1989. ICRISAT Publishing, India 1990, pp. 50-61
- Subbarao GV, Kumar Rao JVDK, Kumar J, Johansen C, Deb UK, Ahmed I, Krishna Rao MV, Venkataratnam L, Hebbar KR, Sai MVSR and Harris D.** Spatial distribution and quantification of rice-fallow in South-Asia-Potential for legumes, ICRISAT, Patancheru, India, 2001.
- Talebi R, Fayaz, F and Jelodar NAB** Correlation and path coefficient analysis of yield and yield components of chickpea (*Cicer arietinum* L.) under dry land condition in the west of Iran. *Asian Journal of Plant Science* 2007, **6**(7):1151-1154.

Wang J, Gan YT, Clarke F and McDonald CL. Response of chickpea yield to high temperature stress during reproductive development. *Crop Science* 2006, **46**: 2171-2178.

Yadav JK, Singh HL and Kumar R Determining selection components in chickpea (*Cicer arietinum* L. Wilczek). *Plant Archives* 2003, **3**: 125-128.

Younis N, Hanif M, Sadiq S, Abbas G, Asghar MJ and Haq MA Estimates of genetic parameters and path analysis in lentil (*Lens culinaris* Medik). *Pakistan Journal of Agricultural Science* 2008, **45**(3): 44 - 48.

Pandey, S., Kureshi, S. P., & Bhatore, A. (2017). Studies on genetic variability and interrelationship among the different traits in exotic lines of lentil (*Lens culinaris* medik). *Plant Archives*, **17**(2), 1164-1170.

Fig.-1: Metrological data recorded during, rabi 2017-18




Table- 1. Analysis of variance for fourteen characters in fifty chickpea genotypes under heat stress condition

SI. NO.	Characters	Mean Sum of Squares Replication (d. f.=2)	Treatment (d. f.=49)	Error (d. f.=98)
1.	Days to 50% flowering	54.25	91.71**	4.90
2.	Days to maturity	6.21	48.27**	5,19
3.	Chlorophyll index	4.12	51.08**	13.83
4.	Canopy Temperature at Vegetative stage	0.38	7.87**	2.30

5.	Canopy Temperature at pod filling stage	9.53	16.15**	4.98
6.	Plant height (cm)	8.17	113.97**	11.32
7.	Primary branches per plant	0.21	0.46**	0.02
8.	Secondary branches per plant	0.03	2.30**	0.13
9.	Total no. of pods per plant	1.61	79.26**	8.90
10.	Effective pods per plant	0.61	38.49**	4.02
11.	Biological yield per plant	0.16	41.90**	1.22
12.	Harvest index	2.71	28.36**	3.93
13.	100-seed weight	0.06	124.77**	0.81
14.	Grain yield per plant	0.09	3.97**	0.40

* and ** Significant at 5% and 1% levels of significance, respectively.

Table 2: Estimates of genetic parameters for fourteen quantitative characters in fifty genotypes of chickpea under heat stress condition

SI. NO.	Characters	σ^2_g	σ^2_p	GCV	PCV	h^2 (bs) %	GA	GA as % of mean
1.	Days to 50% flowering	28.94	33.84	7.76	8.39	85.52	10.25	14.78
2.	Days to maturity	14.36	19.54	3.81	4.44	73.47	6.69	6.73
3.	Chlorophyll index	12.42	26.25	5.95	8.65	47.30	4.99	8.43
4.	Canopy Temperature at Vegetative stage	1.86	4.16	5.13	7.69	44.64	1.88	7.07

5.	Canopy Temperature at pod filling stage	3.72	8.71	5.22	7.98	42.74	2.59	7.03
6.	Plant height (cm)	34.22	45.54	13.56	15.65	75.14	10.44	24.22
7.	Primary branches per plant	0.15	0.17	21.19	22.26	90.72	0.76	41.59
8.	Secondary branches per plant	0.72	0.85	17.55	19.06	84.89	1.61	33.32
9.	Total no. of pods per plant	23.45	32.36	18.73	21.99	72.48	8.49	32.84
10.	Effective pods per plant	11.49	15.51	19.61	22.78	74.08	6.01	34.76
11.	Biological yield per plant	13.56	14.78	22.91	23.93	91.74	7.26	45.22
12.	Harvest index	8.14	12.07	8.41	10.25	67.45	4.83	14.23
13.	100-seed weight	41.32	42.13	29.79	30.08	98.08	13.11	60.78
14.	Grain yield per plant	1.19	1.59	20.14	23.30	74.70	1.94	35.85

σ^2_g =Genotypic variances, σ^2_p = Phenotypic variances, **GCV**=Genotypic coefficient of variation, **PCV**= Phenotypic coefficient of variation, **h²(bs)** = heritability in broad sense and **GA**=genetic advance.




Fig. 2: Histogram showing estimates of Genetic Parameters for fourteen quantitative characters in chickpea under heat stress condition

UNDER PEEH

Table 3: Genotypic and phenotypic correlation for 14 characters in fifty genotypes of chickpea under heat stress condition

Characters		Days to 50% flow.	Days to maturity	Chlorophyll index	CT @ VS	CT @ PFS	Plant height	primary branches plant ⁻¹	Sec. branches plant ⁻¹	Total no. of Pods plant ⁻¹	Effective pods plant ⁻¹	Biological yield plant ⁻¹	Harvest Index	100-seed weight	Grain yield plant ⁻¹
Days to 50% flowering	G	1.000	0.822**	-0.486**	0.491**	0.387**	0.373**	-0.119	0.020	-0.132	-0.073	-0.219**	-0.088	-0.041	-0.264**
	P	1.000	0.711**	-0.336**	0.313**	0.211**	0.310**	-0.120	0.027	-0.098	-0.049	-0.194*	-0.086	-0.039	-0.217**
Days to maturity	G	1.000	1.000	-0.433**	0.599**	0.384**	0.324**	0.064	0.159	-0.096	-0.130	-0.190*	-0.161*	-0.121	-0.261**
	P	1.000	1.000	-0.270**	0.332**	0.190*	0.211**	0.052	0.106	-0.094	-0.120	-0.159	-0.121	-0.100	-0.199*
Chlorophyll index	G			1.000	-0.165*	-0.429**	-0.147	0.280**	-0.006	-0.188*	-0.131	0.170*	0.183*	0.249**	0.255**
	P			1.000	-0.117	-0.163*	-0.136	0.183*	-0.010	-0.106	-0.087	0.131	0.097	0.164*	0.167*
CT @ VS	G				1.000	0.435**	0.094	-0.003	-0.036	-0.005	0.076	-0.175*	-0.084	0.149	-0.210**
	P				1.000	0.199*	0.114	0.051	0.011	-0.010	0.033	-0.098	-0.021	0.084	-0.090
CT @ PFS	G					1.000	-0.239**	-0.058	0.010	-0.202*	-0.147	-0.447**	-0.343**	-0.355**	-0.612**
	P					1.000	-0.124	0.052	-0.156	0.076	0.095	-0.177*	0.023	-0.238**	-0.149
Plant height	G						1.000	0.167*	-0.113	-0.039	-0.054	0.183*	-0.362**	0.299**	0.047
	P						1.000	0.129	-0.062	0.019	0.007	0.181*	-0.206*	0.263**	0.089
Pri. branches plant ⁻¹	G							1.000	0.442**	-0.091	-0.112	0.150	0.032	0.236**	0.176*
	P							1.000	0.438**	-0.090	-0.110	0.140	0.028	0.223**	0.149
Sec. branches plant ⁻¹	G								1.000	0.334**	0.201	0.276**	0.121	0.305**	0.405**
	P								1.000	0.358**	0.251**	0.306**	0.187*	0.263**	0.433**
Total no. of Pods plant ⁻¹	G									1.000	0.814**	0.308**	-0.184*	-0.068	0.280**
	P									1.000	0.862**	0.376**	0.097	-0.060	0.436**
Effective pods plant ⁻¹	G										1.000	0.134	-0.111	-0.064	0.117
	P										1.000	0.229**	0.138	-0.057	0.306**
Biological yield plant ⁻¹	G											1.000	-0.462**	0.569**	0.920**
	P											1.000	-0.256**	0.539**	0.895**
Harvest Index	G												1.000	0.185*	-0.091
	P												1.000	0.146	0.188*
100-seed weight	G													1.000	0.723**
	P													1.000	0.614**

* , ** Significant at 5% and 1% levels of significance, respectively;

CT@VS= Canopy temperature at vegetative stage; CT@PFS= Canopy temperature at pod filling stage

Table 4: Direct (diagonal) and indirect genotypic and phenotypic effects of different characters on grain yield in chickpea under heat stress condition

Characters		Days to 50% flowering	Days to maturity	Chloroph yll index	CT @ VS	CT @ PFS	Plant height	primary branches plant ⁻¹	Secondary branches plant ⁻¹	Total no. of Pods plant ⁻¹	Effective pods plant ⁻¹	Biological yield plant ⁻¹	Harvest Index	100- seed weight	Grain yield plant ⁻¹
Days to 50% flowering	G	0.011	0.009	-0.005	0.005	0.004	0.004	-0.001	0.0002	-0.0014	-0.001	-0.002	-0.001	-0.0004	-0.264
	P	-0.002	-0.001	0.001	-0.0005	-0.0004	-0.0005	0.0002	0.0000	0.0002	0.0001	0.0003	0.0001	0.0001	-0.217
Days to maturity	G	-0.007	-0.009	0.004	-0.005	-0.003	-0.003	-0.0005	-0.001	0.001	0.001	0.002	0.001	0.001	-0.261
	P	0.001	0.002	-0.0004	0.0005	0.0003	0.0003	0.0001	0.0002	-0.0001	-0.0002	-0.0002	-0.0002	-0.0002	-0.199
Chlorophyll index	G	0.001	0.0005	-0.001	0.0002	0.0005	0.0002	-0.0003	0.000	0.0002	0.0002	-0.0002	-0.0002	-0.0003	0.255
	P	-0.001	-0.001	0.004	-0.0005	-0.001	-0.001	0.001	0.000	-0.0004	-0.0004	0.0005	0.0004	0.001	0.167
CT @ VS	G	0.005	0.006	-0.002	0.010	0.004	0.001	0.000	-0.0003	-0.0001	0.001	-0.002	-0.001	0.001	-0.210
	P	0.005	0.005	-0.001	0.015	0.003	0.002	0.001	0.0002	-0.0001	0.0005	-0.001	-0.0003	0.001	-0.090
CT @ PFS	G	-0.010	-0.010	0.011	-0.011	-0.026	0.006	0.002	-0.0003	0.005	0.004	0.012	0.009	0.010	-0.612
	P	0.002	0.002	-0.001	0.002	0.009	-0.001	0.0004	0.001	0.001	0.001	-0.002	0.0002	-0.002	-0.149
Plant height	G	-0.009	-0.008	0.004	-0.002	0.006	-0.025	-0.004	0.003	0.001	0.001	-0.005	0.009	-0.007	0.047
	P	0.002	0.001	-0.001	0.001	-0.001	0.006	0.001	-0.0004	0.0001	0.000	0.001	-0.001	0.002	0.089
Pri. branches plant⁻¹	G	0.004	-0.002	-0.009	0.0001	0.002	-0.005	-0.031	-0.014	0.003	0.003	-0.005	-0.001	-0.007	0.176
	P	0.004	-0.002	-0.006	-0.002	-0.002	-0.004	-0.033	-0.014	0.003	0.004	-0.005	-0.001	-0.007	0.149
Sec. branches plant⁻¹	G	0.002	0.012	-0.0005	-0.003	0.001	-0.009	0.033	0.075	0.025	0.015	0.021	0.009	0.023	0.405
	P	0.002	0.007	-0.001	0.001	0.010	-0.004	0.027	0.062	0.022	0.016	0.019	0.012	0.016	0.433
Total no. of Pods plant⁻¹	G	0.0001	0.0001	0.0002	0.000	0.0002	0.000	0.0001	-0.0003	-0.001	-0.001	-0.0003	0.0002	0.0001	0.280
	P	0.001	0.001	0.001	0.0001	-0.001	-0.0002	0.001	-0.004	-0.012	-0.010	-0.005	-0.001	0.001	0.436
Effective pods plant⁻¹	G	0.0003	0.0005	0.0005	-0.0003	0.0005	0.0002	0.0004	-0.001	-0.003	-0.004	-0.0005	0.0004	0.0002	0.117
	P	-0.0005	-0.001	-0.001	0.0003	0.001	0.0001	-0.001	0.003	0.009	0.010	0.002	0.001	-0.001	0.306
Biological yield plant⁻¹	G	-0.227	-0.197	0.176	-0.181	-0.462	0.189	0.155	0.286	0.319	0.139	1.035	-0.479	0.589	0.920
	P	-0.193	-0.158	0.130	-0.097	-0.176	0.180	0.138	0.303	0.373	0.227	0.992	-0.254	0.535	0.895
Harvest Index	G	-0.031	-0.057	0.065	-0.030	-0.121	-0.128	0.012	0.043	-0.065	-0.040	-0.163	0.353	0.065	-0.091
	P	-0.037	-0.052	0.042	-0.009	0.010	-0.089	0.012	0.081	0.042	0.060	-0.111	0.431	0.063	0.188
100-seed weight	G	-0.002	-0.006	0.012	0.008	-0.018	0.015	0.012	0.015	-0.003	-0.003	0.028	0.009	0.050	0.723
	P	-0.0002	-0.0005	0.001	0.0005	-0.001	0.001	0.001	0.001	-0.0003	-0.0003	0.003	0.001	0.005	0.613

G (R SQUARE = 0.9964 RESIDUAL EFFECT = 0.0601) CT@VS= Canopy temperature at vegetative stage
P (R SQUARE = 0.9886 RESIDUAL EFFECT = 0.1070) CT@PFS= Canopy temperature at pod filling stage




Fig. 3: Phenotypic path diagram for grain yield per plant under heat stress condition at Prayagraj during *rabi*, 2017-18.