

**Cypress (*Cupressus macrocarpa*) Leaf Powder
Modulates Metabolism of Maize Weevil
Sitophilus zeamais Motschulsky (Coleoptera:
Curculionidae)**

ABSTRACT

Aims: Maize grains are severely damaged by maize weevil *Sitophilus zeamais*. Exploiting botanicals against synthetic insecticides as a control strategy has gained prominence in recent decades. The aim of the study was to investigate the potential of cypress (*Cupressus macrocarpa*) to exert some control on *S. zeamais*.

Methodology: Maize and *S. zeamais* were collected from Muea market in Buea, Cameroon. The maize was heat sterilized and all debris removed. *Sitophilus zeamais* was reared for three generations before use. Ten *S. zeamais* were stored with 100 g of maize and 0 g, 2 g, 5 g, and 10 g of shade-dried cypress leaf powder was added in three replicates. Data was collected weekly for 5 weeks and analysis of variance was done. The study was conducted in a laboratory of the Teaching and Research farm in the University of Buea, Cameroon, in a randomized complete design.

Results: Mortality of *S. zeamais* was significantly affected ($F = 126.67$, $df = 3, 8$, $p < .001$) by different doses of cypress leaf powder. The mean mortality was 0%, 40.47%, 50.6% and 70.67% for control, 2 g, 5 g and 10 g of cypress leaf powder, respectively. Grain damaged (%) and holed grain were also influenced by cypress leaf powder. The highest grain damaged and holed grain were from the control treatment. A strong positive correlation ($r = 0.814$, $p < .001$) was observed between grain damaged and holed grain while a strong negative correlation was observed between mortality and holed grain ($r = -0.926$, $p < .001$), and grain damaged ($r = -0.913$, $p < .001$).

Conclusion: It is revealed that cypress leaf powder can alter some metabolic activities of *S. zeamais*, thus it can play a role in an integrated pest management program (IPM) of *S. zeamais* in maize storage.

Keywords: Cameroon, holed grain, damaged grain, IPM, metabolic, mortality, *Sitophilus zeamais*.

1. INTRODUCTION

Maize (*Zea mays* L.) is one of the most important staple in the world, particularly in Africa where it is commonly grown by smallholder farmers [1][2]. Like many stored food grains, damage of stored maize grains is a very serious global problem [3]. The situation is worse in developing countries with poor-resource farmers and inadequate storage facilities. Africa is suffering from 20-30 % postharvest losses [4]. with some farmers experiencing 100% postharvest losses in severe infestations. The principal objective of storage is to guarantee food security and that agricultural products are preserved for the realization of price

stabilization, industrial input provision, enhancement of a nations reputation and seeds for planting [5]. These objectives are hardly attained in developing nations due to a plethora of factors amongst which insect pests play a vital role, causing, in some cases, irreparable damage, resulting in major economic losses [6]. Maize grains are vulnerable to many insect pests during storage. Insect pests from the order Coleoptera and Lepidoptera are the most damaging to stored maize grains [7]. [8] and [9] mentioned that lesser grain borer *Rhyzoperthadominica*(Fabricius), weevil complex *Sitophilus* spp. khapra beetle *Trogodermagranarium*Everts, red flour beetle *Triboliumcasteneum*(Herbst), larger grain borer *Prostephanustruncatus* Hornand *Callosobruchusspp.* are very important coleopterous pests. The maize weevil *Sitophiluszeamais*Motchulsky (Coleoptera: Curculionidae) is one of the major primary pests of stored maize in Africa [1][10]. Females lay eggs into the grain which on hatching, the larva feeds towards the inside of the grain until pupa. The adults emerge and eat their way towards the testa causing exit holes resulting into insect damaged grain [11]. *Sitophilus zeamais* infestation results into severe losses due to direct consumption of maize grains, in addition, their activity increases temperature and moisture conditions thus accelerating the growth of mould such as mycotoxins [12], leading to quality deterioration. It is estimated that about 50% maize grain loss is due to *S. zeamias* infestation [13]. Consequently, the control of this pest is imperative. Presently, farmers rely heavily on the use of synthetic insecticides to control *S. zeamias* in Africa. In traditional farm stores in Africa, the use of synthetic insecticides for grain protection has been partially successful [14]. However, questions have been raised about the safety and prolonged use of synthetic pesticides as pests control strategy - synthetic pesticides treadmills have resulted into negative environmental impacts, harm on non-target species, development of resistance in some insect species, increased cost of application, direct hazards from direct exposure and high residues in food and erratic supply of these insecticides in developing countries [10][13][14][15][16]. These problems have rekindled the interest in the re-evaluation of traditional botanical pests control methods which are cheaper, readily available and environmentally friendly [17]. Researches on botanical on *S. zeamais* are many and varied, with emphasis on plant types and plant parts (roots, rhizomes, bulbs, stems, tree bark, leaves, fruits, seeds) and methods of delivery (ash, powder, fumigant, in solvents) (Table 1). In line with previous studies (e.g Table 1), the current study evaluates the mortality potential of cypress leave powder on the maize grain weevil *Sitophilus zeamais*.

Table 1. List of some plants in Africa evaluated for their insecticidal properties on *Sitophilus zeamais*

Scientific name	Common name	Family	Part(s) evaluated	Manner of exposure	Reference
<i>Azadirachta indica</i>	Neem	Meliaceae	Seeds	Powder	[18]
<i>Jatropha curcas</i> L.	Physic nut	Euphorbiaceae	Leave, bark, seeds	Powder	[19]
<i>Citrus sinensis</i> L.	Orange	Rutaceae	Peels	Powder	[19]
<i>Capsicum frutescens</i> L.	Chilli	Solanaceae	Fruits	Powder	[20][21]
<i>Allium cepa</i>	Onion	Liliaceae	Bulb	Powder	[20]
<i>Piper guineense</i>	Black pepper	Piperaceae	Seeds	Powder	[13][20][21]
<i>Allium sativum</i> L.	Garlic	Liliaceae	Bulb	Powder	[22]
<i>Nicotianatabacum</i> L.	Tobacco	Solanaceae	Leaves	Powder	[22]
<i>Ageratum conyzoides</i>	Billygoatwee	Asteraceae	Leaves	Solvents	[23]

	d				
<i>Lantana camara</i>	Lantana	Verbenaceae	Leaves	Solvents	[23]
<i>Chromolaenaodorata</i>	Siam weed	Asteraceae	Leaves	Solvents	[23]
<i>Curcuma longa</i> L.	Turmeric	Zingiberaceae	Rhizome	Solvents	[24]
<i>Gnetum africanum</i>	Eru	Gnetaceae	Leaves	Solvents	[24]
<i>Ocimum basilicum</i>	Basil	Lamiaceae	Leaves	Powder	[25]
<i>Ricnodendron heudelotii</i>	Njangsa	Euphorbiaceae	Leaves, bark, roots	Powder	[26]

2. MATERIAL AND METHODS

2.1. Local and experimental design

This study was conducted in one of the laboratory of the Teaching and Research Farm, House of the Faculty of Agriculture and Veterinary Medicine of the University of Buea. It is located at the foot of mount Cameroon in Fako Division, South West Region of Cameroon, at latitude/longitude 4° 9' 33" N/9° 14' 12" E. It was conducted at an average temperature of 27°C and 65% relative humidity. The experiment was laid out in a complete randomized design (CRD) with 4 treatments and 3 replicates. The study was conducted in 2014.

2.2. Maize and insect collection

1.5kg of dry maize was bought from traders in the Muea market located in Buea municipality, south west region, Cameroon. Cracked, broken grains and other debris were sorted out from the others. The maize grains were heat sterilized at 40-50 °C for 20 mins to kill any residual eggs. Adults *Sitophilus zeamais* were obtained from the Muea market and reared under ambient condition of 12:12 photoperiods, temperature of 27°C and humidity of 65%. The insects were stored in sterilized (as above) maize grains for 70 days during which time three generations must have passed.

2.3. Plant powder preparations

Fresh plant material (leaves) of *Cupressus macrocarpa* Wilma (Pinales: Cupressaceae) were collected from Bamenda, north west region of Cameroon (between latitudes 4°50' to 5°20'N and longitudes 10°35' to 11°59'E of the Green Wish Meridian). The leaves were shade-dried in one of the rooms (25 °C, 60-65% RH) of the teaching and research farm house of the faculty of agriculture and veterinary medicine of the University of Buea for 2 weeks to avoid volatilization of the active ingredients. On drying, the leaves were ground into powder by a Victoria grinding mill and parked into a black polythene bags for future use.

2.4. Experimental setup, Data Collection and data analyses

2.4.1. Mortality test

Four different doses of cypress leaf powder (0g, 2g, 5g, and 10g) were measured in the laboratory using an analytical balance and introduced into different containers holding 100 g of sterilized maize grains. 10 adult weevils were introduced in the grains and mortality was checked weekly for 3 weeks. The insects were considered dead on failure to respond to three probing using a blunt dissecting probe [15]. At the end of the third week, a cumulative mean mortality was calculated using equation 1.

$$\text{Cumulative mean mortality (\%)} = \frac{\sum_{i=1}^3 x_i}{n} \times 100 \dots\dots\dots \text{eq. 1.}$$

Where n is the number of maize weevil per trial (10) and x_i is the mean weekly mortality of maize weevil per treatment in the ith week.

2.4.2. Percent Holed and Percent Damaged grain

Ten unsexed adult weevils were stored in 100 g of maize with the four different doses. Two weeks after treatment (DAT) the adult live insects were removed from treated and control containers. The number of grains adult exit holes and damaged grain was counted after 4 weeks after infestation for the calculations of percentages of grain damaged and holed grains. Percentages were calculated using equation 2 (percent damaged grain) and equation 3 (percent holed grain).

$$\text{Percent damaged grain} = \frac{\text{Number of damaged grains} \times 100}{\text{Total number of grains}} \dots \text{eq. 2.}$$

$$\text{Percent Holed grain} = \frac{\text{Number of Holed grains}}{\text{Total number of grains}} \times 100 \dots \text{eq. 3.}$$

2.5. Data analysis

Normality and homogeneity of variance tests were conducted using Kolmogorov-Smirnov test in SPSS (ver 23) and Glantz' modification of Levene's test on Microsoft Excel add-in Daniel's XL Toolbox version 7.3.4 (<http://www.xltoolbox.net>), respectively, prior to one-way Analysis of Variance (ANOVA) test for statistical significance. The Bonferroni-Holm post hoc test on the MS Excel add-in was used separate the significantly different means at an alpha (α) level of 0.05. A correlation analysis was done to ascertain how the parameters vary concurrently. A hierarchical cluster analysis was used to understand to evaluate the similarity of the treatments considering all parameters simultaneously. The Paired group (UPGMA) algorithm and the Euclidean similarity index were used. The correlation and cluster analyses were conducted using Paleontological Statistics (PAST ver. 3.25).

3. RESULTS AND DISCUSSION

3.1. Mortality test

The weekly mortality of *S. zeamais* as a result of exposure to different doses of cypress leaf powder is shown in table 2. The mortality was significantly different ($F = 14.55$, $df = 3, 8$, $p = .0013$) after week 1. About 3, 2, and 1 weevil died in 10 g, 5 g and 2 g of cypress leaf powder, respectively in the first week (Table 2). The number of weevils that died in the second week from cypress leaf powder treatments were not significantly different although different from the control treatment ($F = 13.83$, $df = 3, 8$, $p = .0016$). After week 3, the mean mortality of *S. zeamais* was significantly highest from 10 g of cypress leaf powder compared to 2 g of cypress leaf powder and control treatments ($F = 15.0$, $df = 3, 8$, $p = .0012$). It is clear from the findings in table 2 that cypress leaf powder exhibits some significant mortality to *S. zeamais*. No *S. zeamais* mortality was recorded from control treatment. [17]reported that cypress leaf powder showed some repellency to maize weevil *S. zeamais*. The current study suggests that, in addition to repellency effect, cypress leaf powder has some significant mortality potential on *S. zeamais*. Many botanicals are known to exhibit some mortality and repellency to *S. zeamais*. For instanc, [25]reported that basil (*Ocimumbasilicum*) leaf, bark and an admix of leaf and bark powder killed and repelled *S. zeamais*.

The cumulative mean percentage mortality of *S. zeamais* after three weeks of exposure to different doses of cypress leaf powder is shown in table 2. There was a significant difference in the cumulative mean mortality (%) of *S. zeamais* ($F = 126.67$, $df = 3, 8$, $p < .001$). The cumulative mean mortality (%) was 0%, 40.47%, 50.70% and 70.67% for control, 2 g, 5 g and 10 g of cypress leaf powder, respectively (Table 2).The cumulative mean mortality revealed that *S. zeamais* mortality was dose dependent. The higher the dose of the cypress leaf powder, the higher the cumulative mean mortality of *S. zeamais*. More *S. zeamais* died in 10 g of cypress leaf powder treatment than from the other treatments. In fact, the cumulative mean mortality from 10 g of cypress leaf powder treatment was almost twice that from 2 g of cypress leaf powder treatment. It seems that *S. zeamais* mortality is plant-type, plant-part and dose dependent. [27]reported a progressive increase in *S. zeamais* mortality with increasing dose (0.1, 0.2.0.3 and 0.4 g/50g of maize grain) of powdered seeds of black

pepper *Piper guineense* after three weeks of exposure. The current study is on par with those of [23][25].

Table 2. Mean mortality (\pm sd) of maize weevil weeks after treatment with cypress leaf powder

Dose (g)	Week 1	Week 2	Week 3	Cumulative percentage (%)
0	0 \pm 0.0 a	0.0 \pm 0.0 a	0.0 \pm 0.0 a	0.0 \pm 0 a
2	1.3 \pm 0.57 b	1.7 \pm 0.57 b	2.3 \pm 0.57 b	40.47 \pm 5.7 b
5	1.7 \pm 0.57 bc	1.7 \pm 0.57 b	2.7 \pm 0.57 bc	50.70 \pm 4.7 b
10	2.7 \pm 0.58 c	2.0 \pm 0.00 b	3.0 \pm 0.0 c	70.67 \pm 5.8 c

Mean mortality of 10 maize weevil treated with different doses (g) of cypress leaf powder. Means in the same column with the same letter(s) are not significantly different by Bonferroni-Holm post hoc test at $p < 0.05$. sd – standard deviation

3.2. Grain damaged (%)

The percentage maize grain damaged by *S. zeamais* exposed to different doses of cypress leaf powder is shown in figure 2. A one-way ANOVA revealed that there was a statistical significant difference ($F = 32.72$, $df = 3, 8$, $p < .001$) in maize damaged (%) by *S. zeamais*. The highest grain damaged was 71.0% from control treatment, followed by 60.66%, 52.33% and 43.66% from 2 g, 5 g and 10 g of cypress leaf powder, respectively (Figure 1). Some significant amount of grain damaged was observed from all treatments, although that of 10 g of cypress leaf powder was below 50%. This shows that cypress leaf powder prevented quality degradation of maize grains. The current study is in accordance with that of [9] who reported decreasing maize grain damage by *S. zeamais* with increasing dose of powders of *Lippia javanica* dried leaves and *Spirostachys Africana* wood. According to [20] quality deterioration of maize grains has severe economic implications especially for poor-resourced farmers in the developing countries.




Figure 1. Mean maize grain damaged (%) by *Sitophilus zeamais* after five weeks of exposure to different doses of cypress (*Cupressus macrocarpa*) leaf powder. Means with the same letter(s) are not significantly different according Bonferroni-Holm post hoc test ($\alpha = .05$).

3.3. Percent holed grain (%)

The percentage of holed maize grains due to *S. zeamais* activity after exposure to different doses of cypress leaf powder is shown in in [figure 2](#). There was a significant difference ($F = 17.26$, $df = 3, 8$, $p = .0007$) in holed grains (%). The least holed grain was 49.67% from 10 g of cypress leaf powder treatment. It was followed by 54.67%, 67.0% and 86.0% from 5 g and 2 g of cypress leaf powder, and control treatment, respectively. A similar pattern to grain damage (%) was observed for holed grain (%). A significant percentage of maize grain had holes, although that for 10 g of cypress leaf powder treatment was below 50%; all others were above 50%. Indeed, the percentage holed grains from control treatment was twice that from 10 g of cypress leaf powder treatment. Holed grain (%) showed an inverse relationship to dose of cypress leaf powder. Holed grains is an indication that the grain content has been consumed by *S. zeamais*. Such holed grains do not eventually germinate. [27]reported that germination of stored maize grains increased with increased doses of *Piper guineense*, due to reduced *S. zeamais* activity. Inability of stored maize grains to germinate due to holed grains falls short of one of the fundamental principles of storage [5]




Figure 2. Mean holed grain (%) of maize by *Sitophilus zeamais* after five weeks of exposure to different doses of cypress (*Cupressus macrocarpa*) leaf powder. Means with the same letter(s) are not significantly different according Bonferroni-Holm post hoc test ($\alpha = .05$).

3.4. Correlation matrix for three variables measured

A correlation analysis was conducted to ascertain how the measured parameters vary together. The correlation matrix is show in [figure 3](#). A strong positive correlation [$r(18) = 0.814$, $p < .001$] between maize grain damaged and holed grain was observed. On the other hand, there was a strong negative correlation between cumulative mortality and grain damaged [$r(18) = -0.913$, $p < .001$]. In addition, a strong negative correlation was observed between cumulative mortality and holed grain [$r(18) = -0.926$, $p < .001$] ([Figure 3](#)). The correlation matrix reveals two key points; (i) increased mortality of *S. zeamais* results into less and less maize grain damage (%) and holed grains (%), in other words, there is an indirect proportional relationship between *S. zeamais* mortality and maize grains (%), and (ii) there is a direct proportional relationship between grain damage (%) and holed grain (%), in other words, any attempt to minimize grain damaged can equally minimize holed grain (%).

The finding of the current study is in accordance with that of [27] who reported increased mortality of *S. zeamais* due to its exposure to powder of *Piper guineense* resulted into decreased maize grain damaged and eventually higher germination rates. Similar findings have been reported by [28] even though inert materials were used.




Figure 3. Correlation matrix of some measured parameters of *Sitophilus zeamais* activity after exposure to different doses of cypress (*Cupressus macrocarpa*) leaf powder in stored maize

3.5. Similarity

The result of the hierarchical cluster analysis is shown in the dendrogram in **figure 4**. Two distinct clades were observed; one containing cypress leaf powder treatments and the other containing control treatment. The observations of 10 g of cypress leaf powder were most conserved. The dendrogram clearly shows that results from 2 g and 5 g were more similar than 2 g and 10 g of cypress leaf powder. The findings from the similarity study (cluster analysis) further support the idea that an active component or components in the cypress leaf powder actually influenced the metabolism of *S. zeamais* and is dose-dependent with 10 g of cypress powder being the best dose resulting into higher mortality and a reduced maize grain damaged (%). The results are consistent with those of other studies [10][13][14][17]. and purports that cypress leaf powder can play a phenomenal role in the management of *S. zeamais*.




Figure 4. Similarity between different doses of cypress (*Cupressus macrocarpa*) leaf powder treatments on the activities of maize grain weevil *Sitophilus zeamais*.

4. CONCLUSION

The main goal of this study was to evaluate the potential of cypress leaf powder to alter some metabolic processes of *S. zeamais*. The study has shown that mortality of *S. zeamais* due to cypress leaf powder exposure is dose-dependent with 10 g of cypress leaf powder being the preferred dose. In addition, it also demonstrated that increasing mortality of *S. zeamais* will eventually reduce maize grain damage. The evidence from this study suggest that cypress leaves adds to the growing list of botanicals that can play a tremendous role in an integrated pest management program for *S. zeamais*.

REFERENCES

- [1] Nukenine EN, Monglo B, Awasom I, Ngamo LST, Thuenguem FFN, Ngassoum MB. Farmer's perception on some aspects of maize production and infestation levels of stored maize by *Sitophilus zeamaiz* in the Ngaoundere region of Cameroon. *Cameroon Journal of Biology, and Biochemical Sciences*. 2002;12(1):18-30.
- [2] Ntonifor NN, Oben EO, Konje CB. Use of selected plant-derived powders and their combinations to protect stored cowpea grains damage by *Sitophilus zeamais*. *ARP NJ. Agricultural Journal of Biological Science*. 2010;5:13–21.
- [3] Upadhyay RK and Ahmad S. Management strategies for control of stored grain insect pests in farmer stores and public ware house. *World Journal of Agricultural Science*. 2011;7(5):527–549.
- [4] Owusu EO, Osafo WK, Nutsukpui ER. Bioactivity of candlewood, *Zanthoxylum xanthoxyloides* (LAM.) solvent extracts against two stored-product insect pests. *African Journal of Science and Technology*. 2007;8:17–21.
- [5] Lale NES. *Stored product entomology and acarology in Tropical Africa*. Mole Publications, Nigeria Ltd, Maiduguri, Nigeria. 2002;4-5.

- [6] Denloye AAB and Makanjuola WA. Insecticidal promises of plant terpenoids for the control of insect pests of stored grains in the 21st Century. *Journal of Research Revelations in Science*. 2001;2:271–288.
- [7] Firdissa E and Abraham T. Effect of some botanical and other materials against the maize weevil, *Sitophilus zeamais*Motschulsky on stored maize, *In: Maize Production Technology for the Future: Challenge and Opportunities: Proceedings of the Sixth Eastern and Southern Africa Regional Maize Conference, 21 – 25 September 1998, Addis Ababa, Ethiopia*. 1999;101–109.
- [8] Shafique M and Chaudry MA. Susceptibility of maize grains to storage pest. *Pakistan Journal of Zoology*. 2007;39(2):77–81.
- [9] Chikukura L, Mvumi BM, Chikonzo R. Chenzara C. Evaluation of selected indigenous pesticidal plant powders against stored maize and cowpeas insect pest. *African Crop Science Proceedings*. 2011;10:189–192.
- [10] Zhou HN, Zhao NN, Shu Shan D, Yang K, Cheng FW, Zhi LL, Yan JQ. Insecticidal activity of the essential of *Lonicera japonica* flower buds and its main constituent compounds against two grain storage insects. *Journal of Medicinal Plant Research*. 2012;6(5):912–917.
- [11] Tadesse A and Basedow T. Laboratory and field studies on the effect of natural control measures against insect pests in stored maize in Ethiopia. *Journal of Plant Disease Protection*. 2005;112(2):156–172.
- [12] Tripathi AK, Prajapati V, Verma N, Bahl JR, Bansal RP, Khanuja SPS. Bioactivity of the leaf essential oil of *Curcuma longa* (Var. Ch-66) on three species of stored product beetles (Coleoptera). *Journal of Economic Entomology*. 2002;95(1):183–189.
- [13] Udo IO, Ekanem MS, Inyang EU. Laboratory evaluation of west African black pepper (*Piper guineense*) seed powder against maize weevil (*Sitophilus zeamais* Mots.). *Munis Entomology and Zoology*. 2011;6(2):1003–1007.
- [14] Ogendo JO, Deng AL, Belmain SR, Walker DJ, Musandu AAO. Effect of insecticidal plant materials, *Lantana camara* L. and *Tephrosiavogelii* Hook, on the quality parameters of stored maize grains. *Journal of Food Technology in Africa*. 2004;9:29–36.
- [15] Oben-Ofori D, Reichmuth CH, Bekele J, Hassanali A. Biological activity of 1,8 Cineole, a major component of essential oil of *Ocimumkenyense* (Ayobangira) against stored product beetles. *Journal of Applied Entomology*. 1997;121:237–243.
- [16] Achiri DT, Akotsen-Mensah C, Jallow M, Gbewonyo WSK, Afreh-Nuamah K. High levels of some insecticides on cabbage samples (*Brassica oleracea* L. varCapitata) from rural and urban farmers in Ghana. *J. Environmental Sci. Toxicology and Food Technology*. 2016;10(4):97-103.
- [17] Achiri DT. and Njweng MA. Bioactivity of cypress leaf powder (*Cupressusmacrocarpa*) on cowpea weevil (*Callosobruchusmaculatus*Fabr. Coleoptera: Bruchidae) and maize weevil (*Sitophilus zeamais*Motschulsky, Coleoptera: Curculionidae) in stored maize grains in Cameroon. *Inter. J. Interdisciplinary and Multidisciplinary Studies*. 2015;2(4):1–10.
- [18] Parugrug AM and Roxas C. Insecticidal action of five plant against maize weevil, *Sitophilus zeamais*Motsch. (Coleoptera: Curculionidae). *King Mongkut's Institute of Technology Ladkrabang Science and Technology Journal*. 2008;8(10):24–38.
- [19] Suleiman M and Yusuf MA. The potential of some plant powders as biopesticides against *Sitophilus zeamais* (Motsch.) (Coleoptera: Curculionidae) and *Callosobruchusmaculatus* (F.) (Coleoptera: Bruchidae) on stored grains: a review. *Bayero Journal of Pure and Applied Science*. 2011;4(2):204–207.
- [20] Adedire CO. and Ajayi TS. Assessment of the insecticidal properties of some plant extracts as grain protectants against the maize weevil, *Sitophilus zeamais*Motschulsky. *Nigerian Journal of Entomology*. 1996;13:93–101.
- [21] Asawalam EF, Emosairue SO, Ekeleme F, Wokocho RC. Insecticidal effects of powdered parts of eight Nigerian plant species against maize weevil, *Sitophilus zeamais*Motschulsky (Coleoptera: Curculionidae). *Electronic Journal of Environmental, Agricultural and Food Chemistry*. 2007;6(11):2526–2533.

- [22] Danjumma BJ, Majeed Q, Manga SB, Yahaya A, Dike MC, Bamaiyi L. Effect of some plant powders in the control of *Sitophilus zeamais* Motsch (Coleoptera: Curculionidae) infestation on maize grains. American-Eurasian J. Sci. Res. 2009;4(4):313–316.
- [23] Bouda H, Tapondjou LA, Fontem DA, Gumedzoe MYD. Effect of essential oils from leaves of *Ageratum conyzoides*, *Lantana camara* and *Chromolaena odorata* on mortality of *Sitophilus zeamais* (Coleoptera, Curculionidae). Journal of Stored Products Research. 2001;37:103–109.
- [24] Asawalam EF and Chukwuekezie AL. Control of maize weevil, *Sitophilus zeamais* (Motschulsky) using extracts of *Gnetum africanum* (Afang) leaves and *Curcuma longa* L. (Turmeric) rhizomes. Inter J. Agricultural Sci. 2012;2(9):263-265.
- [25] Mwangangi BM and Mutisya DL. Performance of Basil powder as insecticide against maize weevil, *Sitophilus zeamais* (Coleoptera: Curculionidae). Discourse Journal of Agriculture and Food Science. 2013;1(11):196–201.
- [26] Epidi TT, Udo IO, Osakwe JA. Susceptibility of *Sitophilus zeamais* Mots. And *Callosobruchus maculatus* F. to plant parts of *Ricinodendron heudeloti*. J. Plant Pro. Res. 2009;49(4):411–415.
- [27] Asawalam EF. and Emosairue SO. Comparative efficacy of *Piper guineense* (Schum and Thonn) and pirimiphos methyl on *Sitophilus zeamais* (Motsch.). Tropical and Subtropical Agroecosystems. 2006;6:143–148.
- [28] Gemu M, Getu E, Tadess T, Yosuf A. Management of *Sitophilus zeamais* Motschulsky (Coleoptera: Curculionidae) and *Sitotrogacerealella* (Olivier) (Lepidoptera: Gelechiidae) using locally available inert materials in Southern Ethiopia. Discourse Journal of Agriculture and Food Science. 2013;1(6):111-117.