

**Pedogeomorphological categorization of selected soils in Mbogo - Komtonga irrigation scheme, Mvomero District, Morogoro Region, Tanzania**

**Abstract**

Pedogeomorphological categorization of selected soil profiles developed on alluvial deposits in Mbogo - Komtonga traditional irrigation scheme, Mvomero District in Morogoro Region, was carried out during February 2017. Using standard grids, pedogeomorphic approach and standard manuals, detailed soil survey was conducted which enabled delineation of soil mapping units from which the representative profiles were identified, described and sampled. Eighteen samples were collected at a depth of 0 – 30 cm and from each horizon of the selected soil Master pits and analyzed for physico – chemical characterization. Based on FAO soil survey system of classification, the representative profiles were classified as Eutric Fluvisols and/or Eutric Cambisols. The pedon was deep to moderately deep, well to moderately well drained, with brownish black clay top soils or dull yellowish brown soil colors with sub soils stratified with fS, C, CL and SCL. Top soil pH was strongly to medium acid to medium or slightly acid sub soils. OC showed no decline in soil quality. N was very low to low, P and K levels were medium to low or very low; CEC was high to very high in all the profiles.  $Ca^{2+}$  and  $Mg^{2+}$  in the top soils were high to very high and very low or low to medium in the sub soils.  $Na^+$  was rated as low to medium in the top soils of all profiles and low to very low in the sub soils. Base saturation was > 50 % and was rated as high. Topsoil Bd and total porosity were ideal to medium. AWC was medium and water storage capacity (AWSC) was good and sufficient for paddy production and other upland crops. These results suggest that where the soil parameters were low to very low as for N, P and SOM should be included in the overall soil fertility management program. Soil reaction may be regulated during irrigation development by provision of sufficient drainage, discharge and flood control structures and minimum application of lime if required.

**Key words:** Irrigation, alluvial plain, geomorphology, master pits, classification, mapping units

**Abbreviations:** CEC = Cation exchange capacity; AWC = Available water capacity, AWSC = Available water storage capacity, SOM = Soil organic matter, OC = Organic carbon, Bd = Bulk density, N = Nitrogen, P = phosphorus, K = potassium, Ca = calcium, Mg = magnesium, fS = fine sand, C = clay, CL = clay loam, SCL = sand clay loam

## 1 INTRODUCTION

Agriculture sector in Tanzania is by far the largest sector of the economy. It accounts for 24.7 per cent of the GDP, 20 per cent of traditional export earnings, 95 per cent of food requirement, employs 75 per cent of the population and food contributes about 55.9 per cent of the inflation basket [1]. Comparatively, the financial sector is only roughly 10 per cent of the size of the agriculture sector [2]. Although there are specific and main traditional export crops in Tanzania such as cashewnuts (*Anacardium occidentale*), maize (*Zea mays* L) and rice (*Oryza sativa* L) have recently assumed the role of both food and cash crops also exported during the years of surplus. Maize (*Zea mays* L.) is an important staple food for the majority of Tanzanians [3] and about 80 per cent of it is produced by small - scale farmers grown on over 4.9 million ha [4; 5]. Between 65 and 80 per cent of all maize is consumed within the producing households and only 20 to 35 per cent enters commercial channels. It has been identified as a key crop to enhance food production, income, poverty alleviation and food security [6]. Maize provides 60 per cent of dietary calories, over 30 per cent house income and more than 50 per cent of utilizable protein [7; 8]. Estimates suggest that there might be 150 million Tanzanians by 2050, and so, the National demand for maize will have to grow in the future to meet demand of the growing population in response to growth of national Gross Domestic Product (GDP) at nearly 7 per cent per annum. Some studies have reported that Food security must account for opportunities to increase production against projected changes in demand associated with population growth and changing diets, need to reduce the environmental footprint of agriculture, and limited availability of land suitable for crop production [9; 10; 11].

Rice (*Oryza sativa* L) is the second most important crop after maize (*Zea mays* L). Tanzania is the second largest producer of rice in Southern Africa after Madagascar, with production level of 818,000 tons [12] or 2.2 million tons currently. The cultivated area is 681,000 ha and this represents 18 % of Tanzania's cultivated land. About 71 % of the rice grown in Tanzania is produced under rain fed conditions, where irrigated land presents 29 % of the total land with most of it in small scale traditional irrigations with the average yield of 1 - 1.5 t ha<sup>-1</sup> [13]. About half of the production is concentrated in Morogoro, Shinyanga, and Mwanza regions and virtually, 99 % of rice is grown by smallholders in Tanzania, although some of them are part of large - scale rice irrigation schemes that were formerly state - managed farms [14]. Despite the importance of maize and rice, its production is challenged with amongst others to low investment, low soil fertility, and unsustainable agricultural practices leading to land depletion.

Land depletion is caused by inappropriate land use and soil management practices, including poor cropping and farming systems, shortening and or elimination of the fallow period, insufficient and inadequate use of farm manures and fertilizers, nutrients mining and soil erosion [15]. Soil deprivation and underutilization of appropriate mineral elements in crop production portends food security in Tanzania, Mvomero District inclusive [16]. The main reasons of land deprivation include depletion of plants mineral elements, deletion of whole crop residues, use of low levels of mineral elements during crop production and inadequate soil conservation practices [17] and longer cultivation [18]. These factors has been the main reason for low soil fertility with resultant impact on crop production and productivity including grain quality, cost of production and the increased risk of soil erosion. Maintaining long - term soil fertility through conventional agriculture has certain limitations [19]. For example, studies on a continental soil nutrient balance in 38 sub - Saharan countries involving 35 crops [20] has reported that soil nutrient balances were negative for N, P, K mineral elements with mean annual losses of 22 kg N, 2.5 kg P and 15 kg K ha<sup>-1</sup>. This indicates that improving the production and productivity of agriculture in Mbogo Komtonga for example, is

80 greatly dependent on efficient utilization and management of soils [21]. Different soil types exhibit varying  
81 characteristics due to differences in micro - morphological, morphological, physical, chemical and  
82 mineralogical properties [22]. Variations in soil forming factors and processes operating on different parent  
83 materials, under different climatic, topographic, and biological conditions over varying periods would cause  
84 these variations [23].

85  
86 Soil categorization and classification therefore helps to generate required information for land use planning  
87 and soil management purposes. Soil surveys are important for soil characterization and classification  
88 purposes and aids in the creation of data bases on soil morphology, physical and chemical properties [23;  
89 24; 25]. This information is important for determining agricultural potential, limitations and possible  
90 management options for the soils in a particular area thereby helping in selection of the best agricultural  
91 enterprises suitable for that area [26; 27]. Irrigation projects can be planned and developed based on  
92 information obtained from soil characterization and classification. Area specific soil fertility management  
93 strategies, aimed at increasing crop production, can be developed for a particular area using soil survey  
94 data instead of using general fertilizer recommendations. Information on soil characterization and  
95 classification can be utilized widely by land use planners, agriculture researchers, extension staff,  
96 development agents and farmers in order to sustainably increase agriculture production.

97  
98 A detailed study of the soil characteristics and classification will provide baseline information on the  
99 physical, chemical and mineralogical properties of the soil for crop production, land use planning and  
100 management. Despite the fact that Mbogo - Komtonga irrigation scheme in Mvomero District is an intensive  
101 producer of rice and maize there is no soil pedogeomorphological characterization and classification that  
102 have been done on the soils of the area. Soil pedogeomorphological characterization and classification of  
103 the Mbogo - Komtonga irrigation scheme are very important in providing the needed basic information on  
104 soils of the area. Thus, this study aims to characterize the soils of the area based on their  
105 pedogeomorphological characteristics, physico - chemical properties and their classification according to  
106 the FAO – Unesco Soil Map of the World system of classification [28]. The results emanating from the  
107 study will provide information on the soil fertility trends and will serve to guide activities related to the  
108 management of the existing land resources for sustainable agricultural production in Mvomero District  
109 Therefore, the objective of this study was to characterize the soils under maize production in Mbogo -  
110 Komtonga, Mvomero District, Morogoro Region and to recommend management practices required for  
111 sustainable crop production.


## 112 113 **2 MATERIALS AND METHODS**

### 114 115 **2.1 Description of the study area**

116  
117 The study was carried out in Mbogo Komtonga traditional irrigation scheme located in Mvomero District,  
118 Morogoro Region, Tanzania. It is bordered by Kichangani village to the North, Nguu Mountains in the West,  
119 Diwale/Mbulumi River in Kisala village to the East, and Kigugu village to the South. Administratively, the  
120 project area is located at Mbogo - Komtonga village, Sungaji ward, Turiani division, Mvomero District,  
121 Morogoro Region (Fig 1). Agricultural practices in Mbogo - Komtonga irrigation scheme are both traditional  
122 irrigation and rainfed. The main crop grown in this area is rice (*Oryza sativa* L) mainly as food and cash  
123 crop. According to the interviewed farmers, hand hoe is the overall dominant tool for land preparation.  
124 Failure of crops in these areas is due to prolonged flood during rainy season, nutrient leaching and

125 inadequate irrigation water in the dry season, suggesting irrigation development. Generally, when irrigation  
 126 water is needed it is not sufficiently available and when sufficiently available, there is no drainage. For rice  
 127 cultivation, early planting starts in December and harvested in May while late planting starts in January and  
 128 harvested in June. After the main rainy season, most farmers use residual moisture to grow maize, cassava  
 129 and horticultural crops. The current average production of paddy ranges between 2.5 – 3.0 t ha<sup>-1</sup>. Among  
 130 other factors, low crop yield is attributed to low or no use of agricultural inputs, lack and or poor irrigation  
 131 infrastructure and lack of drainage during the rainy season. Rainfall in the study areas is bi-modal with 46.2  
 132 % of the total rains falling between March through May and about 44.5 % light rains falling between  
 133 November through February. The total average annual rainfall is about 970 mm. Temperature, RH (%),  
 134 potential evaporation and other climate variables representative of the study areas are presented in Table 1  
 135 and Fig 2. The mean temperature varies from 21.8°C in July to 27.0°C in February. All pedons have an  
 136 isohyperthermic soil temperature regime (STR) and udic soil moisture regime. The monthly average relative  
 137 humidity (RH) varies from 58.8 (i.e. October) to 77.9 % (i.e. April). The potential evaporation is about 1,799  
 138 mm per annum and varies widely throughout the year from 93.5 to 206.9 mm per month in June and  
 139 December respectively.

140



141

142 **Fig 1: Location of the study area**

143

144

**Table 1: Climatic data representative for Mvumi and Mbogo - Komtonga Irrigation schemes**


| Description                    | J     | F     | M     | A     | M    | J    | J     | A     | S     | O     | N     | D     |
|--------------------------------|-------|-------|-------|-------|------|------|-------|-------|-------|-------|-------|-------|
| Rainfall (mm m <sup>-1</sup> ) | 119.4 | 107.7 | 163.6 | 204.4 | 80.3 | 16.5 | 9.3   | 14    | 11.8  | 38.6  | 81.3  | 132.7 |
| T mean max (°C)                | 32.2  | 32.5  | 31.9  | 29.9  | 28.8 | 28.1 | 27.7  | 28.5  | 30.2  | 31.6  | 32.2  | 32.3  |
| T mean min (°C)                | 21.6  | 21.5  | 21.3  | 20.8  | 19.1 | 16.4 | 15.8  | 16.4  | 17.5  | 18.8  | 20.3  | 21.5  |
| T mean (°C)                    | 26.9  | 27    | 26.6  | 25.3  | 24   | 22.2 | 21.8  | 22.4  | 23.9  | 25.2  | 26.3  | 26.9  |
| Evap. (Pan) mm m <sup>-1</sup> | 192   | 177.1 | 160.5 | 110.3 | 96.3 | 93.5 | 105.4 | 117.5 | 161.2 | 186.1 | 191.7 | 206.9 |
| 0.5ETo (mm m <sup>-1</sup> )   | 96    | 88.6  | 80.3  | 55.1  | 48.2 | 46.8 | 52.7  | 58.8  | 80.6  | 93    | 95.9  | 103.4 |
| RH mean (%)                    | 65.4  | 65.2  | 69.8  | 77.9  | 75.7 | 70.5 | 68.5  | 65.8  | 60.3  | 58.8  | 60.3  | 63.2  |
| SH (hrs.)                      | 7.9   | 7.7   | 6.8   | 5.8   | 5.9  | 6.5  | 6.3   | 6.6   | 7.5   | 8.1   | 8.2   | 7.9   |
| WS (km day <sup>-1</sup> )     | 252   | 232.9 | 172.7 | 89    | 85.3 | 99.4 | 120.6 | 150.3 | 185.5 | 187.4 | 238   | 261.6 |

145

**Source:** Mtibwa, Ilonga, Dakawa, Dakawa Rice farm and Morogoro Meteorological weather stations. Total annual rainfall ≈ 970 mm, Total annual Evaporation (Pan) ≈ 1,799 mm

146

147



148

149 **Fig 2:** Water balance and the determination of the growing period for Mbogo – Komtonga irrigation  
 150 schemes.

151

152 **2.2 Field methods**

153

154 Soil survey was planned by means of gridlines. A GPS device was used to carry out boundary survey, set  
 155 out grid lines, prepare or reconfirm a base map, as well as recording the coordinates and elevations of all  
 156 field observations. During the fieldwork, two kinds of observations were made. These were auguring and  
 157 profile (master) pits observations. Auger observation was taken as an identification of the taxonomic unit for  
 158 which a particular pedon belong. The standard depth was taken as 120 cm but extended further to 150 cm  
 159 whenever necessary and possible. A total of one hundred and ten (110) augers were observed at a depth  
 160 of 120 cm. Core samples were also sampled from 0 – 50 cm and 50 – 100 cm soil depth of each profile.  
 161 Fourteen (14) core samples were then sampled from 7 profiles. The second observation was full pit or full  
 162 profile. This was done after the establishment of the important soil sets. A total of three (3) profiles and one  
 163 (1) minipit were opened and described respectively. These representative profiles were MB - Pa1, MB -  
 164 Pa2 and MB - Pa3. The approximate volume of a full master pit was 150 cm x 150 cm x 120/150 cm. These  
 165 observations were concisely described according to the **FAO guidelines (1977)** for soil description and were  
 166 carefully entered abreviatively on a pre - prepared data form. Soil classification was done by using the FAO  
 167 – UNESCO soil map of the world (1988). Overall, 50 disturbed soil samples were collected for physical -  
 168 chemical characterization. Of the total disturbed samples, 41 were from master pits and 9 were collected as  
 169 composite soil samples from a soil depth of 0 – 30 cm. Geomorphologically, the proposed irrigation scheme  
 170 fall into one landscape, that is, the Plain or Floodplain [29]. It is essentially a flat area with moderate to  
 171 imperfectly drainage condition. Most of clearly drainable sections of rivers flow from NW to SE direction and  
 172 a few flows southerly. The geology of the area can generally be described as having alluvium deposits  
 173 probably originating from the high plateau. The high plateau is covered by red brown and in places, light

174 grey earth particularly on the flat ridges. Deep weathering of the gneisses which may have originated  
175 during the Neogene's period is a common feature on the plateau. The lowlands have a thick cover of black  
176 cotton soil (mbugas) which in places are replaced by light coloured sandy soils which is partly alluvial. Light  
177 grey clays are common in the marshy areas and are probably found in several layers within the soil profile  
178 observed in the lowlands suitable for agricultural purposes. Table 2 presents the salient features of the  
179 study sites.

UNDER PEER REVIEW

180  
181  
182

**Table 2:** Salient features of the study area

| Soil Profile | MU       | Village          | Scheme     | District | Coordinates      |                | Alt. (m) | Parent Material | Landform           | Land use  | Soil classification           | SMR   | STR             |
|--------------|----------|------------------|------------|----------|------------------|----------------|----------|-----------------|--------------------|---|-------------------------------|-------|-----------------|
| MB - SP1     | MB - Pa1 | Mbogo - Komtonga | Irrigation | Mvomero  | North<br>9316954 | East<br>344246 | 364      | Alluvial        | flat to undulating | Rice cultivation  | Eutric Fluvisols (FAO, 1988); | Ustic | Isohyperthermic |
| MB - SP2     | MB - Pa2 | Mbogo - Komtonga | Irrigation | Mvomero  | 9317507          | 373442         | 363      | Alluvial        | flat               | Sugarcane production  | Eutric Fluvisols              | Ustic | Isohyperthermic |
| MB - SP3     | MB - Pa3 | Mbogo - Komtonga | Irrigation | Mvomero  | 9317750          | 344231         | 362      | Alluvial        | flat to undulating | Rice cultivation, ploughed ready for transplanting/broadcasting | Eutric Fluvisols              | Ustic | Isohyperthermic |

183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202



## 203 2.3 Laboratory methods

204  
205 Laboratory methods used in the determination of different physico - chemical characteristics in the study area are  
206 summarized in Table 3.

207  
208 **Table 3:** Physical chemical analysis of the collected soil samples

| No | Parameter  | Analysis Method  | Reference  |
|----|--|--|--|
| 1  | Bulk density   | Core method  | Day, 1965  |
| 2  | Particle density   | Particle density was calculated using the mass of the solid particles and the volume they occupy. Mass of the solid particles was obtained by weighing the solid particles and likewise the volume was determined from the mass and density of water displaced by the sample | Soil Survey Staff, 2014                              |
| 3  | Total Porosity   | Total soil porosity was calculated by using the bulk and particle density data   |  |
| 4  | Soil moisture retention characteristics  | Sand Kaolin Box for low suction values and pressure apparatus for high suction values  | Okalebo, et al., 2002; Nelson and Sommers, 1982      |
| 5  | Particle size analysis   | Hydrometer method  | Nelson and Sommers, 1982; Bremner and Mulvaney, 1982 |
| 6  | Textural classes   | USDA textural triangle   | Thomas, 1996   |
| 7  | Soil pH  | Measured potentiometrically in water and 1 N KCl at a ratio of 1:2.5 weight to volume basis  | Nelson and Sommers, 1982; Chapman, 1965.             |
| 8  | Electrical conductivity (EC)   | Measured on a 1:2.5 soil: water suspension using electrical conductivity meter   | IUSS Working Group WRB, 2014.                        |
| 9  | Organic carbon   | Walkley and Black wet oxidation  | Obasi et al., 2015                                   |
| 10 | Organic matter   | By organic carbon conversion by multiplying with a factor of 1.724   | Khan et al., 2012                                    |
| 11 | Total N  | Micro-Kjeldahl digestion distillation  | Kebedey, et. al., 2015                               |
| 12 | Available phosphorus   | Bray and Kurtz-1 for low pH soils (pH water < 7) and Olsen for high pH soils (pH water > 7)  | Uwingabire, et al., 2016, Uwitonze, et al., 2016.    |
| 13 | Cation exchange capacity of soil (CEC soil) and exchangeable bases                             | Determined by saturating soil with neutral 1 M NH <sub>4</sub> OAc (ammonium acetate) and the adsorbed NH <sub>4</sub> <sup>+</sup> were displaced using 1 M KCl and then determined by Kjeldahl distillation  | [USDA-NRCS, 2016].                                   |
| 14 | Exchangeable bases (Ca <sup>2+</sup> , Mg <sup>2+</sup> , Na <sup>+</sup> and K <sup>+</sup> ) | Atomic absorption spectrophotometer (AAS)  | Lal and Shukla, 2005.                                |
| 15 | total exchangeable bases (TEB)   | Calculated arithmetically as the sum of the four exchangeable bases (Ca <sup>2+</sup> , Mg <sup>2+</sup> , Na <sup>+</sup> and K <sup>+</sup> ) for a given soil sample.   |  |

## 209 3 RESULTS AND DISCUSSION

### 210 3.1 Soil pedogeomorphological characteristics and genesis in Mbogo Komtonga Irrigation 211 Scheme 212 213 214

215 Salient morphological characteristics of the studied profiles are given in Table 4. All studied soil profiles  
216 were deep to moderately deep and water table was estimated at > 180 cm deep. Drainage was observed to  
217 be moderately well drained in profiles MB - P1, well drained at MB - P2 and well to moderately well drained  
218 at MB - P3. Floods were reported to be common in April - May. All profiles had clay (C) texture in its first  
219 horizon or the 0 – 30 cm soil depth and cracks were observed from the surface to 50 cm soil depth. With  
220 the exception of MB - P3 which had no fine sand (fS) texture in the third horizon (i.e. 50 – 90/120 cm), the  
221 rest of the master pits i.e. MB – P1 & P2 had fine sand (fS) texture material on third horizon. However, MB  
222 – P3 was more stratified compared with MB – P1 & P2 as C was underlain by sandy loam followed by Sand  
223 clay loam followed by gravelly sand material. All top soils of the studied master pits had brownish black  
224 colour. Whereas MB – P1 was dominated by brownish black colour in most of its horizons, MB – P2 had  
225 more mixed/complex colour and MB – P3 was dominated by dull yellowish brown colour. Consistency in  
226 these profiles ranged from hard when dry to soft or loose. For instance, profile MB – P1 & P2 had hard  
227 consistency when dry in the first two horizons compared with the only first horizon in profile MB – P2  
228 ascribed to the clay nature of the soil. Generally, the soil structure of the master pits was rated as strong to  
229 moderate medium and coarse sub angular blocky. But MB – P3 was dominated by weak medium and



230 coarse sub angular blocky. Infiltration rate was rated as moderate or moderate to high in profile MB – P3.  
231 Horizon boundary attributes varied within the pedons, whereby distinctness ranged from abrupt to gradual,  
232 but topography was dominantly smooth and wavy. Taken together, morphology and genesis of the studied  
233 soils were typical of alluvial soil formation.

UNDER PEER REVIEW

234 **Table 4:** Main pedogeomorphological features of the studied soil profiles in Mbogo Komtonga, Mvomero District, Tanzania  
 235

| Soil pedons <sup>1</sup> | Horizon             | Depth (cm)  | Texture | colour           |                  | Consistence |         |         | Structure     | Pores       | Roots          | Rock fragment        | Horizon boundary |
|--------------------------|---------------------|-------------|---------|------------------|------------------|-------------|---------|---------|---------------|-------------|----------------|----------------------|------------------|
|                          |                     |             |         | dry              | Moist            | Dry         | moist   | wet     |               |             |                |                      |                  |
| Mbogo MB-P1              | Ap                  | 0–30        | C       | brb (10YR 3/2)   | brb (10YR 2/2)   | h           | fr      | s & p   | str m+c sbk   | cm, cf+vf   | cf             | na                   | cs               |
|                          | Bw <sub>1t1</sub>   | 30–50       | C       | brb (10YR 3/2)   | brb (10YR 2/2)   | h           | fr      | s & p   | str m+c sbk   | cf+m, mf+vf | vff            | na                   | cs               |
|                          | Bw <sub>2</sub>     | 50 - 80     | fS      | na               | duybr (10YR 7/3) | fi          | fr      | s & p   | sg            | cm          |                | s freq irr qrtz + Fe | cs               |
|                          | B <sub>3W2t2</sub>  | 80–116      | C       | na               | brb (7.5YR 4/2)  | na          | fr      | s & p   | mo, m + f sbk |             | cm, mf+vf      | na                   | cs               |
|                          | B <sub>4W3</sub>    | 116–140     | SCL     | na               | brb (7.5YR 3/2)  | na          | fr      | ns & np | wk, m & f sbk | mf+vf       |                | na                   | cs               |
| B <sub>5W4</sub>         | 140–180             | fS          | na      | duybr (10YR 7/4) | na               | l           | ns & np | sg      | mf            |             | na             | na                   |                  |
| MB-P2                    | Ap                  | 0 – 27/32   | C       | gbr (7.5YR 4/2)  | brb (7.5YR 3/2)  | h           | fr      | s & p   | str m+f sbk   | cf+vf       | vfc, cm, mf+vf | na                   | cw               |
|                          | Bw <sub>1</sub>     | 27 – 60/74  | SC      | br (7.5YR 4/4)   | dbr (7.5YR 3/3)  | h           | fr      | ss & sp | mo m & c sbk  | vfm+vf      | cm, f+vf       | na                   | cw               |
|                          | Bw <sub>2</sub>     | 60 – 90/120 | fS      | gybr (10YR7/3)   | gybr (10YR 6/4)  | l           | na      | ns & np | wk sg         | mf          | ff + vf        | na                   | cw               |
|                          | Bw <sub>3t1g1</sub> | 90 – 160    | C       | duyo (10YR 7/4)  | g (10YR 4/1)     | h           | fr      | s & p   | str m+f sbk   | cf+vf       | vff            | na                   | cs               |
|                          | Bw <sub>4t2g2</sub> | 160 – 180   | SC      | na               | br (10YR 5/2)    | na          | fr      | s & p   | wk m+f sbk    | fm, cf+mvf  | na             | na                   | na               |
| MB-P3                    | Ap                  | 0 – 28      | C       | duybr (10YR5/3)  | brb (10YR3/2)    | h           | fr      | ss & sp | str m+c sbk   | mf+vf       | cf+mvf         | na                   | cs               |
|                          | Bw <sub>1</sub>     | 28 – 80     | SL      | duybr (10YR6/4)  | duybr (10YR5/4)  | s           | vfr     | ns & np | w m+c sbk     | mf+vf       | vfc+vff        | na                   | cw               |
|                          | Bw <sub>2g1</sub>   | 80 – 126    | SCL     | na               | gybr (10YR 6/4)  | na          | fr      | ns & np | w m+f sbk     | fm, mf+vf   | vff            | na                   | cs               |
|                          | Bw <sub>3g2</sub>   | 178 – 126   | SCL     | na               | duybr (10YR4/3)  | na          | fr      | ns & np | w m+c sbk     | cm, mf+vf   | vff+vfc        | na                   | cs               |
|                          | Bw <sub>4</sub>     | 178+        | grS     | na               | duybr (10YR7/3)  | na          | vfr     | ns & np | sg            | na          | na             | na                   | na               |

236 **Soil texture:** C = clay, SC = sandy clay, SCL= sandy clay loam, SL = sandy loam, fS = fine sand, grS = gravelly sand; **Soil colour:** brb = brownish black, duybr = dull yellowish  
 237 brown, g = gray; gbr = grayish brown; gybr = greyish yellowish brown, dbr = dark brown, br = brown; **Soil consistence:** **Dry:** s = soft; h = hard; l = loose; na = not applicable.  
 238 **Moist:** fr = friable; vfr = very friable; l = loose; **Wet:** ns = non-sticky; ss = slightly sticky; s = sticky; np = non-plastic; sp = slightly plastic; p = plastic. **Structure:** **Grade:** Str = strong;  
 239 mo = moderate; w = weak; sg = structure less single grained. **Size:** f = fine; m = medium; c = coarse. **Type:** sbk = sub angular blocky; **Pores:** Abundance: f = few; c = common; a  
 240 =abundant. **Size:** vf = very fine; f = fine; m = medium; common. **Roots:** **Colour:** ybr = yellowish brown; br = brown; dbr = dark brown; gybr = greyish yellowish brown; duyo = dull  
 241 yellowish orange; duybr = dull yellowish brown; g = grey; brb = brownish black; **Type:** Fe = iron; **Horizon boundary:** **Distinctness:** a = abrupt; c = clear; g = gradual.  
 242 **Topography:** s = smooth; w = wavy.

## 243 3.2 Soil physical characteristics

244

### 245 3.2.1 Soil texture, silt clay ratio, bulk density (BD) and total porosity (TP)

246

247 Table 5 presents data on soil physical properties including texture, silt clay ratio, bulk density and porosity  
 248 of the studied soils. Soil texture is the most stable physical characteristics of the soil. It influences a number  
 249 of other soil properties such as structure, consistence, bulk density (Bd), soil moisture regime, permeability,  
 250 root penetration, infiltration rates, runoff rate, erodibility, workability, root penetration and fertility. In Mbogo  
 251 – Komtonga soil profiles MB - P1 & P2 had clay top soils overlying clay or sand clay sub soils while profile  
 252 MB - P3 had clay topsoil overlying sandy loam sub soils. Generally, profile MB – P2 had much heavier  
 253 texture with the exception of the second horizon when compared with the other two master pits, implying it  
 254 would behave differently from these two in terms of physical and chemical properties. For example, clayey  
 255 texture is associated with high water retention capacity and high nutrient supply [30]. Profile MB - P2 would  
 256 probably offer more favourable conditions for paddy than the other two profiles. Clay content decreased  
 257 more or less with depth in all pedons providing some indication of non-uniformity in clay eluviation -  
 258 illuviation. Silt/clay ratio, an indicator of soil susceptibility to erosion was less than the threshold of 0.4 [31],  
 259 as in profile MB – P1 implying moderate resistance to erosion. Silt/clay ratios in the other two profiles i.e.  
 260 MB – P2 & P3, showed higher values than the threshold value.

261

262 **Table 5.** Some selected physical properties of master pits from Mbogo irrigation scheme

263

| Profile No.  | Horizons            | Horizons Depth (cm) | <2 | 20-50             | 50-2000 | Textural Class | Si/C ratio | Bulk Density | Particle Density | Total Porosity |
|--------------|---------------------|---------------------|----|-------------------|---------|----------------|------------|--------------|------------------|----------------|
|              |                     |                     |    | ( $\mu\text{m}$ ) |         |                |            |              |                  |                |
| <b>Mbogo</b> |                     |                     |    |                   |         |                |            |              |                  |                |
| MB-P1        | Ap                  | 0–30                | 58 | 24                | 18      | C              | 0.41       | 1.209        | 2.65             | 54.4           |
|              | Bw <sub>1t1</sub>   | 30–50               | 66 | 2                 | 32      | C              | 0.03       | 1.216        | 2.65             | 54.1           |
|              | B <sub>2</sub>      | 50 - 80             | 6  | 2                 | 92      | fS             | 0.33       | 1.684        | 2.65             | 36.5           |
|              | B <sub>3w2t2</sub>  | 80 –116             | 72 | 12                | 16      | C              | 0.17       | 1.501        | 2.65             | 43.4           |
|              | B <sub>4w3</sub>    | 116 –140            | 24 | 2                 | 74      | SCL            | 0.08       | 1.446        | 2.65             | 45.4           |
|              | B <sub>5w4</sub>    | 140 –180            | 6  | 2                 | 92      | fS             | 0.33       | 1.684        | 2.65             | 36.5           |
| MB-P2        | Ap                  | 0 – 27/32           | 60 | 12                | 28      | C              | 2.33       | 1.223        | 2.65             | 53.8           |
|              | Bw <sub>1</sub>     | 27 – 60/74          | 34 | 8                 | 58      | SC             | 7.25       | 1.364        | 2.65             | 48.5           |
|              | Bw <sub>2</sub>     | 60 – 90/120         | 6  | 0                 | 94      | fS             | 0.00       | 1.688        | 2.65             | 36.3           |
|              | Bw <sub>3t1g1</sub> | 90 –160             | 62 | 8                 | 30      | C              | 3.75       | 1.222        | 2.65             | 53.9           |
|              | Bw <sub>4t2g2</sub> | 160 –180            | 44 | 6                 | 50      | SC             | 8.33       | 1.311        | 2.65             | 50.5           |
| MB-P3        | Ap                  | 0 – 28              | 56 | 12                | 32      | C              | 2.67       | 1.241        | 2.65             | 53.2           |
|              | Bw <sub>1</sub>     | 28 – 80             | 18 | 6                 | 76      | SL             | 12.67      | 1.492        | 2.65             | 43.7           |
|              | Bw <sub>2g1</sub>   | 80 – 126            | 20 | 6                 | 74      | SCL            | 12.33      | 1.472        | 2.65             | 44.5           |
|              | Bw <sub>3g2</sub>   | 126 – 178           | 22 | 8                 | 70      | SCL            | 8.75       | 1.451        | 2.65             | 45.2           |
|              | Bw <sub>4</sub>     | 178+                |    |                   |         | grS            |            |              | 2.65             | 100.0          |

264 C = clay, SC = sandy clay, SCL= sandy clay loam, SL = sandy loam, fS = fine sand, grS = gravelly sand

265

266 Changes in bulk density for a given soil can alert soil managers and or extension agents to changes in soil  
 267 quality and ecosystem function. Bulk density reflects the soil's ability to function for structural support, water  
 268 and solute movement, and soil aeration. It is also used to express soil physical, chemical and biological  
 269 measurements on a volumetric basis for soil quality assessment and comparisons between management  
 270 systems. Bulk densities above thresholds indicate impaired function. Generally, in highly productive soils,  
 271 Bd range from 1.0 – 1.5 g cm<sup>-3</sup> (i.e. fine to medium texture) and 1.10 to 1.65 g cm<sup>-3</sup> (i.e. coarse textured

272 soils) also see Table 6 with potential root restriction occurring at  $\geq 1.4 \text{ g cm}^{-3}$  for clay and  $\geq 1.6 \text{ g cm}^{-3}$  for  
273 sandy soils [32]. In Mbogo - Komtonga study area, soil texture in most of the topsoil of the representative  
274 profiles was dominated by clay (C) or clay loam (CL) or sand clay loam (SCL). Bulk density (Bd) and total  
275 porosity (Pt) are very important factors in the determination of root penetration and proliferation. In some  
276 soil profile horizons, soil texture is used to determine the soil Bd that is used to calculate the total porosity.  
277 Whereas the Bd of the surface soils in Mbogo - Komtonga it range from  $1.21 \text{ g cm}^{-3}$  -  $1.24 \text{ g cm}^{-3}$ , Bd  
278 ranged from 1.21 to 1.68 in profile MB – P1; 1.22 – 1.69 in MB – P2 and  $1.24 - 1.49 \text{ g cm}^{-3}$  in MB – P3.  
279 These correspond to total porosity of 36.5 – 54.4 %; 36.3 – 53.9 % and 43.7 – 53.2 % respectively. The  
280 data showed that the Bd increased with depth [33; 34; 35] and was medium in 67 % (slightly above  
281 adequate but not restrictive) and ideal for plant growth in 33 % of the study area. Similarly, the data  
282 indicated that the lower the Bd, the higher the porosity and vice versa (see Table 5). High bulk density (Bd)  
283 is an indicator of low soil porosity and soil compaction; poor environment for root growth, reduced aeration  
284 and undesirable changes in hydrologic function such as reduced water infiltration rates [33; 36; 37]. The  
285 comparatively higher Bd in 67 % of the study areas in Mbogo was probably due to less aggregation, clay  
286 (heavy) textural class of the area, fewer roots and compaction caused by the overlaying layers. Similarly,  
287 higher Bd can be caused by consistently ploughing or disking to the same depth; allowing equipment traffic  
288 especially on wet soil; using a limited crop rotation without variability in root structure or rooting depth; and  
289 incorporating, burning, or removing crop residues. In order to reduce the chances of high bulk density and  
290 compaction, soil disturbance and production activities when soils are wet should be minimized, field/farm  
291 roads for farm equipment should be designed and constructed, sub-soiling to disrupt existing compacted  
292 layers, and practices that maintain or increase SOM should be adopted at least once in three years

### 293 294 **3.2.2 Field Capacity, Permanent Wilting Point, Available Water and Available Water Capacity,**

295  
296 Field capacity (FC) is the water remaining in soil after it has been thoroughly saturated and allowed to drain  
297 freely, usually for one to two days (Table 6). Permanent wilting point (PWP) is the moisture content of soil  
298 at which plants wilt and fail to recover when supplied with sufficient soil water. It is an indicator of soil's  
299 ability to retain water and make it sufficiently available for plant use. Available water capacity (AWC),  
300 usually expressed as a volume fraction, percentage, or depth (cm), is the maximum amount of water held in  
301 soil between its field capacity (at pF 2.0) and permanent wilting point (at pF 4.2). In Mbogo – Komtonga,  
302 AWC range between  $107 \text{ mm m}^{-1}$  and  $144 \text{ mm m}^{-1}$  rated as medium (Table 7). However, water storage  
303 capacity (AWSC) was  $200 \text{ mm m}^{-1}$  (inferred) considered as good and sufficient for paddy production and  
304 other upland crops. ~~AWC is used to develop water budgets, predict droughtiness, design and operate  
305 irrigation systems, design drainage systems, protect water resources, and predict yields. Water availability  
306 (WA) is an important indicator because plant growth and soil biological activity depend on water for  
307 hydration and delivery of nutrients in solution. Runoff and leaching volumes are also determined by storage  
308 capacity and pore size distribution.~~ Lack of AW reduces root and plant growth, and can lead to plant death  
309 if sufficient moisture is not provided before PWP. Poor AW is caused by conventional tillage operations; low  
310 residue crop rotations, and burning, burying, harvesting, or otherwise removing plant residues; heavy  
311 equipment traffic on wet soils, and grazing systems that allow development of livestock loafing areas and  
312 livestock trails. In order to improve AWC in soils, farmers should grow high residue crops, cover crops,  
313 reduce soil disturbing activities, and manage residue to protect and increase SOM. When feasible, tillage,  
314 harvest, and other farming operations requiring heavy equipment can be avoided when the soil is wet so as  
315 to minimize compaction. Compacted layers can be ripped to break them and expand the depth of the soil  
316 available for root growth.

317

318 **Table 6:** Soil physical characteristics of the selected sites in Mbogo - Komtonga proposed irrigation schemes  
 319

| Location       | MU      | $B_d$            | $P_d$ | $P_t$ | Texture | FC  | PWP | AW  | AWSC |
|----------------|---------|------------------|-------|-------|---------|-----|-----|-----|------|
|                |         | ( $g\ cm^{-3}$ ) | (%)   | (%)   |         |     |     |     |      |
| Mbogo Komtonga | MB - P1 | 1.21             | 2.65  | 54.4  | C       | 426 | 285 | 140 | 200  |
|                | MB - P2 | 1.22             | 2.65  | 53.8  | C       | 381 | 273 | 107 | 200  |
|                | MB - P3 | 1.24             | 2.65  | 53.2  | C       | 386 | 275 | 111 | 200  |

320 MU = Mapping Unit;  $B_d$  = Bulk density;  $P_d$  = Particle density; WSC = Water storage capacity; AW =  
 321 Available water; HC = Hydraulic conductivity;  $P_t$  = Total Porosity; SL = Sandy loam; SCL = Sand clay loam;  
 322 CL = Clay loam; C = Clay.

323

### 324 3.3. Chemical properties of the studied pedons in Mbogo Komtonga, Mvomero District

325

#### 326 3.3.1 Soil pH

327

328 Results of soil pH and other chemical properties of the soils of the studied representative master pits of  
 329 Mbogo Komtonga are presented in **Table 7**. Soil pH influences the rate of plant nutrient release by  
 330 weathering, suitability of all materials in the soil, and amount of nutrients ions stored on the cation  
 331 exchange complex **due to the fact that pH affects the form of nutrient ions in soils thus affecting plant**  
 332 **availability**. Before nutrients can be used by plants they must be dissolved in the soil solution. The pH is  
 333 therefore a good guide for predicting which plant nutrients are deficient. Soils tend to become acidic as a  
 334 result of (1) rainwater leaching away basic ions (Ca, Mg, K and Na); (2) formation of a weak organic acid as  
 335 a result of  $CO_2$  from decomposing OM and root respiration dissolving in soil water; (3) formation of strong  
 336 organic and inorganic acids, such as nitric ( $HNO_3$ ) and sulphuric acid ( $H_2SO_4$ ), from decaying OM and  
 337 oxidation of ammonium ( $NH_3$ ) and sulphur (S) fertilizers. Strongly acid soils are usually the result of the  
 338 action of these strong organic and inorganic acids. The pH of top soils of the studied soil profiles in Mbogo -  
 339 Komtonga irrigation scheme ranged from 5.4 to 6.0. This was rated as strongly acid to medium acid [38].  
 340 Similarly, the pH of the sub soils ranged from 5.8 – 6.2 and was rated as medium acid to slightly acid [36;  
 341 38]. The strong to medium acid observed in the tops soils of these profiles could be ascribed to low amount  
 342 of bases by leaching during water table fluctuations and water percolation during flooding periods and soil  
 343 nutrients mining [36; 39; 40]. The data also showed that pH increased with depth in the studied profiles as  
 344 likewise reported in [41; 42]. The nature of the observed acidity in the top soils of the representative profiles  
 345 threatens the availability of mineral elements such as P which is readily available in soils with pH centred at  
 346 6.5. For example under low pH, P is precipitated due to dissolution of Al and Fe mineral elements leading  
 347 to its fixation and further soil pH depression [39]. However, most plant mineral elements are available in the  
 348 pH range of approximately 6.5 – 7.0 [43]. Similarly, soil pH can influence plant growth by its effect on the  
 349 activity of beneficial micro-organisms [44]. For example, bacteria that decompose SOM are hindered in  
 350 strong acid soils which in turn prevent OM from breaking down. As a result, OM is accumulated un-  
 351 decomposed or unbroken consequently tying up of nutrients such as N making them unavailable to plants.  
 352 In order to reverse this trend, it is recommended to carryout liming in such soils by using limestone/calcium  
 353 carbonate ( $CaCO_3$ ) at a rate of 3 - 4 t  $ha^{-1}$  to raise the pH from the current status ( $5.4 \leq pH \leq 6.0$ ) to a pH  
 354 range of between 6.5 – 7.0 (Hausenbuiller,1978). Other material that can also be used is calcium oxide  
 355 ( $CaO$ ) also known as quick lime with Calcium Carbonate Equivalent (CCE) of 179 %. Although this material  
 356 gives quick results, care should be taken as it is difficult to apply for it irritates the eyes. Electrical  
 357 conductivity (EC) is a measure of relative salt concentration or salinity, and too much salt in the soil can  
 358 interfere with root function and nutrient uptake [45; 46]. EC values of the top soils ranged between 0.23

359 and 0.32 (dS m<sup>-1</sup>) and 0.02 – 0.04 (dS m<sup>-1</sup>) in the sub soils of the studied master pits horizons implying that  
360 all the soils were non-saline.

361

### 362 **3.3.2 Organic Carbon**

363

364 Organic carbon (OC) or Soil Organic Matter (SOM) in the soil is important because humidified OM  
365 molecules may react with mineral colloids and contribute to the stabilization of soil aggregates. While SOM  
366 favours water retention capacity and adsorption of fulvic and humic compounds by Fe<sup>2+</sup> and Al<sup>3+</sup> oxide, it  
367 also prevents their crystallization hence decreasing fixation power with regards to phosphates at  
368 unfavourable pH values. SOM provides much of the CEC, and, surface soils contain large quantity of plant  
369 nutrients with storehouse considered as slow release of nutrient especially so by N. Results of organic  
370 carbon (OC) determination from the top soil (0 - 30 cm) of the representative master pits in Mbogo -  
371 Komtonga ranged from 24.7 g kg<sup>-1</sup> to 40.0 g kg<sup>-1</sup> (**Table 8**). This corresponds to 42.5 g kg<sup>-1</sup> to 69.3 g kg<sup>-1</sup>  
372 SOM. Organic carbon in most of the profiles showed systematic trend of decreasing with depth. Since SOM  
373 content was calculated from SOC [47], these parameters **have** similar trend. It is generally accepted that a  
374 threshold for SOM in most soils is 34 g kg<sup>-1</sup> below which decline in soil quality is expected to occur [48].  
375 With the observed data all values were above the proposed threshold limits, suggesting that no decline in  
376 soil quality for Mbogo - Komtonga irrigation scheme [49]

377

378

UNDER PEER REVIEW

379 **Table 7:** Some selected chemical properties of master pits from Mbogo irrigation scheme  
 380

| Profile No.  | Horizons            | Horizons Depth<br>(cm) | pH               |     | EC<br>dS.m <sup>-1</sup> | OC   | OM<br>(%) | N    | C/N<br>ratio | Av. P<br>mg.kg <sup>-1</sup> |
|--------------|---------------------|------------------------|------------------|-----|--------------------------|------|-----------|------|--------------|------------------------------|
|              |                     |                        | H <sub>2</sub> O | KCl |                          |      |           |      |              |                              |
| <b>Mbogo</b> |                     |                        |                  |     |                          |      |           |      |              |                              |
| MB-P1        | Ap                  | 0–30                   | 5.8              | 4.9 | 0.32                     | 3.80 | 6.54      | 0.20 | 21           | 4.80                         |
|              | Bw <sub>1t1</sub>   | 30–50                  | 6.0              | 4.6 | 0.10                     | 1.16 | 2.00      | 0.13 | 9            | 2.46                         |
|              | B <sub>2</sub>      | 50 - 80                | 6.7              | 5.1 | 0.02                     | 0.07 | 0.12      | 0.05 | 1            | 1.06                         |
|              | B <sub>3w2t2</sub>  | 80 –116                | 6.4              | 4.8 | 0.10                     | 0.97 | 1.67      | 0.10 | 10           | 0.88                         |
|              | B <sub>4w3</sub>    | 116 –140               | 6.6              | 5.1 | 0.04                     | 0.65 | 1.12      | 0.07 | 9            | 3.82                         |
|              | B <sub>5w4</sub>    | 140 –180               | 6.8              | 5.6 | 0.04                     | 0.09 | 0.15      | 0.04 | 2            | 0.80                         |
| MB-P2        | Ap                  | 0 – 27/32              | 6.0              | 5.2 | 0.24                     | 4.03 | 6.93      | 0.18 | 23           | 5.47                         |
|              | Bw <sub>1</sub>     | 27 – 60/74             | 6.2              | 4.7 | 0.05                     | 0.55 | 0.95      | 0.11 | 5            | 0.87                         |
|              | Bw <sub>2</sub>     | 60 – 90/120            | 6.6              | 5.0 | 0.02                     | 0.07 | 0.12      | 0.04 | 2            | 1.62                         |
|              | Bw <sub>3t1g1</sub> | 90 –160                | 6.2              | 4.3 | 0.05                     | 0.98 | 1.69      | 0.17 | 6            | 1.91                         |
|              | Bw <sub>4t2g2</sub> | 160 –180               | 6.4              | 4.6 | 0.05                     | 0.55 | 0.95      | 0.09 | 6            | 3.70                         |
|              | MB-P3               | Ap                     | 0 – 28           | 5.4 | 4.3                      | 0.23 | 2.47      | 4.25 | 0.15         | 17                           |
|              | Bw <sub>1</sub>     | 28 – 80                | 5.8              | 4.3 | 0.04                     | 0.36 | 0.62      | 0.07 | 5            | 2.24                         |
|              | Bw <sub>2g1</sub>   | 80 – 126               | 5.7              | 4.2 | 0.04                     | 0.43 | 0.74      | 0.07 | 6            | 2.54                         |
|              | Bw <sub>3g2</sub>   | 126 – 178              | 5.6              | 4.1 | 0.05                     | 1.08 | 1.86      | 0.06 | 17           | 1.79                         |

381 pH = soil reaction, EC = Electrical conductivity, OC= Organic carbon, OM = Organic matter, N = Nitrogen,  
 382 C/N = Carbon Nitrogen ratio, Av.P = Available P

383  
 384 **3.3.3 Total Nitrogen**  
 385

386 Inadequate amount of N in the soil is the primary factor that limits plant growth and development in many  
 387 parts of the world [50; 51). Nitrogen levels in the studied soil Master pits were low to medium with values  
 388 ranging from 1.5 – 2.0 g kg<sup>-1</sup> in top soils and 0.4 – 0.6 g kg<sup>-1</sup> in the sub soils. These values were rated as  
 389 very low to low [38]. According to [38] guidelines, the proposed threshold value for N in most crops in  
 390 Tanzania is 2 g kg<sup>-1</sup> soil. The results show that of the studied Master pits only MB – P1 had N which was at  
 391 least on the threshold value but MB – P2 & P3 were below the threshold value. The observed low or  
 392 medium N in the surveyed areas may probably be influenced by microbial activity in the soil and the very  
 393 low or low soil pH [49; 52; Table 9]. So, any activity envisaged to improve the soil pH, SOM quality as well  
 394 as microbial activities can, consequently, lead to an increase in N in the soil [49]. The low to very low levels  
 395 of N in the surveyed areas suggests application of ammoniocal form of N, which resists better to leaching  
 396 caused by rainfall or irrigation as the case may be in the surveyed areas. As far as humification is  
 397 concerned, an average C/N ratio of 10 (i.e. 8 - 12) is considered as optimal [36, 46]. The C/N ratio of top  
 398 soils of the representative master pits ranges from 17 – 21 and was rated as moderate to poor quality  
 399 SOM. It is generally accepted that C/N ratios between 8 and 12 are considered to be the most favourable,  
 400 implying a relatively fast mineralisation of N from the organic materials. With the exception of MB-P3 which  
 401 registered C/N ratio of 17 rated as moderate or medium quality SOM, the rest of the representative Master  
 402 pits (MB – P1 & P2) had C/N ratio outside the suggested range and were rated as poor quality SOM.  
 403 However, the C/N ratio observed in the sub soils of all Master pits ranged from 1 – 10 in MB – P1, 2 – 6 in  
 404 MB – P2 and 5 - 17 in MB – P3 which was rated as medium and good quality SOM. According to [36] and  
 405 [46], C/N ratio of 10:1 indicates good quality organic material, although they cautioned that C/N ratio might  
 406 not be a good indicator of soil fertility, and thus encouraged use of individual C and N values instead.



### 407 3.3.4 Available Phosphorus (Pav)

408

409 The data from the top soil of the representative Master pits (MB – P1, P2 & P3) in Mbogo - Komtonga  
410 irrigation scheme shows that available P range from 0.87 – 5.47 mg kg<sup>-1</sup> rated as low (Table 9). Likewise  
411 the data in the sub soils range from 0.80 – 3.82 mg kg<sup>-1</sup> also rated as low. Phosphorus (P) is an essential  
412 macro element for plant growth, hence an important soil fertility indicator. In agriculture, management of P  
413 is second only to management of N in its importance for the production of healthy and profitable crop  
414 yields. An average P level of 7 mg kg<sup>-1</sup> is considered optimal below which P deficiency symptoms are likely  
415 to occur in most crops. Based on the generally accepted threshold P level, all the observed P values in  
416 Mbogo - Komtonga are considered to be below the critical range and will definitely need measures to  
417 reverse the trend. The generally low P availability manifested in all the mapping units in Mbogo - Komtonga  
418 (Table 9) suggests that management of P in these areas is critical for sustainable agricultural development.

419

### 420 3.3.5 Exchangeable bases, cation exchange capacity and per cent base saturation

421

422 Results of exchangeable bases, cation exchange capacity and per cent base saturation in the  
423 representative Master pits in Mbogo Komtonga are presented in Table 8. Potassium (K) in the top soils  
424 ranged from 0.62 cmol (+) kg<sup>-1</sup> (MB - P3) to 2.97 cmol (+) kg<sup>-1</sup> (MB - P2) rated as medium to very high. In  
425 the sub soils, exchangeable K ranged from 0.03 (MB – P1) – 0.06 cmol (+) kg<sup>-1</sup> (MB – P3) and were rated  
426 as low to very low. In general terms, a response to K fertilizers is likely when a soil has an exchangeable K  
427 value of < 0.2 cmol (+) kg<sup>-1</sup> soil and unlikely when it is above 0.4 cmol (+) kg<sup>-1</sup> soil [Table 10; 38; 53]. The  
428 data shows that K is unlikely to respond to Mbogo – Komtonga Irrigation scheme. Exchangeable Ca<sup>2+</sup> in  
429 the topsoil of the representative Master pits ranged from 12.6 cmol (+) kg<sup>-1</sup> (MB – P3) – 29.64 cmol (+) kg<sup>-1</sup>  
430 (MB – P2) rated as high to very high. In the sub soils, Ca<sup>2+</sup> ranged from 0.0 (MB – P3) 2.2 (MB – P1) rated  
431 as very low to low. [54] Proposed that in most of the crops, the recommended threshold level of Ca<sup>2+</sup> is 5  
432 cmol (+) kg<sup>-1</sup>. It is generally acknowledged that field conditions that limit Ca<sup>2+</sup> uptake produce lower crop  
433 yields compared with field conditions that do not limit Ca<sup>2+</sup> uptake [55]. Based on the critical limits, and Ca<sup>2+</sup>  
434 levels at the top soils, it is unlikely to have Ca<sup>2+</sup> deficient of for most crops as it lies below the proposed  
435 critical limits. Exchangeable Mg<sup>2+</sup> in top soils of the representative Master pits in Mbogo - Komtonga range  
436 from 4.25 cmol (+) kg<sup>-1</sup> (MB – P1) – 5.07 cmol (+) kg<sup>-1</sup> (MB – P2), rated as high to very high. In the sub  
437 soils, Mg ranged from 0.38 (MB – P2) – 0.59 (MB – P1) rated as low to medium [38]. The recommended  
438 value of Mg<sup>2+</sup> in most crops is 2 cmol (+) kg<sup>-1</sup> [56]. These data suggests that based on the top soil data, the  
439 studied area have sufficient Mg<sup>2+</sup> supplies for crop growth even though there is irregular decrease of  
440 exchangeable Mg with depth. Topsoil exchangeable Na<sup>+</sup> and or exchangeable sodium percentage (ESP)  
441 the levels as well as the electrical conductivity (EC) in the representative Master pits (MB – P1, P2 & P3) in  
442 the study area are presented in Table 10. The results indicates that the levels of Na<sup>+</sup> in the top soils  
443 corresponds to 0.17 (MB – P2) – 0.45 cmol (+) kg<sup>-1</sup> (MB – P3). These values were rated as low to medium  
444 [31]. Exchangeable Na in the sub soils ranged from 0.08 (MB – P2) – 0.38 (MB – P1) rated as very low to  
445 low [38]. The values of Na beyond which crop growth and development is impaired is less than 1 cmol (+)  
446 kg<sup>-1</sup> [31]. The corresponding ESP range from 0.4 – 1.7 % rated as non-sodic. The critical values of ESP  
447 above which most crops are affected are established at 15 % [57]. These results suggest that the surveyed  
448 areas have no threat to sodicity problems [31; 46].

449

450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473

### 3.3.6 Cation exchange capacity (CEC)

Cation exchange capacity (CEC) refers to the exchange phenomenon of positively charged ions (cation) at the surface of the negatively charged colloids [58]. It is often used as a characteristic in the determination of the nutrient retention soil quality. The higher the CEC, the more capable the soil is to retain nutrients. High CEC means more nutrients are held on the soil, decreasing their mobility and uptake whereas low CEC means that more nutrients are in the soil solution, making them available to plants but also increasing the likelihood of leaching. Studies have shown that soils with CEC values of between 6 - 12 cmol (+) kg<sup>-1</sup> soil are poor in exchangeable bases [38]. CEC values in the topsoil of the representative Master pits in Mbogo - Komtonga irrigation scheme are as shown in Table 8. Results showed that CEC values ranged between 27.02 cmol (+) kg<sup>-1</sup> (MB - P3) – 44.8 cmol (+) kg<sup>-1</sup> (MB - P2) and were rated as high to very high [38]. The high to very high CEC could be related to the clay mineral and soil organic matter (SOM) or organic carbon (OC) present in these soils. However, it is recommended to apply both manure/compost manure and the required amount of inorganic fertilizer. By adding inorganic fertilizer, one increases the humus content of the soil and consequently resulting into a higher or maintenance of higher CEC hence a better retention of nutrients. The data also showed that percent base saturation (BS) of the representative Master pits varied irregularly and the trend with depth was not clear within the soil profiles. However, the top soils of the representative Master Pits recorded relatively higher topsoil values in MB - P1 and MB - P2 than in MB - P3. Based on [46], % BS in all the representative Master pits were rated as high and fertile soils because the BS were greater than 60 %.

**Table 8:** Exchangeable cations and related properties the studied soils.

| Profile No.  | Horizons                         | Horizons Depth (cm) | Exchangeable bases |      |      |      |      | CECsoil | BS (%) |
|--------------|----------------------------------|---------------------|--------------------|------|------|------|------|---------|--------|
|              |                                  |                     | Ca                 | Mg   | K    | Na   | TEB  |         |        |
| <b>Mbogo</b> |                                  |                     |                    |      |      |      |      |         |        |
| MB-P1        | Ap                               | 0–30                | 22.46              | 4.25 | 1.15 | 0.39 | 28.3 | 41.0    | 68     |
|              | Bw <sub>1t1</sub>                | 30–50               | 20.36              | 4.57 | 0.28 | 0.38 | 25.6 | 32.4    | 77     |
|              | B <sub>2</sub>                   | 50 - 80             | 2.99               | 0.67 | 0.03 | 0.14 | 3.8  | 10.5    | 68     |
|              | B <sub>3</sub> W <sub>2t2</sub>  | 80–116              | 18.66              | 4.53 | 0.28 | 0.35 | 23.8 | 30.2    | 77     |
|              | B <sub>4</sub> W <sub>3</sub>    | 116–140             | 8.58               | 2.06 | 0.10 | 0.24 | 11.0 | 18.2    | 71     |
|              | B <sub>5</sub> W <sub>4</sub>    | 140–180             | 2.20               | 0.59 | 0.03 | 0.14 | 3.0  | 7.8     | 82     |
| MB-P2        | Ap                               | 0 – 27/32           | 29.64              | 5.07 | 2.97 | 0.17 | 37.9 | 44.8    | 82     |
|              | Bw <sub>1</sub>                  | 27 – 60/74          | 9.58               | 2.80 | 0.12 | 0.26 | 12.8 | 18.0    | 78     |
|              | Bw <sub>2</sub>                  | 60 – 90/120         | 1.80               | 0.38 | 0.04 | 0.08 | 2.3  | 7.0     | 84     |
|              | Bw <sub>3t1</sub> g <sub>1</sub> | 90 –160             | 14.97              | 4.61 | 0.21 | 0.36 | 20.2 | 24.8    | 79     |
|              | Bw <sub>4t2</sub> g <sub>2</sub> | 160 –180            | 10.18              | 3.78 | 0.17 | 0.33 | 14.5 | 20.6    | 71     |
| MB-P3        | Ap                               | 0 – 28              | 12.57              | 5.02 | 0.62 | 0.45 | 18.7 | 27.0    | 66     |
|              | Bw <sub>1</sub>                  | 28 – 80             | 0.00               | 1.61 | 0.10 | 0.14 | 1.9  | 6.9     | 62     |
|              | Bw <sub>2</sub> g <sub>1</sub>   | 80 – 126            | 6.59               | 1.89 | 0.08 | 0.09 | 8.7  | 16.4    | 66     |
|              | Bw <sub>3</sub> g <sub>2</sub>   | 126 – 178           | 6.19               | 1.78 | 0.06 | 0.18 | 8.2  | 16.6    | 63     |

474 Ca = Calcium, Mg = Magnesium, K = Potassium, Na = Sodium, TEB = Total exchangeable bases, CEC = Cation exchange  
475 capacity, BS = Base saturation

476 **3.3.7 Nutrient balance**

477  
 478 The availability of nutrients for uptake by plants depends not only upon absolute levels but also on relative  
 479 amounts of individual elements. According to [45], a good trend is with  $\text{Ca}^{2+}$  higher than  $\text{Mg}^{2+}$ , and  $\text{Mg}^{2+}$   
 480 higher than  $\text{K}^+$  (i.e.  $\text{Ca}^{2+} > \text{Mg}^{2+} > \text{K}^+$ ). With the exception of  $\text{Na}^+$  that was more or less greater than  $\text{K}^+$ , the  
 481 exchangeable cations in the Mbogo – Komtonga irrigation scheme followed that trend:  $\text{Ca}^{2+} > \text{Mg}^{2+} > \text{K}^+ <$   
 482  $\text{Na}^+$  at MB - P1, P2 & P3 (Table 9). A similar observation has been reported elsewhere by [59; 60; 61 and  
 483 62]. Results from this study showed that the Ca/TEB ratios in the topsoil of the representative Master pits  
 484 ranged from 0.67 to 0.80 (Table 9). With the exception of MB – P3 Master pit which did not show any  
 485 decreasing trend with depth, the data showed a generally decreasing trend in MB – P1 & P2. According to  
 486 [31] Ca/TEB critical ratios beyond which the uptake of Mg, K and other bases are affected is pegged at  
 487 more than 0.5 and Ca induced deficiency of Mg and/or K become clear. This result suggests that Ca may  
 488 alter uptake and induce deficiency of K at all the studied Master pits (i.e. MB – P1, P2 & P3). Ca/Mg ratios  
 489 ranged from 2.5 to 5.9 in the top soils and generally decreasing with depth in MB – P1 & P2 but not in MB  
 490 – P3 (Table 9) suggesting that Ca content was greater in top soils compared with the sub soils [31]. Studies  
 491 has shown that a critical range of between 2 and 4 in Ca/Mg ratio was considered as optimal for plant  
 492 growth [38] suggesting that only MB – P3 has values within the critical range. With regards to Mg/K ratios  
 493 the top soils of the representative Master Pits ranged from 1.7 to 8.1 and showed irregular pattern with soil  
 494 depth. This result suggests that MB – P1 & P2 are within the recommended critical range for optimal  
 495 nutrient uptake by plants [31]. The percentage K/TEB ratios ranged between 3.3 (MB – P3) to 7.9 % (MB –  
 496 P2) and decreased with soil depth in the Master pit. These top soil values were above 2 % (Table 9) and  
 497 were considered favourable for most tropical crops just as reported in [31]. As the K/TEB ratios are greater  
 498 than 2 % in all the top soils of the studies Master pits, problems of K – deficiency is unlikely [63]. However,  
 499 sub surface K/TEB ratio was probably lower and K deficiency is likely to happen.

501 **Table 9:** Nutrient ratios of the studied soils.

502

| Profile No.  | Horizons                         | Horizons Depth (cm) | Ca/TEB | Ca/Mg | Mg/K  | % (K/TEB) |
|--------------|----------------------------------|---------------------|--------|-------|-------|-----------|
| <b>Mbogo</b> |                                  |                     |        |       |       |           |
| MB-P1        | Ap                               | 0–30                | 0.80   | 5.28  | 3.70  | 4.07      |
|              | Bw <sub>1t1</sub>                | 30–50               | 0.80   | 4.46  | 16.32 | 1.09      |
|              | B <sub>2</sub>                   | 50 - 80             | 0.78   | 4.46  | 22.33 | 0.78      |
|              | B <sub>3</sub> W <sub>2t2</sub>  | 80 –116             | 0.78   | 4.12  | 16.18 | 1.18      |
|              | B <sub>4</sub> W <sub>3</sub>    | 116 –140            | 0.78   | 4.17  | 20.60 | 0.91      |
|              | B <sub>5</sub> W <sub>4</sub>    | 140 –180            | 0.74   | 3.73  | 19.67 | 1.01      |
| MB-P2        | Ap                               | 0 – 27/32           | 0.78   | 5.85  | 1.71  | 7.85      |
|              | Bw <sub>1</sub>                  | 27 – 60/74          | 0.75   | 3.42  | 23.33 | 0.94      |
|              | Bw <sub>2</sub>                  | 60 – 90/120         | 0.78   | 4.74  | 9.50  | 1.74      |
|              | Bw <sub>3t1</sub> g <sub>1</sub> | 90 –160             | 0.74   | 3.25  | 21.95 | 1.04      |
|              | Bw <sub>4t2</sub> g <sub>2</sub> | 160 –180            | 0.70   | 2.69  | 22.24 | 1.18      |
| MB-P3        | Ap                               | 0 – 28              | 0.67   | 2.50  | 8.10  | 3.32      |
|              | Bw <sub>1</sub>                  | 28 – 80             | 0.00   | 0.00  | 16.10 | 5.41      |
|              | Bw <sub>2</sub> g <sub>1</sub>   | 80 – 126            | 0.76   | 3.49  | 23.63 | 0.92      |
|              | Bw <sub>3</sub> g <sub>2</sub>   | 126 – 178           | 0.75   | 3.48  | 29.67 | 0.73      |

503 Ca/TEB = Calcium to Total exchangeable bases, Ca/Mg = Calcium to Magnesium ratio, Mg/K = Magnesium to Potassium ratio, K/TEB =  
 504 Potassium to Total Exchangeable Bases

505 **3.3.8 Soil classification**

506

507 Based on the field and laboratory data, the soils were classified as Eutric Fluvisol and Vertic Cambisol in  
 508 the FAO soil classification system [28; Table 10]

509

510 **Table 10:** Classification of the studied soil Master Pits in Mbogo Komtonga irrigation scheme, Mvomero  
 511 District, Tanzania.

512

| PROFILE | Diagnostic horizons | Other diagnostic features  | FAO UNESCO SOIL MAP OF THE WORLD CLASSIFICATION (1988) |                     |                         |
|---------|---------------------|--|--|---------------------|-------------------------|
|         |                     |  | Soil unit  | Major soil grouping | Soil subunits           |
| MB - P1 | Ochric A, Argic B   | Flat to undulating, deep to moderately deep, moderately well drained, brownish black clay over dull yellowish brown fine sand soil over brownish black clay over brownish black sand clay loam over dull yellowish brown fine sand soil. Water table is estimated at >180 cm. Floods reported to be common in April/ May. The soil texture is heavy at the first two horizons but lighter down the profile. Cracks observed on the surface to 50 cm soil depth, medium acid, Ustic moisture regime, stratification, Isohyperthermic STR  | Fluvisols  | Eutric Fluvisols    | Gleyi - Eutric Fluvisol |
| MB - P2 | Ochric A, Cambic B  | Almost flat, deep to moderately deep, well drained, greyish brown clay over brown to dark brown sand clay loam over greyish yellow fine sand over dull yellowish orange clay over brown sand clay soil. Water table was estimated at >180 cm. Floods in the area occurs in March/April or November /December. Vertical cracks were observed from the surface to 50 cm soil depth. Animal burrows (crotovinas) were observed in the profile from 0 - 74 cm depth. medium acid, Ustic moisture regime, stratification, Isohyperthermic STR | Fluvisols  | Eutric Fluvisols    | nd                      |
|         |                     |  | Cambisols  | Vertic Cambisols    |                         |
| MB - P3 | Ochric A            | Almost flat, deep to moderately deep, moderately well drained to well drained, dull yellowish brown clay over dull yellowish brown sand clay loam over greyish yellow brown sand clay loam over dull yellowish brown gravel sand soil. Water table was estimated at >150 cm. No previous history of floods in the area was reported. No cracks were observed. Animal burrows were observed from 0 - 126 cm depth, strongly acid, Ustic moisture regime, stratification, Isohyperthermic STR  | Fluvisols  | Eutric Fluvisols    | nd                      |

513 STR = Soil temperature regime, MB = Mbogo

514

515

516 **4 Conclusions and Recommendations**

517

518 In conclusion, the soils in the study area were classified as Eutric Fluvisols and Vertic Cambisols in FAO  
519 Soil classification Systems. The soil was deep to moderately deep, well to moderately well drained on flat  
520 or almost flat to undulating topography. The soil is stratified, developed under isohyperthermic soil  
521 temperature and ustic soil moisture regimes. Typically, the soil was dominantly brownish black in colour  
522 and clay texture top soils with gray to dull yellowish brown with overall stratification of fine sand, clay, sand  
523 clay and sand clay loam sub soils. Physically, bulk density was slightly above adequate but not restrictive in  
524 67 % of the studied profiles and ideal Bd in 33 % of the profiles. Chemically, soil reaction shows strongly to  
525 medium acid in the top soils but decreasing to medium to slightly acid with depth. Soil organic carbon was  
526 in good quality and all values were above the critical limits. N, P and K were very low, low or medium to  
527 very low. Ca and Mg were high to very high in the top soils but decreased with depth. CEC was high to very  
528 high and the soils were rated as non sodic with low to medium exchangeable Na<sup>+</sup>. Ca/Mg ratio showed that  
529 67 % were outside the critical range, Mg/K ratio was within the critical range in 33 % of the studied pedons.  
530 The top soils indicate that K deficiency is unlikely as K/TEB was greater than two (> 2) which is a generally  
531 accepted critical value. Taken together, the observed strongly to medium acid in the top soils in the study  
532 area calls for soil reaction management to acceptable levels through liming by using either Calcitic or  
533 Dolomitic lime whichever is relevant, available and affordable. Likewise, the low to very low NPK observed  
534 in the studied area suggests that such mineral elements should be incorporated in the fertilizer  
535 management program during and or after the irrigation development in the area.

536

537 **COMPETING INTERESTS DISCLAIMER:**

538 Authors have declared that no competing interests exist. The products used for this research are  
539 commonly and predominantly use products in our area of research and country. There is absolutely no  
540 conflict of interest between the authors and producers of the products because we do not intend to use  
541 these products as an avenue for any litigation but for the advancement of knowledge. Also, the research  
542 was supported by the National Irrigation Commission through Morogoro Zonal Irrigation Office in  
543 Tanzania and the Expanded Rice Production Program (ERPP).

544 **Conflict of interest statement**

545

546 Authors declare that they have no conflict of interest.

547

548

549 **4** **References**

- 550
- 551 [1] United Republic of Tanzania (URT). Accelerating Pro-Poor Growth in the Context of Kilimo  
552 Kwanza. URT, Dar es Salaam, Tanzania. 2009; 13-23.
- 553 [2] NBS. National Bureau of Statistics, Ministry of Finance Dar es Salaam and Office of Chief  
554 Government Statistician President's Office, Finance, Economy and Development Planning  
555 Zanzibar. , March, 2013; 264
- 556 [3] USAID. Maize Value Chain Assessment. 63 - slide Power Point Presentation by Technoserve, Dar  
557 es Salaam to NAFKA, 'Feed the Future'. 2010.
- 558 [4] Pauw K, and Thurlow J. "Agricultural Growth, Poverty, and Nutrition in Tanzania." Food Policy  
559 2011; 36 (6): 795–804.
- 560 [5] FAO. FAOSTAT. 2014; Online statistical database (retrieved November 2014) (available at  
561 <http://faostat.fao.org>).
- 562 [6] Homann-Kee Tui S, Bandason E, Maute F, Nkomboni D, Mpofo N, Tanganyika J, van 730 Rooyen  
563 A, Gondwe T, Dias P, Ncube S, Moyo S, Hendricks S, Nisrane F. Optimizing 734 Livelihood and  
564 Environmental Benefits from Crop Residues in Smallholder Crop-Livestock Systems 732 in  
565 Southern Africa. ICRISAT, Socio - economics Discussion Paper Series. 2013; 11. ICRISAT,  
566 Patancheru. 733 ICRISAT, 2010. ICRISAT Eastern and Southern Africa, 2009 Highlights. Nairobi.
- 567 [7] IITA. International Institute of Tropical Agriculture. 2009. Maize. Online: available at  
568 <http://www.iita.org/maize>. Accessed in August 29, 2014.
- 569 [8] Amani HKR. Agricultural Development and Food Security in Sub - Saharan Africa: 2004; Tanzania  
570 Country Report. [http://www.fao.org/tc/tca/work05/ Tanzania.pdf](http://www.fao.org/tc/tca/work05/Tanzania.pdf) (Accessed on 10 August, 2012)
- 571 [9] Kenneth GC, Achim D, Daniel TW, and Haishun Y. Meeting Cereal Demand while Protecting  
572 Natural Resources And Improving Environmental Quality. Annu. Rev. Environ. Resour. 2003;  
573 28:315–58 doi: 10.1146/annurev.energy.28.040202.122858
- 574 [10] Godfray HCJ, Beddington JR, Crute IR, Haddad L, Lawrence D, Muir JF, Pretty J, Robinson S,  
575 Thomas SM, and Toulmin C. Food Security: The challenge of feeding 9 billion people. Science.  
576 2010; 327:812–818.
- 577 [11] Foley JA, Ramankutty N, Brauman KA, Cassidy ES, Gerber JS, Johnston M, Mueller ND,  
578 O'Connell C, Ray DK, West PC, Balzer C, Bennett EM, Carpenter SR, Hill J, Monfreda C, Polasky  
579 S, Rockström J, Sheehan J, Siebert S, Tilman D & Zaks DPM. Solutions for a cultivated planet.  
580 Nature. 2011; 478: 337-342
- 581 [12] Match Maker Associates (2010). Value Chain Analysis of Rice and Maize in Selected Districts in  
582 Tanzania: A research report submitted to Agricultural Council of Tanzania. 24-89pp.
- 583 [13] Kibanda NJM. Rice research and production in Tanzania. [Internet]. African Japanese Plenary  
584 Workshop on Sustainable Rice Production, Alexandria, Egypt. 2008; Available from:  
585 <http://www.jesty.edu.eg>. (Accessed 20 May, 2009).
- 586 [14] NBS (National Bureau of Statistics). Results of the 2002-03 National Agricultural Sample Census:  
587 Volume II. National Bureau of Statistics, Ministry of Agriculture and Food Security, Ministry of  
588 Cooperatives and Marketing, and Ministry of Livestock. Dar es Salaam. 2007.
- 589 [15] Pascal AP. Tanzania's fertilizer assessment. A paper presented at the International Training  
590 Program on Fertilizer Policy and Marketing Strategies in Africa at Naura Springs Hotel, Arusha,  
591 Tanzania May 20 - 24th, 2013.
- 592 [16] Henao J, and Baanante CA. Nutrient depletion in the agricultural soils of Africa. International Food  
593 Policy Research Initiative, 2020 Vision, Brief No. 62, 1999; Muscle Shoals, Alabama.



- 594 [17] Bojo J, and Cassels D. Land Degradation and Rehabilitation in Ethiopia: **Area assessment**. AFTES  
595 Working paper No. 17. World Bank, Washington DC, 1995.
- 596 [18] Wu J, Norvell WA, Hopkins DG, Smith DB, Ulmer MG, Welch RM. Improved prediction and  
597 mapping of soil copper by kriging with auxiliary data for **cation - exchange** capacity. Soil Sci. Soc.  
598 Am. J. 2003; 67: 919–927
- 599 [19] Charpentier H, Doumbia S, Coulibaly Z, and Zana O. Stabilizing agriculture in northern and central  
600 Cote d'Ivoire: what are the new farming systems? Agriculture-et-Développement. 1999; 21: 4-70.
- 601 [20] Stoorvogel JJ, and Smaling EMA. Assessment of soil nutrient depletion in sub - Saharan Africa  
602 1983-2000. Report 28. The Winand Staring Centre for integrated land, soil and water research  
603 (SC-DLO), Wageningen.1990.
- 604 [21] Waddington SR, Murwira HK, Kumwenda JDT, Hikwa D, and Tagwira F. Soil fertility research for  
605 smallholder **maize - based** systems in Malawi and Zimbabwe. Harare: SoilFertNet/CIMMYT, 1998;  
606 312p.
- 607 [22] Ukut AN, Akpan US, Udoh BT. Characterization and classification of soils in steep sided hills and  
608 sharp-crested ridges of Akwa Ibom State, Nigeria. **Net J. Agric. Sci.** 2014; 2(2): 50-57
- 609 [23] Soil Survey Division Staff. Soil survey manual. United States Department of Agriculture; 1993.
- 610 [24] Bennett JG. An analysis of soil fertility systems in the hills of Nepal. A Joint project between Lumle  
611 Agriculture Research Center and the Natural Resources Institute, Kent, UK. 1995; Pp. 87.
- 612 [25] Sharu MB, Yakubu M, Noma SS, Tsafe AI. Characterization and classification of soils on an  
613 agricultural landscape in Dingyadi District, Sokoto State, Nigeria. **Nigerian J. Basic and Appl. Sci.**  
614 2013; 21(2):137-147.
- 615 [26] Karuma AN, Gachene K, Charles K, Msanya BM, Mtakwa PW, Amuri N, Gicheru PT. Soil  
616 morphology, physicochemical properties and classification of typical soils of Mwala District, Kenya.  
617 **Inter. J. Plant & Soil Sci.** 2015; 4(2):156-170.
- 618 [27] Kebeney SJ, Msanya BM, Ng'etich WK, Semoka JM, Serrem CK. Pedological characterization of  
619 some typical soils of Busia County, Western Kenya: Soil morphology, physico - chemical  
620 properties, classification and fertility trends. **Inter. J. Plant & Soil Sci.** 2015; 4(1):29-44
- 621 [28] FAO. FAO - UNESCO soil map of the world, revised legend, 1988; Technical Paper 20
- 622 [29] Chao DJ. Brief explanation of the geology of Kidatu sheet No. QDS 217. 1983.
- 623 [30] Akpan-Idiok AU, and Ogbaji PO. Characterization and Classification of Onwu River Floodplain  
624 Soils in Cross River State, Nigeria. **Inter. J. Agric. Res.** 2013; 8:107-122.
- 625 [31] Uwitonze P, Msanya BM, Mtakwa PW, Uwingabire S, and Sirikare S. Pedological Characterization  
626 of Soils Developed from Volcanic Parent Materials of Northern Province of Rwanda. Agric. Forestry  
627 and Fisheries. 2016; 5 (6): 225-236.
- 628 [32] Aubertin GM and Kardos LT. Root growth through porous media under controlled conditions. 1.  
629 Effect of pore size and rigidity. Soil Sci. Soc. Am. Proc. 1965; 29:290–293.
- 630 [33] Brady NC, and Weil RR. The Nature and Properties of Soils, 2008; 12th edition. Prentice-Hall, Inc.,  
631 Upper
- 632 [34] Lal R, and Shukla MJ. Principles of Soil Physics. Marcel Dekker, Inc. USA. 2005; 682p.
- 633 [35] Obasi SN, Onweremadu EU, Egbuche CT, and Iwuanyanwu UP. Characterization and  
634 Classification of Selected Rice Soils of Tropical Rainforest Region, Southeastern Nigeria. Agric.  
635 Forestry and Fisheries. 2015; 4 (3): 46-50.
- 636 [36] Landon JR. Booker Tropical Soil Manual: **A handbook** for soil survey and agricultural land  
637 evaluation in the tropics and subtropics. John Wiley and Sons: New York. 1991; 450p.



- 638 [37] USDA-NRCS. Soil bulk density/moisture/aeration. Soil quality kit-Education for educators.  
639 Accessed in October, Saddle River, NJ, USA. 2016; 881 p.
- 640 [38] NSS. Laboratory procedures for routine soil analysis. 3rd edition. Ministry of Agriculture and  
641 Livestock Development, National Soil Service (NSS), ARI Mlingano, Tanga, Tanzania. 1990.
- 642 [39] Brady NC, and Weil RR. The Nature and Properties of Soils. (13th edition). Macmillan Publishing  
643 Co., Upper Saddle River, New Jersey. 2002; pp.905.
- 644 [40] McKenzie NJ, Jacquier DJ, Isbell RF, and Brown KL. Australian Soils and Landscapes: An  
645 Illustrated Compendium. CSIRO Publishing: Collingwood, Victoria. 2004; pp.387.
- 646 [41] Khan ZH, Hussain MS, and Ottner F. Morphogenesis of Three Surface - Water Gley Soils from the  
647 Meghna Floodplain of Bangladesh. *Dhaka University J. Biol. Sci.* 2012; 21 (2): 17-27.
- 648 [42] Massawe BHJ. Digital Soil Mapping and GIS based Land Evaluation for Rice Suitability in  
649 Kilombero Valley, Tanzania. PhD thesis. Graduate School of Ohio State University. 2015.
- 650 [43] Prasad R and Power JF. Soil Fertility Management for Sustainable Agriculture. Boca Raton, FL:  
651 CRC Press LLC. 1997; 356p
- 652 [44] Adamchuk VI and Mulliken J. Precision agriculture site-specific of soil pH (FAQ), extension paper  
653 713, historical materials from University of Nebraska-Lincoln. 2005; EC05-705.
- 654 [45] EUROCONSULT. Agriculture Compendium for Rural Development in the Tropics and Sub-Tropics.  
655 3rd Revised edition. Elsevier, Amsterdam. Oxford. New York. Tokyo. 1989; 740pp.
- 656 [46] Msanya BM, Kimaro DN, Kimbi GG, and Munisi ALM. 2001. Land resources inventory and  
657 suitability assessment for the production of the major crops in the eastern part of Morogoro Rural  
658 District, Tanzania. Department of Soil Science, 3: 1 - 69.
- 659 [47] Walkley A, Black CA (1934). Acid extractable Zn in soil in relation to the occurrence of Zn  
660 deficiency symptoms of corn: A method of analysis. *Soil Sci. Soc. Am. Proc.* 12: 143-148.
- 661 [48] Loveland P and Webb J. Is there a critical level of organic matter in the agricultural soils of  
662 temperate regions: a review. *Soil & Tillage Res.* 2003; 70: 1 - 18.
- 663 [49] Makoi JHJR and Ndakidemi PA. Selected chemical properties of soil in the traditional irrigation  
664 schemes of the Mbulu district, Tanzania. *Afr. J. Agric. Res.* 2008; 3 (5): 348 – 356
- 665 [50] Vermeer JG, and Berendse F. The relationship between nutrient availability, shoot biomass and  
666 species richness in grassland and wetland communities. *Vegetation.* 1983; 53: 121-126.
- 667 [51] Tilman D. Plant dominance along an experimental nutrient gradient. *Ecol.* 1984; 65:1445 – 1453.
- 668 [52] Facelli JM and Pickett STA. Plant litter: its dynamics and effects on plant community structure. *Bot.*  
669 *Rev.* 1991; 5:1 – 32.
- 670 [53] Anderson JM. Carbon dioxide evolution from two temperate deciduous woodland soils. *J. Appl.*  
671 *Ecol.* 1973; 10: 361 – 378.
- 672 [54] Marx ES, Hart J, Stevens RG. Soil test interpretation guide. Oregon State University Extension  
673 Services. Oregon State University, Corvallis. 1996; 7pp
- 674 [55] Smiciklas KD, Mullen RE, Carlson RE, Knapp AD. Drought-induced stress effect on soybean  
675 seed calcium and quality. *Crop Sci.* 1989; 29: 1519-1523.
- 676 [56] Schwartz HF, Pastor-Corrales MA, eds. Bean production problems in the tropics, 2nd edn. Centro  
677 Internacional de Agricultura Tropical (CIAT), Cali, Colombia. 1989; 726 p.
- 678 [57] Lebron I, Suarez DL, Schaap MG. Soil pore size and geometry as a result of aggregate-size  
679 distribution and chemical composition. *Soil Sci.* 2002;167: 165 - 172.
- 680 [58] Peverill KI, Sparrow LA, and Reuter DJ. 'Soil Analysis. An Interpretation Manual.' (CSIRO  
681 Publishing: Collingwood.) 1999.

- 682 [59] Msanya BM, Kimaro DN, and Magoggo JP. Characteristics of two pedons and their implication for  
683 environmental management in parts of Mbinga District, Tanzania. Paper presented at the first  
684 annual Faculty of Agriculture Research Conference. August 28-30th. Sokoine University of  
685 Agriculture, Morogoro, Tanzania. 1995.
- 686 [60] Magoggo JP, Msanya BM, and Kimaro DN. Environmental profile for agricultural production and  
687 development of conservation strategies in Mahenge Village, Mbinga District, Tanzania. Miombo  
688 Woodland Research. 1996.
- 689 [61] Meliyo JL. Pedological investigation and characterization in Litembo village, Mbinga District,  
690 Tanzania. MSc (Agric). Dissertation, Sokoine University of Agriculture, Morogoro, Tanzania. 1997;  
691 113p.
- 692 [62] Msanya BM, Kaaya AK, Araki S, Otsuka H. and Nyadzi, GI. Pedological characteristics, general  
693 fertility and classification of some benchmark soils of Morogoro District, Tanzania. Tanzania. Afr. J.  
694 Sci. and Techn. (AJST) Science and Engineering Series. 2003; 4 (2): 101-112.
- 695 [63] Makoi JHJR, Mmbaga H. Soil Fertility Characterization in Mvumi and Mbogo - Komtonga Irrigation  
696 Schemes in Kilosa and Mvomero Districts, Morogoro Region, Tanzania, 2018.