

Original Research Article

Relative abundance and status of arthropod pests and diseases in honey bees at Ari district of southwestern Ethiopia

ABSTRACT

Formatted: Tab stops: 2.47", Left

Aims: To determine the species composition, relative abundance and status of honey bee pests and diseases in Ari district of Inka zone of southern western Ethiopia.

Study design: Four major honey producing peasant associations (PAs) representing diverse ecological zones were selected purposively, while the sub-localities, villages and beekeepers were selected randomly using a nested design

Place and Duration of Study: Survey was conducted between 1, august – 30, December 2019 with the interval of one month in major honey producing peasant associations representing diverse ecological zones of Ari district of south western Ethiopia.

Methodology: The assessment on abundance and status was made from of half kilogram of fresh and dry honey comb samples were taken from 720 farmer's hives of four peasant associations using taxonomic key of books related to arthropod pests and diseases of honey bee.

Results: Five species of predators and parasitoids belonging to four orders were recorded. Of these 76.63% accounts for small hive beetles, followed ants (38.31%) and wasps (19.92%). Besides, six type's diseases of honey bee caused by different pathogens were recorded. Of which, Chalk brood, other virosis, European foulbrood and American foulbrood, respectively were the most frequently occurring. In terms of the status, the diseases of honey bee were in the following orders, i.e., Chalk brood > other virosis > European foulbrood > American foulbrood > Sac Brood Virus, while their pests were in the following orders, i.e., Small hive beetles > Ants > Wasps > Wax moths > Spiders.

Conclusion: Arthropod pests and diseases were the most important constraints of honey colony and honey production in the study areas. Thus, it is important to design and implement effective management strategies to reduce loss of honey colony and honey production in the study area.

Keywords: Honey bee, pests, diseases, beekeeping, status, hives, abundance

1. INTRODUCTION

Beekeeping is an appropriate and well accepted farming technology that has been suited to extensive range of tropical Africa. It is important in diverse way including in production of honey, bee wax, queen, bee colonies and other products such as pollen, royal jelly, bee venom and propolis in cosmetics and medicines. Additional role of beekeeping is pollination of food crops and many plant species used for conservation of the natural environment and can be integrated the other activities like crop production, animal husbandry and horticulture crops [1, 2]. Apiculture is a deep rooted and long standing activity in Ethiopia where honey bee colonies are kept in traditional hives in backyard and in forest using different management practices. It accounts 1.3% of agricultural GDP, and currently one out of 10 rural households engaged in it and its contribution is substantial to rural income generation [3, 4].

However, beekeeping and honeybee colony establishment has been declining extremely in contemporary beekeeping in Ethiopia because of different abiotic and biotic factors. Among biotic constraints, pests and diseases are reported to be fundamental factors that are responsible toward significant decline of honey bee colony and beekeeping [5].

Consequently, there is a need for undertaking extensive research in such area (pests and diseases of honey bee) that affect honeybee health [6, 7]. Hence, conducting consistent (routine) surveillances and monitoring as well as research regarding the diseases and pests honey bee and attempting to fill this gap accordingly is the key steps to protect honeybee colonies, honey production and beekeeping from harmful diseases and pests [8]. Besides, survey and research

on pests and disease of honey bee might also provide baseline information that could help designing their effective preventive and management strategies that could easily adopted by beekeepers. Furthermore, though some research works associated with honey bee colony health (pests and diseases of honey bee) was conducted in different parts of the country, there is still lack of relevant and documented information regarding the relative abundance, species composition and the status of honey bee pests and diseases in the study area, in particular and Ethiopia, in general.

Therefore, to at least address some of the aforementioned gaps the current survey is designed with the objective of determining the species composition, relative abundance and status of honey bee pests and diseases in Ari district of Inka zone of south western Ethiopia.

2. MATERIAL AND METHODS

2.1 The Study Area Descriptions

The study was conducted in South Ari Woreda or district of South Omo Zone of Southwest Ethiopia. The woreda is located 678 km away from Addis Ababa (the capital city of Ethiopia), and its area is 1521 km², 547 km away from Hawassa (the capital city of SNNPR state), 17 km away from Jinka (the capital city of South Omo Zone). The woreda is sub-divided in to 46 kebeles and four urban centers, Gazer (the capital city of Ari woreda), Metser, Tolta and Wubhamer. South Ari is bordered on the south by Benatsemy, on the west by Mago river, on the north by Basketo (special woreda), on the northeast by Gamogofa zone and on the east by Male. (Figure 1). The study was conducted from 1, August to 30, December, 2019.




Figure 1 Map of the study site

3.2 The Study Design and Sampling Procedure

3.2.1 The Study Design

Out of 46 kebeles (smallest administrative unit of Ethiopia) in the district, four (4) kebeles or peasant associations (PAs) representing diverse ecological zones (Kure, Komer, Gedir and Sheshir) were selected purposively with the assistance of the Ministry of Agriculture (MOA) sub-kebele staff, while the sub-localities, villages and beekeepers were selected randomly using a nested design as adopted by earlier researchers [9, 10].

3.2.2 Sampling Procedure

From each peasant associations (PAs), about three sub-localities were randomly selected and from each sub-locality, three villages were selected at random with the assistance of the Ministry of Agriculture (MOA) sub-kebele staff as adopted by earlier researchers [9, 10]. Kebeles were selected purposefully based on abundance of honey production, such that those kebeles growing honey largely were selected for the survey, while villages, representative farmers and their hives were selected randomly.

3.3 Determination of the Species Composition, Relative Abundance and Status of Insect Pests

Out of 36 randomly selected villages in the aforementioned manner, a total of 720 farmers apiaries were selected randomly (two from each village) and 0.5 kg of fresh and dry honey comb samples were taken and the presence of pests and diseases in apiaries were tested, identified as well as their pictures were taken. In the selected apiaries, inspection of honeybee colonies and sampling of randomly picked life worker bees and bee brood (egg, larva and pupa both live and decayed) were also carried out for identification diseases and pests of honey bee under laboratory conditions using morphometric as indicted by earlier scientists [11, 12, 3, 14, 15]. Samples of pests and parasites that were found in or around hives were collected for laboratory conformation. Samples taken by brushing the bees off the comb through large mounted funnel or directly in universal bottle, killed, preserved in 70% methyl alcohol and labeled accordingly.

Each sample at each sampling date from each farmer's apiaries at each villages of the sampling site was collected in sampling bag or jar, labeled with necessary information and kept for further identification of insect pests and diseases. The samples of pests, workers and honey combs were brought to Zoology laboratory Jinka University for identification. The brood examination was conducted by random opening of brood cells. The brood removed from the cell with a fine forceps and the cells were inspected for the presence of pests.

The samples from different farmers' apiaries of villages of the same peasant association were sub sampled further after thoroughly mixing them to come up with a standard of 10 g (0.1kg) sample. Then, the procedures and taxonomic keys of the books related with honey bee arthropod pests and diseases by different scientists (Shimanuki et al 1990; Shimanuki and Knox 1991; Ritter and Akwatanakul, 2006; Sammataro and Yoder; 2011; FAO, 2018) were used to refer for identification purpose through assistance of trained Entomologist of Arba Minch University. Besides, keys and pictures from on line available literatures were also used for identification. Then after, insect pests and diseases were sorted according to their orders, families and species, and counted for each subsample honey from each of different farmer's traditional apiaries from each village, in each case noting the number.

Then after, the species composition, abundance and status of pests and diseases were determined using the formulas indicated below as adopted by previous researcher [16]. The average of the sum total of species of arthropods (insect pests) and diseases collected from sub-samples was used to determine the species composition, abundance and status of pests and diseases. For assessing pest's infestations, the main variables have been included abundance, relative abundance and Constance (frequency) of species found in samples as suggested by earlier researcher [16]. Abundance refers to the total number individuals of a species divided by the total number of samples (in this case the total number of kilograms) and it is expressed by the following formula:

Total number of individuals of species

Abundance of species = _____

Total number of samples

The relative abundance of species is expressed by the percentage of individuals of the species in a total number of observed individuals as shown in the following formula:

Number of individuals of a species

Relative Abundance of species = _____ x 100

Total number of observed individuals

Constance (frequency) expresses the percentage of species occurrence. It is obtained by the relationship between the number of samples containing the species and the total number of samples. The following formula is expressing this relationship.

Number of samples in which the species occurred

Constance of species = _____ x 100

Total number of sample

2.4 Data management and analysis

All the data collected from field survey were managed and analyzed using Microsoft Excel version 2013 and Statistical Program for Social Sciences (SPSS) version 16. Descriptive statistics (mean and percentage) were used for compiling and computing data collected on relative abundance, species composition and the status of honey bee pests and disease. Significant differences between means were separated by Tukey's honestly significant difference (THSD) test at 95% confidence interval level.

3. RESULTS AND DISCUSSION

3.1 The Relative Abundance, Species Composition and the Status of Honeybee Pests in Ari District

Table 1 and 2 demonstrated different categories of insect pests (predators or parasitoids) honey bee that were recorded from honey bee samples collected from different apiaries of Ari district.

Accordingly, five species of natural enemies (predators and parasitoids) belonging to four orders with in five family were recorded. Of these 76.63% accounts for small hive beetles from order Coleopteran with in the family Nitidulidae, followed by ants (38.31%) and wasps (19.92%) from order Hymenoptera with in two families Formicidae and Vespidae, respectively. However, the remaining 11.87 % and 8.81% were the greater wax moths and spiders of family Pyralidae and Theridiida of orders Lepidoptera and Araneae, respectively. Similarly, pervious researcher [17] reported that as different insect pests from different orders such as Coleopteran (including *Aethina tumida* of family Nitidulidae), Lepidoptera (including *Galleria mellonella* and *Achroea grisella* of family Pyralidae) and Hymenoptera (including *Dorylus* spp of family Formicidae and *Vespa* species of family Vespidae) seriously affects honey bee colony and honey production.

Of the aforementioned five species of predators and parasitoids of honey bee recorded, small hive beetles, followed by ants were the most abundant as they account for about 27.78 and 13.89 mean numbers of individuals, respectively per 10 g of honey sample. They were also the most frequently occurring as they occurred in 94.44% and 83.33%, respectively per 10 g of honey sample. Following these species, wasps and wax moths, respectively were the next abundant and frequently prevalent. However, spiders were the least abundant and less frequently occurring (Tables 1 and 2).

Generally, in terms of abundance, relative abundance, frequency of occurrence and status (economic importance), the pests of honey bee recorded from honey samples collected from Ari district were found to be in the following orders, i.e., Small hive beetles > Ants > Wasps > Wax moths > Spiders (Table 1). These natural enemies cause significant loss in apiculture by destroying combs, stores, hive and taking foraging of worker as they leave hive and by being true parasitoids by raising their offspring in bodies of bees [17]. The various pests honey bee recorded from Ari district in the current study is in agreement with the report of [18], in which ants, beetle, moths, lice, termites, mites, and large vertebrate animals such as amphibians, reptile, lizards, birds, mammals like honey badgers and mice were recognized as pests of honey bees in Ethiopia. In accordance to various arthropod pests in the current study area earlier scientist [8] also reported that as different types ants, wax moth (greater and lesser wax moths), mice, birds (different types), honey badger, wasps, death's head hawk moth, bee lice (*braula coeca*), beetles (different types), lizards, toads/frog, prey-mantis, spiders, *Pseudo scorpions* (chelifer species) were registered as the major honeybee pests locally in Ethiopia. Similarly, previous researcher [19] also reported as different types honey pests such as honey badger, spider, bee-eating birds, bee lice, beetles, wasps, Death Head hawks, moth mice and lizards were recognized in Bale zone of southeastern Ethiopia.

Furthermore, the categorization and determination of their economic status (importance) in the present study is in agreement with the works or reports of previous scientists [11, 12, 3, 14, 15], in which all the identified species honey bee pests were indicated as they significantly affect honey bee colony and honey production in different part of the world either indirectly by causing loss of honey (feed of the brood) or directly being parasitoid or predators of honey bee.

Moreover, in accordance with currently recorded pests of honey bee, it was reported that both adults and larvae small hive beetle were serious pests that weakened honey bee colonies or honey supers and as they multiply in huge numbers as well as their larvae tunnel through comb to eat brood, ruin stored honey, and ultimately destroy infested colonies. Ants were also reported to be most troublesome to honey bees and bee keeping sector and as they eats or carries off any comb contents, honey, pollen and brood [18]. Among ants, *Dorylus fulvus* was reported to be one of important honey bees' enemies that caused a serious problem [20]. Wax moths also reported to causes significance damage in colony of honeybees in several African countries [18].

Table 1. The species composition of honeybee pests recorded in Ari district

Pests of honey bee	Number predators or parasitoids	Abundance (average no. insects/ 10g of honey sample)	Relative abundance (%)	Frequency (% of samples containing each species)	Status
Small hive beetles	1000	27.78A	76.63A	94.44A	Major
Wasps	260	7.22C	19.92C	77.78C	Major
Wax moths	155	4.3D	11.87D	55.56D	Major
Ants	500	13.89B	38.31B	83.33B	Major
Spiders	115	3.19E	8.81E	50E	Major
Total	2030				

Means followed by the different letters within columns (upper case letters) are significantly different, $p < 0.05\%$ using Turkey's studentized range test (HSD).

Table 2 Taxonomic position of honey bee pests recorded in Ari district

Common name of pests of honey bee	Scientific name	Family	Order	Type of the pest
Small hive beetles	<i>Aethina tumida</i>	Nitidulidae	Coleoptera	Predator
Oriental hornets; Wasps	<i>Vespa orientalis</i>	Vespidae	Hymenoptera	Parasitoid
Greater wax moths	<i>Galleria mellonella</i>	Pyralidae	Lepidoptera	Predator
Driver ants (safari ants)	<i>Dorylus species</i>	Formicidae	Hymenoptera	Predator
Black widow; Spiders	<i>Latrodectus mactan</i>	Theridiida	Araneae	Predator

3.2 Prevalence honeybee disease in Ari district

Table 3 demonstrated different types of diseases and the associated pathogens of honey bee that were recorded from honey bee samples collected from different apiaries of Ari district.

Accordingly, six types diseases caused by different species of pathogens (five identified and one non identified pathogens) belonging to three types pathogenic agents were recorded. Among these diseases caused by different pathogens, Chalk brood, other Virosis, European foulbrood and American foulbrood, respectively were the most frequently occurring as they appeared 83.33, 66.67, 58.33 and 55.56 percent's, respectively per 10 g of honey sample collected Ari district. But, Stone brood and Sac Brood Virus (SBV) were less frequently appearing as they occurred 41.67 % and 27.78%, respectively per 10 g of honey sample collected.

Generally, in terms of frequency of occurrence and the status (economic importance), the diseases of honey be recorded from honey samples collected from Ari district were found to be in the following orders, i.e., Chalk brood > Other virosis > European foulbrood > American foulbrood > Stone Brood > Sac Brood Virus (SBV) (Table 3). The different diseases and their causative agents of honey bee recorded from Ari district in the present study is partially in accordance with the report of previous researcher [18], in which the Honey bee (*Aphis mellonifera* L.) is indicated to be prone to infection by fungal, bacterial and protozoan pathogenic organisms and as honey bee diseases in Ethiopia include Chalkbrood diseases that is caused by pathogenic fungi, *Ascospaera aphidis*, Nosematosis that is caused by *Nosema apis* and amoeba that is caused by a single protozoa *Malpighamoeba mellificae*. Besides, the categorization and determination of

their frequency occurrence and relative importance in Ari district the current study is in accordance with the works or reports of previous scientists [11, 12, 3, 14, 15], in which all the identified species honey bee diseases and their causative agents were shown cause significant loss of honey bee colony and honey production in different part of the world either indirectly by causing loss of honey (feed of the brood) or directly by affecting honey bee colony through being Entomopathogens of the different stages of honey bee.

Table 3. Prevalence honeybee diseases recorded in Ari district

Diseases of honey bee	Disease agents	Type	Number of samples infested	Frequency (% of samples containing each species)	Status
European foulbrood	<i>Melissococcus pluton</i>	Bacterial	21	58.33	Major
Chalk brood	<i>Ascosphera apis</i>	Fungal	30	83.33	Major
American foulbrood	<i>Paenibacillus larvae</i>	Bacterial	20	55.56	Major
Stone brood	<i>Aspergillus flavus</i>	Fungal	15	41.67	Major
Sac brood Virus (SBV)	<i>Virus Picorna-like</i>	Viral	10	27.78	Inter
Other Virosis	Not distinguished	Viral	24	66.67	Major

Inter =Intermediate

5. CONCLUSION

From the present survey five species of natural enemies (predators and parasitoids) belonging to four orders with in five families were recorded. Of these 76.63% accounts for small hive beetles from order Coleopteran with in the family Nitidulidae, followed by ants and wasps. Besides, six types diseases caused by different species of pathogens (five identified and one non identified pathogens) belonging to three types pathogenic agents were recorded. Among these diseases caused by different pathogens, Chalk brood, other Virosis, European foulbrood and American foulbrood, respectively were the most frequently occurring.

Generally, in terms of the status (economic importance), the diseases of honey bee recorded from honey samples collected from Ari district were found to be in the following orders, i.e., Chalk brood > Other virosis, > European foulbrood > American foulbrood > American foulbrood > Sac Brood Virus (SBV), while the pests of honey bee recorded were found to be in the following orders, i.e., Small hive beetles > Ants > Wasps > Wax moths > Spiders.

Accordingly along with other constraints, arthropod pests and diseases were found to be the most economically important constraints of honey colony establishment and honey production by **bee keepers** in the study areas of Jinka zones. Thus, it is important to design and implement safe and effective management strategies to reduce loss of honey colony and honey production, and hence income as well as food insecurity of poor farmers in the study area. Besides, provision of training to farmers and extension workers on safe handling of apiculture as well as management practices of arthropod pests and diseases of honey bee in the study area is urgently needed by any concerned bodies.

REFERENCES

1. Gidey Y, Mekonen T. Participatory technology and constraints assessment to improve the livelihood of beekeepers in Tigray Region, northern Ethiopia. *Momona Eth J Sci.* 2010; 2(1):76-92.
2. Mekonnen E. Characterization of honey production and marketing systems, challenges and opportunities In Ada Berga District, West Shoa Zone, Oromia, Ethiopia. A MSc thesis submitted to College of Agriculture and Environmental Sciences Post Graduate Program, Bahir Dar University, Ethiopia; 2016.

3. Aynalem AT, Mekuriaw ZZ. Study on the beekeeping situation, the level of beekeepers knowledge concerning local honeybee subspecies, their productive characteristics, and behavior in Eastern Amhara Region, Ethiopia. *Advances in Agriculture*; 2014.
4. Sebsib A, Yibrah T. Beekeeping practice, opportunities, marketing and challenges in Ethiopia: Review. *J Dairy & Veter sci*. 2018; 5(3):1-21.
5. Cham DT. Diversity of honey bee *Apis mellifera* subspecies (Hymenoptera: Apidae) and their associated arthropod pests in Cameroon, Doctoral dissertation, University of Nairobi; 2017.
6. Genersch E. Honey bee pathology: current threat to honeybee and beekeeping. *App Micro-bio Biotech*. 2010; 87:87-97.
7. Gemedi DA. Status of Honeybee *Apis mellifera* Bandansii Pests and Pathogens in Seka Chokersa District of Jimma Zone, Ethiopia. *Inter J Res Stud in Biosci*. 2017; 5 (12): 12-27. DOI: <http://dx.doi.org/10.20431/2349-0365.0512003>
8. Begna D. Honeybee diseases and Pests research progress in Ethiopia: A review. *Afri J Inse*. 2015; 3(1):093-6.
9. Emanu G. Studies on the distribution and control of Angoumois grain moth. *Sitotroga cerealella* in Sidama administrative region. M.Sc. Thesis, Alemaya University of agriculture, Alemaya, Ethiopia; 1993. pp. 10.
10. Hiruy B, Emanu G. Insect pests associated to stored maize and their bio rational 248 management options in Sub Sahara Africa. *Int J Acad Res and Dev*. 2018; 3(1):741-748.
11. Shimanuki H, Knox DA, Furgala B, Caron DM, Williams JL. Diseases and pests of honey bees. *Bee Keeping in the United States*; Science and Education Administration, United States Department of Agriculture: Washington, DC, USA; 1990; 335:118-28.
12. Shimanuki H, Knox DA. Diagnosis of honey bee diseases. *Diagnosis of honey bee diseases, Agriculture Handbook Number 690*; 1991. pp. 63.
13. Ritter W, Akwatanakul P. Honey bee diseases and pests: a practical guide. *FAO agricultural and food engineering technical report*; 2006. pp.42
14. Sammataro D, Yoder JA. Honey bee colony health: challenges and sustainable solutions. *CRC Press of Taylor & Francis Group*; 2011. pp.315.
15. FAO. Main bee diseases: Good beekeeping practices. *Food and Agriculture Organization of the United Nations Rome*; 2018. pp. 52.
16. Bueno, VHP, Souza MB. Ocorrência e Diversidade de Insetos Predadores e Parasitóides na Cultura de Couve Brassica oleracea Var. acephala em Lavras MG, Brazil; 1991. Latin.
17. Sarwar M. Insect pests of honey bees and choosing of the right management strategic plan. *Inter J Entom Res*. 2016; 1(2):16-22.
18. Tesfay H. Honey bee diseases, pest and their economic importance in Ethiopia. *Inter J Innov Scient Res*. 2014; 10:527-535.
19. Tesfaye B, Dadi G, Gelgelu T. Assessment of honeybee enemies (pests and predators) in Bale zone, southeastern Ethiopia. *J Agri Ext Rur Dev*. 2017; 9(4):53-61.
20. Desalegn B. Honey bee pest and predators of Ethiopia. 3rd proceedings of national conference of Ethiopia beekeeping association, Addis Ababa, Ethiopia; 2001. pp. 59-67.