

**NURSING DIAGNOSTICS IN THE CHAGAS
CARDIOMYOPATHY CARRIER: INTEGRATIVE
LITERATURE REVIEW**

ABSTRACT

Chagas' cardiomyopathy is a complication of Chagas' disease that presents in the cardiac form, which involves the manifestation of a clinical picture of dilated cardiomyopathy, with global left ventricular dysfunction and heart failure syndrome. In this context, the aim is to describe the Nursing Diagnosis evidenced in the literature in patients with Chagasic Cardiomyopathy. This is a descriptive and exploratory study of the Integrative Literature Review (ILR), with a qualitative approach. We searched the LILACS and PubMed databases for articles published without time limit, in Portuguese, English and Spanish, with the descriptors: Chagasic Cardiomyopathy; Nursing Diagnosis; Nursing Care. Data were analyzed by tabulation. Three articles were sought for the sample. The following Nursing diagnoses were highlighted: Decreased Cardiac Output; Intolerance to Activity; Poor Knowledge; Sleep and Rest impaired; Anxiety; Excessive fluid volume; Risk of impaired skin integrity; Comfort impaired. It is concluded that the diagnoses are directly related to heart failure caused by ventricular damage, and that interventions are essential for the improvement in the quality of life of patients. It is also noteworthy that the applicability of the Systematization of Nursing Care in these patients only improves the quality of service and has repercussions on patient safety, being also a way to promote the health of patients.

Keywords: Chagas Cardiomyopathy; Nursing diagnoses; Nursing care

1. INTRODUCTION

The Chagas disease (CD) is a pathology of infectious origin, in which it presents in the acute or chronic phases, and is still considered by the World Health Organization (WHO) as a neglected disease. It is related to low income and education, and in endemic countries it has a high morbidity and mortality rate, including in Brazil, with focal expression in unequal epidemiological contexts. The spatial distribution of the disease is primarily concentrated in the American continent due to the presence of more than 140 species of the insect vector (Triatominae, Hemiptera, Reduviidae), also considered as "American trypanosomiasis". However, CD has been presented in non-endemic countries, caused by the displacement of infected individuals and by other transmission mechanisms, as a result of the intense process of international migration in this decade [1].

In relation to epidemiology, the occurrence of cases and outbreaks by oral transmission, home vector without colonization and extradomiciliary vector, especially in the Legal Amazon. From 2007 to 2016, confirmed cases of acute Chagas disease were reported in most Brazilian states, representing an annual average of 200 cases. The largest distribution,

33 about 95%, is concentrated in the Northern region, with the state of Pará being responsible
34 for 85% of the cases. Of the main probable forms of transmission in the country, 69% were
35 by oral transmission, 9% by vector transmission, and in 21% the form of transmission was
36 not identified [2].

37

38 Transmission can occur in a variety of ways, including contact with the faeces and/or urine of
39 hematophagous triatomines, ingestion of food contaminated with parasites from infected
40 triatomines, maternal-fetal transmission, blood transfusion or organ transplantation, and
41 laboratory accidents. After the contamination occurs an incubation period of 1 to 2 weeks (in
42 oral transmission this period varies from 3 to 22 days), begins the acute phase, with an
43 average of 8 to 12 weeks. It is characterized clinically by prolonged febrile syndrome related
44 to high parasitemia, with few symptoms or asymptomatic or oligosymptomatic. In addition, it
45 may cause cardiac and central nervous system involvement, but severe forms of acute
46 disease evolve in less than 1% of patients [3].

47

48 In the chronic phase the disease presents with low parasitemia, and are still classified in
49 indeterminate form, that without clinical complications or symptoms, and the determined
50 forms, which have cardiac or digestive affections or both, or neurological which is very rare.
51 It is also noteworthy that 10 to 30% of carriers evolve to the determined forms, relating to the
52 decrease in quality of life and morbidity and mortality [4].

53

54 In this context, it stands out for the forms determined with emphasis on cardiac
55 complications. Chagas' cardiomyopathy involves the manifestation of a clinical picture of
56 dilated cardiomyopathy, with global left ventricular dysfunction and heart failure syndrome.
57 The Latin American Guidelines for the Diagnosis and Treatment of Chagas' Cardiopathy
58 established a categorization for left ventricular dysfunction, which reflects the gradation of
59 the evolutionary stages of heart failure followed in international guidelines for this syndrome.
60 Thus, the chronic phase of CD with cardiomyopathy can be classified into 5 evolutionary
61 stages (A, B1, B2, C and D) of left ventricular dysfunction. Sudden death and the progress of
62 heart failure (HF) are the most common mechanisms that cause death. The most
63 pronounced prognostic aspects are symptoms of advanced HF, cardiomegaly, LV systolic
64 dysfunction and non-sustained ventricular tachycardia, characterizing a severe condition and
65 presenting high mortality [5].

66

67 The patient needs specialized outpatient follow-up to minimize and control the complications
68 that can be modified, being assisted by a multiprofessional team, with the objective of
69 monitoring, treatment and monitoring the progression, in decompensated cases the patient
70 must be hospitalized. In this perspective, the professional Nurse stands out, who works
71 based on Nursing Theories, through the Systematization of Nursing Assistance, which is an
72 instrument for nursing care to be applied through scientific evidence, and which traces
73 Nursing Diagnosis (ND) for each problem evidenced based on the Nursing Theories. The
74 ND are performed with the help of the North American Nursing Diagnosis Association"
75 (NANDA), from the ND tracings the nurse prescribes interventions for each one, with the
76 objective of promoting health, preventing injuries, assisting in treatment and rehabilitation [6].

77

78 Thus, Chagas' cardiomyopathy is a serious condition that affects the quality of life of the
79 patient, with high chances of evolving to death. Thus, the following research question
80 emerged: what are the Nursing Diagnoses evidenced in the literature in patients with
81 Chagasic Cardiopathy. Then, the objective is to describe the Nursing Diagnostics evidenced
82 in the literature in patients with Chagasic Cardiopathy.

83

84

85 **2. MATERIAL AND METHODS**

86

87 This research is an integrative literature review (ILR), descriptive and exploratory, with a
88 qualitative approach. For Soares et al[7] the RIL is a form of research that allows reviewing,
89 criticizing and synthesizing the representative literature on a topic or subject in an integrated
90 manner and capable of generating new approaches and perspectives on the revised subject.
91 This method should be followed in stages: 1 - Preparation of the research question, 2 -
92 Establishment of criteria for inclusion and exclusion of studies/sampling or literature search,
93 3 - Definition of the information to be extracted from selected studies/ categorization of
94 studies, 4 - Evaluation of studies included in the integrative review, 5 - Interpretation of
95 results, 6 - presentation of the review/ synthesis of knowledge [8].

96

97 From the elaboration of the research question: which are the Nursing Diagnoses evidenced
98 in the literature in patients with Chagasic Cardiopathy, the databases used for the research
99 were listed. It was sought in Latin American and Caribbean Literature on Health Sciences
100 (LILACS) and PubMed. It includes original articles, case reports, theses, dissertations and
101 review, with no time limit, in English, Portuguese and Spanish. Incomplete research and
102 abstracts were excluded.

103

104 The search was performed using the descriptors: Chagas' cardiomyopathy; Nursing
105 diagnoses; Nursing care; and crossing the searches with the Boolean operator AND. For the
106 collection and organization of data, it was opted for the tabulation, elaborated by the authors
107 containing the following information: article number, authors, title, databases, year,
108 methodology and Nursing Diagnosis highlighted in the studies.

109

110 For the data analysis, in the first step after the selection of the sample, the data in the table
111 was organized and deepened in the reading, with the objective of highlighting the ND
112 reported in each study, and includes in the tabulation, being a simple descriptive analysis,
113 which from the organization of the data in the chart was possible to show the ND evidenced
114 in the studies, and thus selected for discussion.

115

116

117 **3. RESULTS AND DISCUSSION**

118

119 From the search with the selected descriptors, the LILACS database resulted in 9 articles,
120 and after reading only 3 studies were included for the sample. In PubMed, the search
121 showed 39 articles, but after reading the abstracts it was not possible to include any for the
122 sample, because they did not answer the research question. Thus, three articles were
123 included in the sample of this review. See below the table for the information extracted.

124

125 **Chart 1 - Information collected from the sample and organized in tabulation,**
126 **performed the searches in August. Belém-PA. Brazil, 2019.**

127

Number, Title year/Base	Authors	Methodology	Nursing diagnoses evidenced in the studies
1 – Nursing diagnoses in hospitalized patients due to chronic chagasic cardiopathy 1990	CRUZ, DALM. ARCURI, EAM.	A study was conducted with 30 patients admitted to three hospitals in São Paulo, from the interview, physical examination and medical records.	Decreased cardiac output. Intolerance to Activity. Deficient knowledge. Sleep and Rest impaired.

LILACS			
2- Nursing care for clients with Chagasic cardiomyopathy: a case study 2010 LILACS	Moreira NS, Almeida GN, Oliveira JG.	This was a qualitative research of descriptive nature of the type case study.	Decreased cardiac output; Anxiety.
3- Nursing diagnoses of the nanda of a patient with chagasic cardiomyopathy: a case study 2018 LILACS	SOARES et al.	The study was carried out in a public hospital, in the city of Recife-PE, where the medical history and physical examination of F.M.D., male, 42 years old, was performed and lives in Afogados da Ingazeira.	Excessive liquid volume; Risk of impaired skin integrity. Comfort impaired.

128 **Source:** Authors' research, 2019.

129

130 Regarding the profile of the articles, all used as methodology the Systematization of Nursing
 131 Care to highlight the ND in the patient with Chagasic Cardiomyopathy, 1 study conducted
 132 the research with 30 patients and 2 with only one patient, being a case study, but the 3
 133 articles highlighted the ND. The year of publication was very discrepant between one and
 134 the other, such as the 1st of 1990, 2nd of 2010 and 3rd of 2018. It was also shown the
 135 limitation to this research, because even with the inclusion criterion for study published
 136 without time limits, it was only possible to include 3 studies in the sample, showing the lack
 137 of these studies in relation to the theme with focus on ND.

138

139 From the organization in the chart, it was possible to highlight the ND evidenced in the
 140 studies that were: Reduced Cardiac Output; Intolerance to Activity; Deficient knowledge;
 141 Sleep and Rest impaired; Anxiety; Volume of excessive liquid; Risk of impaired skin integrity;
 142 Harmful Comfort.

143

144 Chagas' cardiomyopathy is a cardiac complication of CD considered severe and that its
 145 repercussions cause an impact on the quality of life and mortality of patients. Complications
 146 can cause: Arrhythmias, Cardiac Insufficiency. Thus, the main ND are related to the
 147 symptoms of the complications that cardiomyopathy causes.

148

149 The first ND evidenced in this review was decreased cardiac output, related to low
 150 ventricular ejection, related to left ventricular dilation. Cardiac insufficiency is a severe
 151 complication of CD, because due to left ventricular dysfunction, the ventricle pumps less
 152 blood than it should, thus emerging the main symptoms of HF, such as dyspnea, limb edema
 153 and fatigue. For the interventions in this ND, it is based on: evaluating vital signs, observing
 154 signs of hypoxia, low perfusion, cardiogenic shock, oxygen saturation (O₂), fluid balance,
 155 fluid restriction, adherence to pharmacological treatment [9].

156

157 Another ND evidenced in this study was the Excessive Liquid Volume, characterized by
 158 dyspnea and edema. Thus, this ND is also directly related to decreased cardiac output, and
 159 requires interventions in conjunction with this ND, which are based on: lower limb elevation,

160 use of diuretics, hydric restriction, avoidance of physical effort, weight control, hydric balance
161 and guidance in relation to adequate pharmacological therapy [10].
162

163 It also showed that ND Activity Intolerance was reported in patients, characterized by
164 limitation of activities, related to low ventricular ejection. In patients with low ventricular
165 ejection, the performance of daily activities is limited, because the basic daily activities
166 demand the use of oxygen by the cells, especially the myocardium, so in low ejection the
167 oxygen demand is reduced and causing tiredness and fatigue in patients. For the
168 interventions, the same measures of decreased cardiac output are basically performed,
169 because the main cause is related to this ND.
170

171 The Harmful Comfort was also highlighted in these patients, characterized by limitations of
172 daily activities related to Cardiac Failure. Thus, the symptoms again affect the quality of life
173 of the patient, interfering even in comfort. The interventions should be: oriented as to their
174 limitations, guide everything about the pathology, guide as to pharmacological therapy, seek
175 positions that improve comfort, guide on leisure practices that cause no effort, promote
176 support, hope and encouragement [11].
177

178 One of the articles in the sample also showed the risk of impaired skin integrity, related to
179 cardiovascular disease edema. Thus, edema in this condition is caused by decreased
180 venous return due to low ejection fraction and low venous pressure. Thus, the skin in this
181 condition presents greater fragility, because the elasticity is impaired and causes a decrease
182 in thickness, increasing the risk of injury by pressure and trauma. For interventions, light
183 walking is recommended to stimulate venous return, elevation of the lower limbs, adherence
184 to pharmacological therapy, fluid restriction, skin hydration, comfort massages [12].
185

186 Sleep and impaired rest were also reported in the studies, which are characterized by
187 interruptions in sleep, related to dyspnea, frequent diuresis. Symptoms such as nocturnal
188 dyspnea, related to dorsal position, cause discomfort for sleep and rest of the carrier, and
189 pharmacological therapy with the use of diuretics stimulates a higher frequency of diuresis,
190 and is also a factor to interfere in the appropriate sleep pattern. Guidance should be given
191 when the diuretic intake during the morning, because the effects during the night will be
192 reduced, as well as the low water intake only during the night, with the aim of reducing
193 urinary output during sleep. In relation to dyspnea in dorsal decubitus, it is recommended the
194 elevation of 30% degrees of the chest, can be performed with the use of two pillows,
195 because this position improves the thoracic expansion and consequently minimizing the
196 respiratory discomfort [13].
197

198 The ND Deficient Knowledge and Anxiety were also described, they are ND related,
199 because the lack of knowledge is linked to anxiety. However, the complications and
200 limitations that the pathology causes have repercussions on feelings of anxiety that can
201 cause other psychological conditions. Deficient knowledge, on the other hand, has
202 repercussions on the appropriate care that the patient should provide to minimize symptoms,
203 such as adherence to pharmacological therapy, hydric restriction, limitations to activities, etc.
204 for interventions, should guide the patient in relation to the pathology, such as complications,
205 limitations and basic daily care with health, adherence to treatment, explaining the harmful
206 effects of non-adherence. When the patient knows all the stages of his pathology, he tends
207 to influence treatment adherence and the implementation of preventive measures in order to
208 minimize more serious complications [14].
209

210 The applicability of Systematization of Nursing Care in patients with Chagas'
211 cardiomyopathy becomes essential for qualified care, so the nursing diagnoses provide

212 greater scientificity in nursing care, and the interventions discussed applied have a direct
213 impact on improving the quality of life of these patients.

214

215 It is also noteworthy that this study presented limitations in relation to the search for articles,
216 because it showed the low inclusion of articles that address the Systematization of Nursing
217 Assistance and Nursing Diagnosis in patients with Chagasic cardiomyopathy, emphasizing
218 that the lack of studies focused on this theme is very large, and that it was necessary to
219 include an article published in 1990, because it was the only one that conducted the
220 research that several patients, the others were case studies, so from this observation we
221 chose to include it in this review. Thus, it is emphasized that there is a need for research on
222 the subject, since the epidemiology shows that cases of Chagasic Cardiomyopathy increase
223 and are related to higher mortality and decreased quality of life of patients.

224

225

226

227 **4. CONCLUSION**

228

229 Through this study it was possible to describe the main Nursing Diagnoses presented by
230 patients with Chagasic Cardiomyopathy, but showed that the ND are specifically related to
231 cardiac complications, such as Cardiac Failure, which is the main complication caused by
232 damage to the ventricles, so it was also discussed about the main nursing interventions
233 applied to each ND.

234

235 Nursing care based on the Systematization of Nursing Care only ensures the quality of care
236 and service, impacting on patient safety, since it is a respective process and based on
237 Nursing Theories, making it a scientific and evidence-based care. Thus, the use of nursing
238 systematization should be emphasized for the applicability of care in patients with Chagasic
239 Cardiomyopathy.

240

241 It is hoped that this study will provide the literature with knowledge for nursing professionals
242 about the care of patients with Chagas' cardiomyopathy, thus offering evidence based on
243 reliable scientific bases, resulting in an improvement in the quality of service of these
244 patients, as well as a way to promote the health of patients, influencing the improvement of
245 the quality of life of these patients.

246

247 **COMPETING INTERESTS**

248

249 Authors have declared that no competing interests exist.

250

251

252 **CONSENT**

253

254 It is not applicable.

255

256 **ETHICAL APPROVAL**

257 It is not applicable.

258

259 **REFERENCES**

260

261 1. Dias JCP, Ramos AN, Luquetti, Gontijo ED, Alejandro, Torres RM, et al. II Brazilian
262 Consensus on Chagas Disease, 2015. Epidemiol Serv Saúde [Internet]. 2016 Jun [cited

263 2019 Aug 24]; 25 (21): 1–10. Available from:
264 http://www.iec.pa.gov.br/template_doi_ess.php?doi=10.5123/S1679-
265 [49742016000500007&scielo=S2237-96222016000500007](http://www.iec.pa.gov.br/template_doi_ess.php?doi=10.5123/S1679-49742016000500007&scielo=S2237-96222016000500007)

266

267 2. Brazil M da S. Epidemiological situation - Chagas disease [Internet]. Health from A to Z.
268 2017 [cited 2019 Aug 28]. Available from: [http://www.saude.gov.br/saude-de-a-z/doenca-de-](http://www.saude.gov.br/saude-de-a-z/doenca-de-chagas/situacao-epidemiologica)
269 [chagas/situacao-epidemiologica](http://www.saude.gov.br/saude-de-a-z/doenca-de-chagas/situacao-epidemiologica)

270

271 3. Brazil M da S. Clinical Protocol and Therapeutic Guidelines Chagas Disease [Internet]. 1st
272 ed. Brasília: Secretariat of Science, Technology and Strategic Inputs; 2018 [cited 2019 Aug
273 24]. 141 p. Available from: <http://conitec.gov.br>

274

275 4. Brazil M da S. Health Surveillance Guide [Internet]. 1st ed. Brasília: Secretariat of Health
276 Surveillance; 2017 [cited 2019 Aug 24]. 286 p. Available from: www.saude.gov.br/bvs

277

278 5. Simões MV, Romano MMD, Schmidt A, Martins KSM, Marin-Neto JA. Cardiomyopathy of
279 Chagas Disease. *Int J Cardiovasc Sci* [Internet]. 2018; 31 (2): 173-89. Available from:
280 http://www.scielo.br/pdf/ijcs/v31n2/en_2359-4802-ijcs-31-02-0173.pdf

281

282 6. John Wiley & Sons L. Nursing Diagnostics NANDA 2015-2017. 10th. Nursing Diagnoses -
283 Definitions and Classification 2015-2017. 2014. 497 p.

284

285 7. Soares C, Hoga L, Peduzzi M, Sangaleti C, Yonekura T, Silva D. Integrative review:
286 concepts and methods used in nursing descriptors. *Rev Esc Nursing USP*. 2014; 48 (2):
287 329–39.

288

289 8. Mendes KDS, CP Silveira RC, Galvão CM. Integrative review: research method for
290 incorporating evidence into health and nursing. *Text Context Enferm* [Internet]. 2008 Dec
291 [cited 2019 Jul 15]; 17 (4): 758–64. Available from:
292 http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0104-
293 [07072008000400018&lng=en&tIng=en](http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0104-07072008000400018&lng=en&tIng=en)

294

295 9. Sousa MM de, Araújo AA de, Freire MEM, Oliveira J dos S, Oliveira SH dos S. Nursing
296 diagnoses and interventions for the person with decompensated heart failure. *J Res Fundam*
297 *Care Online*. 2016; 4 (1): 2733–9.

298

299 10. Gonçalves LWP, Pompeo DA. Application of the outcome-present state test model in
300 patient with congestive heart failure. REME Rev Min Enferm [Internet]. 2016 [cited 2019 Aug
301 18]; 20 (0). Available from: <http://www.gnresearch.org/doi/10.5935/1415-2762.20160047>

302

303 11. Almeida DV de, Oliveira KF de, Oliveira JF de. Most frequent nursing diagnoses in
304 patients admitted to the Coronary Intensive Care Unit. Arq Med Hosp Fac Ciênc Med St
305 House Sao Paulo [Internet]. 2013; 58: 64–9. Available from:
306 <http://archivesmedical.fcmsantacasasp.edu.br/index.php/AMSCSP/article/download/225/255>

307

308 12. Ribeiro MAS, Lages JSS, Lopes MHBM. Nursing diagnoses related to skin: operational
309 definitions. Rev Lat Am Nursing [Internet]. 2012 Oct [cited 2019 Aug 29]; 20 (5): 863–72.
310 Available from: [http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0104-
311 11692012000500007&lng=en&tlng=en](http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0104-11692012000500007&lng=en&tlng=en)

312

313 13. Santos MA dos, Guedes E of S, Barbosa RL, Cruz D of ALM da Sleeping difficulties
314 reported by patients with heart failure. Rev Lat Am Nursing [Internet]. 2012 Aug [cited 2019
315 Aug 29]; 20 (4): 644–50. Available from:
316 [http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0104-
317 11692012000400003&lng=en&tlng=en](http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0104-11692012000400003&lng=en&tlng=en)

318

319 14. Lacerda MS, Cirelli MA, Barros ALBL de, Lopes J de L, Lacerda MS, Cirelli MA, et al.
320 Anxiety, stress and depression in family members of patients with heart failure. Rev from
321 USP's Esc Enferm [Internet]. 2017 [cited 2019 Aug 29]; 51 (0). Available from:
322 [http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0080-
323 62342017000100403&lng=en&tlng=en](http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0080-62342017000100403&lng=en&tlng=en)