

**SELF-BREAST EXAMINATION: KNOWLEDGE AND PRACTICE OF YOUNG FEMALE STUDENTS TOWARDS SELF-BREAST EXAMINATION AND BREAST CANCER**

**ABSTRACT**

Objective of the study is to assess the knowledge towards self-breast examination (SBE) of breasts among young adults in various districts of Sindh province in Pakistan. Descriptive cross sectional study design was used to conduct the study at Larkana. A total of 171 young female students participated in the study including students of matric, intermediate, undergraduate and postgraduate level. Participants were given a self-administered questionnaire designed to assess the knowledge of students towards SBE and Breast cancer. Data was analyzed using descriptive and inferential statistical tests in SPSS (Statistical Package for Social Sciences version 23.0). A total 171 young adults participated in the study. Only a 2.33% were married. Most of the participants (47.36%) had intermediate level education, 43.85% were undergraduate, 8.18% had a family member with breast cancer, 38.6% of the respondents didn't know that the breast cancer is most fatal cancer in women all over the world. A 43.8% considered that family history as a risk factor for the breast cancer, 61.4% considered that pain in the breast is symptom for the breast cancer, 83.63% of participants didn't know how to perform breast self-examination, 88.3% of the participants were not practicing self-breast examination. It was concluded that most of the young students in various districts of Sindh province in Pakistan do not have reasonable knowledge towards Self-breast examination and do not practice the same in routine life.

**Keywords:** *Breast cancer, Self-breast Examination, Female Students, Sindh Province*

## **1. INTRODUCTION**

Cancer is a leading cause of death worldwide in countries of all income levels [1]. The International Agency for Research on Cancer (IARC) estimates that in 2018, there would be 18.1 million new cases and 9.6 million deaths linked to cancer worldwide [2]. Breast cancer typically produces no symptoms when the tumor is small and most easily treated, which is why screening is important for early detection. The most common physical sign is a painless lump. Sometimes breast cancer spreads to underarm lymph nodes and causes a lump or swelling, even before the original breast tumor is large enough to be felt. Breast cancer is typically detected either during a screening examination, before symptoms have developed, or after a woman notices a lump. Most masses seen on a mammogram and most breast lumps turn out to be benign (not cancerous), do not grow uncontrollably or spread, and are not life-threatening. When cancer is suspected, microscopic analysis of breast tissue is necessary for a diagnosis and to determine the extent of spread (stage) and characterize the type of the disease [3]. Meanwhile incidence and mortality rates for most cancers (including lung, colorectum, female breast, and prostate) are decreasing in the United States and many other western countries, same are increasing in many developing countries as an outcome of the trends to adopt western unhealthy lifestyles of smoking, physical inactivity and consumption of calorie-dense food. In developing countries, the incidence of new cancers which was 56% reported I 2008, has been projected to 60% by year 2030 [4]. According to the American cancer Society estimated deaths due to female breast cancer cases were more the 40,000 in US in 2017 [3]. Historically, Asian breast cancer incidence has been lower than the Europe and North America but for recent decades incidence of the same seems significantly increasing in Asian countries. Statistically a significant rise in mortality rate was shown in less developed countries like Malaysia and Thailand [5]. Global Cancer Statistics 2018 revealed that the incidence of breast cancer was top second around the

globe after lung cancer. In 2018, total new cases of the lung cancer were 2,093,876 while the total new cases of breast cancer reported only around five thousand short of the aforementioned having a figure 2,088,84 causing 626,679 number of deaths. This suggests 11.6% of the newly reported cancers and 6.6% of deaths occurred in 2018 [6]. Breast cancer is the most commonly diagnosed cancer in women (24.2%, i.e. about one in 4 of all new cancer cases diagnosed in women worldwide are breast cancer), and the cancer is the most common in 154 of the 185 countries included in GLOBOCAN 2018. Breast cancer is also the leading cause of cancer death in women (15.0%), followed by lung cancer (13.8%) and colorectal cancer (9.5%), which are also the third and second most common types of cancer, respectively; cervical cancer ranks fourth for both incidence (6.6%) and mortality (7.5%) [7]. Cancer is an issue of higher magnitude in both developed and emerging countries, especially breast cancer. In Pakistan, breast cancer is one of the leading cancers. Lack of proper facilities and most importantly the awareness among the patients in Pakistan, there is progress and prevalence of breast cancer. Moreover there is plenty of data available on breast cancer worldwide to estimate survival of breast cancer patients but unluckily there is no such study in Pakistan. There are several reasons for that too, inability of reaching out to those cancer patients who doesn't seek medical treatment. Higher costs of cancer treatment and fewer hospitals especially designed for cancer treatment [8]. The incidence of breast cancer in Pakistan is highest in Asians after Jews in Israel and 2.5 times higher than that in neighboring countries like Iran and India, accounting for 34.6% of female cancers. The Pakistani population is deficient in information regarding breast cancer etiology and epidemiology [9]. Among Asian countries especially in Pakistani population there has been alarming increase in the incidence of breast cancer. Out of total 213,377 surgical specimens registered during the above mentioned period 53,012 specimens were breast cancers, according to the data published by Karachi Cancer Registry in 2000, for population of the Karachi South district during 1995-1997 [10]. The breast cancer is more common in middle age women of average age 48 years in Pakistan, whereas, in the western countries, the breast cancer is more common in older age

after 60 years. Other factors include overexposure to estrogen due to early periods of 12 years, late menopause after 55 years, late childbirth at 30 years or no childbirth, short time period of postpartum, breastfeeding for less than two years, previous breast benign lesion of one breast that have metastasized to another non-diseased breast and previous chest radiations mostly at the time of adolescence are at a higher risk, due to continuous growth of breast [11]. Performance of specific BSE components may reduce the risk of death from breast cancer. It is very important to early detect and start the treatment of breast cancer. Periodic screening mammography with or without clinical breast examination reduces rates of death from breast cancer for women aged 50 to 69 years. Breast self-examination (BSE), is self-administered and inexpensive to become aware of any visible changes in breast [12]. The purpose of breast self-examination is for a woman to learn the topography of her breast, know how her breasts normally feel and be able to identify changes in the breast should they occur in the future. In fact, whether breast self-examination alone can reduce the number of deaths from cancer is currently a source of controversy [13].

## **2. MATERIALS AND METHODS**

A descriptive cross-sectional study design was used to conduct study at Larkana Sindh. Data was collected from the young students belonging to various districts and studying at different levels of education (from matric to postgraduate level) in 2018. Young students from different districts of Sindh appeared in a co-curricular event organized by a non-profit organization IQRA ETO. A self-administrative questionnaire was developed including the questions regarding age, district, academic level, and family history of breast cancer, knowledge regarding breast cancer, knowledge and practice of breast self-examination. The questionnaire was revised and validated by panel of 6 experts medical and teaching field; Cronbach's alpha was used for internal consistency which was 0.80. Before collection of the data, a pilot study was done in a class room of matric and intermediate students where 22 students responded the questionnaire in presence of the class teacher and researchers. Some of


the students questioned the abbreviation of SBE which was later written in full form in the questionnaire for the ease of the students. Few students questioned the various names of diagnostic test like CT scan and MRI that they did not understand, which the researchers explained to the students during pilot study and later on while data collection. Students took less than 5 minutes to answer the questionnaire. The approval for data collection was obtained from IQRA ETO, a written approval was given by the “Director Events” of the aforementioned non-profit organization. Before serving the questionnaires, students were verbally explained about the importance and purpose of the study and a verbal consent was obtained. Students were told that their participation is voluntary and data will be confidential and be used for the sake of research only. Neither the name nor the ID number of any student was asked in the questionnaire.

All the students attending the event were the sampling population for the study, using a total population sampling technique all the students appearing in the event were given with the questionnaire and a cover letter explaining the importance of the study with consent form attached, 180 questionnaires were distributed and 174 questionnaires were returned showing high response rate of more than 97%, 3 questionnaires were discarded by the researchers for having incomplete data. The total sample size for the study was 171. The collected data was entered and analyzed using the SPSS (Statistical Package for Social Sciences version 23.0) and Microsoft Office Excel 2010. Descriptive and Inferential statistical tests were used. Data was analyzed using frequency, percentage.

### 3. RESULTS

There were 171 young adults who participated in the study. Most of them belonged to Larkana, Ghotki , Kashmore and Jacobabad districts. Age of the participants ranged from 15 to 28, where 73% of those were 20 years old or below (Figure 1).

Fig. 1. Age of the Participants




Out of 171 participants, only a 2.33% were married. Most of the participants (47.36%) had intermediate level education, 43.85% were undergraduate, 5.2% were matric level and rest of the participants were graduate and above (Table 1).

**Table 1. Marital Status and Academic Level of the Participants**

Marital Status and Academic Level of the Participants		
	No.	%
Academic Level of the Participants		
Matric	9	5.20%
Intermediate	81	47.36%
Undergraduate	75	43.85%
Graduate	3	1.75%
Postgraduate	3	1.75%
Marital Status of the Respondents		
Married	4	2.33%
Unmarried	167	97.66%

Responding to the question regarding risk factors of the breast cancer, 43.8% considered that family history is a risk factor for the breast cancer (Figure 2).


**Fig. 2. Knowledge of the participants towards risk factors of breast cancer**






While answering the question on symptoms of the breast cancer, 61.4% considered that pain in the breast is symptom for the breast cancer (Figure 3).

**Fig. 3. Knowledge of the participants towards symptoms of breast cancer**




More than half of the participants (90 respondents, 56.6%) considered the CBE (clinical breast examination) is the method to diagnose the breast cancer, while 34% considered the mammography as diagnostic tool (Figure 4).

**Fig. 4. Knowledge of the participants towards tools of diagnosis for breast cancer**



Answering the question regarding how often self-breast-examination should be performed, around 2/3<sup>rd</sup> of the participants confessed that they don't know at all, nearly 1/5<sup>th</sup> of the participants considered that self-breast examination should be practiced every month (Figure 5)

**Fig. 5. Frequency of self-breast examination performance**



Surprisingly, 38.6% of the respondents didn't know that the breast cancer in women all over the world. There were 14 respondents (8.18%) who had a family member with breast cancer; rest of the 91.8% had no family history of breast cancer. One of the surprising finding of the study was that the 83.63% of participants didn't know how to perform breast self-examination. The last question in the study was whether the respondent practices self-breast examination herself or not, the finding against this research question is never less than a call for attention; surprisingly 88.3% of the participants were not practicing self-breast examination (Table 2).

**Table 2. Knowledge and practice of participants towards SBE**

	<b>Knowledge and practice of participants towards SBE</b>			
	No.	Yes %	No.	NO %
Q.5 Do you know that breast cancer is the most common cancer in females?	105	61.40%	66	38.60%
Q. Do you have anyone in your family with breast cancer?	14	8.18%	157	91.81%
Q. Do you know how to perform self-Breast Examination?	28	16.37%	143	83.63%
Q. Do you practice Self-breast Examination (SBE)?	20	11.69%	151	88.30%

#### **4. DISCUSSION**

Breast cancer is one of the leading cancers in women with an increasing incidence worldwide. Early detection of breast cancer is highly important improve breast cancer outcome and survival. Performance of breast self-examination is one of the important steps for identifying breast disease at an early stage by the woman herself. Similar to our study, a study was conducted among young students in Ajman. Likewise our study, the frequent age range was 18-22 years and most of the participants (82%) were unmarried. It was found that shows the family history of breast cancer which was 9.2% [13]. In our study, participants were students and 73% of those were 20 years old or below. However our sample size was smaller compared to this study. For the family history, both studies had nearly similar findings as there were 8.18% students in our study who had family member with breast cancer.

In a study conducted in Malaysia among students of a private university where more than half of the participants (55.4%) mentioned that they practiced BSE. young female students have knowledge about breast self- examination, the majority mentioned that radio and TV were their main sources of information (38.2%), followed by family members (21.1%), friends (14.7%), and then newspapers (12.4%) [14]. In contradiction to this study we found that 83.63% our participants didn't know how to perform self-breast examination.

Nearly three quarter of the respondents (87.7%) had heard of BSE, but out of all those only 19.0% of them were performing SBE every month at Ahmadu Bello University. Regarding the sources of information about BSE among respondents, media was found to be most common followed by health workers accounting for 45.5% and 32.2% respectively [15]. A significant difference is obvious in knowledge and practice

of SBE among students in this study, yet again; findings of our study stand in contradiction with this as most of the students in our study lacked the basic knowledge and more than 4/5<sup>th</sup> participants in our study didn't know how to perform SBE.

We had nearly similar finding with a study conducted in Oman, where a vast majority (3/4<sup>th</sup>) of the participants were aware about the presenting symptoms of breast cancer; a total of 145 (92%) and 139 (88%) correctly recognized Breast pain and Lump in the breast as the most common symptoms of breast cancer [16]. Similarly in our study, answering the questions on symptoms of breast cancer 61.4% and 55.7% opted for pain the breast and lump in the breast respectively. Family history as risk factor breast cancer was another finding of our study similar to this study. The most widely known risk factor among the participants was family history of breast cancer (86.62 %) [16]. Likewise, choice of family history as risk factor for breast cancer was on top (43%) in our study.

Similar findings to our study was observed in high school students of Turkey where most common reason for not doing SBE was having “not knowing how to perform Self Breast Examination” among 98% participants [17].

Positive family history of breast cancer was reported by 156(13.2%) women; 420(35.5%) believed advancing age was a risk factor; 1041(87.9%) never had breast self-examination; 1106(93.4%) never had a clinical breast examination; and 1171(98.9%) never had screening mammogram [18]. Likewise; in our study 88.30% do not practice SBE and 83.63% do not know how to perform SBE.

More than 2/3<sup>rd</sup> participants had sound knowledge about the risk factors and signs of breast cancer. Majority was aware of the benefits of mammography. More than 80% had the consensus that breast cancer is curable if detected early and more than 50% thought that a surgeon should be consulted first if lump is palpable [19]. On contrary, findings of our study are almost opposite to this as more than 80% of our participants neither practiced SBE nor they knew how to perform it. Yet again, there is reasonable

justification for this as the respondents in our study were young adults most of them aged below 20 and belong to non-medical background as well, while the participants in mentioned study were practicing medical doctors, consultants, residents, medical officers and nurses. The other most important justification to this can be a social taboo in various districts of Sindh province to discuss and share regarding breast cancer.

## **5. CONCLUSION**

Our study shows that the young students in various districts of Sindh province in Pakistan do not have reasonable knowledge towards Self-breast examination and do not practice the same in routine life. It was also observed that despite having knowledge regarding SBE, female students do not practice Self-breast examination. Since early detection of the breast cancer like any other cancer is very important for effective treatment and management, having reasonable knowledge and starting to practice SBE for young females will be a great tool to prevent and manage breast cancers.

## **6. RECOMMENDATIONS**

Our study has observed that despite there was a reasonable deficiency in knowledge and practice of SBE among students yet there was a positive attitude of students which provide an available ground to initiate the mass media and awareness programs. When more than 80% of the students

Knowledge of BC and BSE practices are highly deficient in this study population. However, there is generally a positive attitude which provides a fertile ground for awareness dissemination in order to improve knowledge and practices. Awareness programs for breast cancer especially for young students should be designed all over the province emphasizing Self breast examination young females.

## **CONSENT**

It was taken verbally from all participants.

## **ETHICAL APPROVAL**

The SMBBMU Larkana Ethics Committee approval has been collected and preserved by the author.

## REFERENCES

1. Torre LA, Siegel RL, Ward EM, Jemal A. Global Cancer Incidence and Mortality Rates and Trends--An Update. *Cancer Epidemiol. Biomarkers Prev.* 2016;25(1):16-27. doi:10.1158/1055-9965.EPI-15-0578.
2. Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA: A Cancer Journal for Clinicians.* 2018;68(6):394-424.
3. ACS. *Breast Cancer Facts & Figures.* American Cancer Society. 2017.
4. Jemal A, Center MM, DeSantis C, Ward EM. Global patterns of cancer incidence and mortality rates and trends. *Cancer Epidemiology and Prevention Biomarkers.* 2010;19(8):1893-907
5. Bhoo-Pathy N, Yip CH, Hartman M, Uiterwaal CS, Devi BC, Peeters PH, Taib NA, van Gils CH, Verkooijen HM. Breast cancer research in Asia: adopt or adapt Western knowledge?. *European Journal of Cancer.* 2013;49(3):703-9.
6. World Health Organization. Cancer burden rise to 18.1 million new cases and 9.6 million cancer deaths in 2018. 2018.
7. Arshad S, ur Rehman M, Abid F, Yasir S, Qayyum M, Ashiq K, Tanveer S, Bajwa M, Ashiq S. Current situation of breast cancer in Pakistan with the available interventions. *Int. J. Biosci.* 2019;11(6):232-40.
8. Amirali Z, Edgar K, Azim ZA, Sadruddin S. Breast cancer in Pakistani females. *i-Manager's Journal on Nursing.* 2014;4(2):11.
9. Bhurgri Y, Bhurgri A, Hassan SH, Zaidi SH, Rahim A, Sankaranarayanan R, Parkin DM. Cancer incidence in Karachi, Pakistan: first results from Karachi cancer registry. *International Journal of Cancer.* 2000;85(3):325-9.



10. Al-Sharbatti SS, Shaikh RB, Mathew E, Al-Biate MA. Breast self examination practice and breast cancer risk perception among female university students in Ajman. *Asian Pac. J. Cancer Prev.* 2013;14(8):4919-23.
11. Harvey BJ, Miller AB, Baines CJ, Corey PN. Effect of breast self-examination techniques on the risk of death from breast cancer. *Canadian Medical Association Journal.* 1997;157(9):1205-12.
12. World Health Organization. Guidelines for the early detection and screening of breast cancer. 2006.
13. Asif HM, Sultana S, Akhtar N, Rehman JU, Rehman RU. Prevalence, risk factors and disease knowledge of breast cancer in Pakistan. *Asian Pacific Journal of Cancer Prevention.* 2014;15(11):4411-6.
14. Al-Naggar RA, Al-Naggar DH, Bobryshev YV, Chen R, Assabri A. Practice and barriers toward breast self-examination among young Malaysian women. *Asian Pac. J. Cancer Prev.* 2011;12(5):1173-8.
15. Gwarzo UM, Sabitu K, Idris SH. Knowledge and practice of breast self-examination among female undergraduate students. *Annals of African Medicine.* 2009;8(1).
16. Al Junaibi RM, Khan SA. Knowledge and Awareness of breast cancer among university female students in Muscat, Sultanate of Oman-A pilot study. *Journal of Applied Pharmaceutical Science.* 2011;1(10):146.
17. Karayurt Ö, Özmen D, Çetinkaya AÇ. Awareness of breast cancer risk factors and practice of breast self examination among high school students in Turkey. *BMC Public Health.* 2008;8(1):359.
18. Sarwar MZ, Shah SF, Yousaf MR, Ahmad QA, Khan SA. Knowledge, attitude and practices amongst the Pakistani females towards breast cancer screening programme. *Screening.* 2015;8:17-9.
19. Kumar S, Imam AM, Manzoor NF, Masood N. Knowledge, attitude and preventive practices for breast cancer among health care professionals at Aga Khan Hospital Karachi. *Journal of the Pakistan Medical Association.* 2009;59(7):474.