

Enhancing Quality of Fodder Sorghum through application of Fe chelate

Abstract

Sorghum (*Sorghum bicolor* (L.) Monech) is one of the important fodder crops for ruminants and it is a dual-purpose crop used as a food and fodder but one of the great limiting factors with the forage sorghum is that it is having an anti-nutritional factor cyanogen, which is extremely toxic to the animals feeding on them. The present experiment was conducted at Tamil Nadu Agricultural University, Coimbatore to study the effect of iron on the quality of fodder crop. The experiment was laid in Factorial CRD with 3 factors namely seven sources of Iron (Fe glycinate citrate, Fe tartrate, Fe glutamate, FeSO₄, Fe-EDDHA, Fe malate) four levels (FeSO₄ - 0, 25, 37.5 and 50 kg ha⁻¹ and Fe chelates - 0, 1, 2.5 and 5 kg ha⁻¹) and two different soils (calcareous and non-calcareous). It was seen that application of Ferrous citrate at 5 kg ha⁻¹ has shown the maximum reduction in cyanogen content and it has shown an increase in crude protein and decreased crude fibre content, which are desirable **qualities** in fodder crop. The variation in the above parameter may be due to the fact that Fe is a constructive component of different enzymes (hematian, cytochroms, propyrin and ferrichrome) that favourably improves the nutritional environment of crop and final yield. All treatments supplied with micronutrients irrespective of the soil has **shown decreased** cyanogen levels below the **threshold** of 200 ppm, which is safe for feeding cattle as green forage under normal cultivation.

Key words: Fe chelates, Sorghum, Cyanogenic glycoside, Crude protein, Crude fibre

Introduction

Micronutrients are important to plant growth and profitability. All the nutrient have their own importance but are used in different amounts and work collectively as activators of many plant functions. These micronutrients help the plants to complete their life cycle (Chaney *et al.*, 1992).

Iron is involved in many physiological processes of plants such as activation of many enzymes which are used in photosynthesis and respiration along with synthesis of chlorophyll. Iron is associated with heme and non-heme proteins such as ferredoxin, main constituent in the chloroplast where it participates in sulphur and nitrogen metabolism and has important role in energy transfer within plants. Deficiency results in chlorosis in young leaves that is shown by yellow and green strips along the length which is avoided by spraying iron complexes (Schmidt, 1999; Tagliavini & Rombola, 2001; Ashemad, 2001).

Sorghum is an important forage crop grown for green as well as dry forage production over wide areas. It is fast growing, palatable, nutritious and mainly utilized as green forage. Being an exhaustive crop, yield and quality of sorghum fodder suffers heavily if proper amount of fertilizer is not applied.

The hydrogen cyanide in sorghum is stored in the form of non-poisonous cyanogenic glycosides. When the leaf tissue is ruptured – such as when chopping for forage or chewing by animals- these glycosides come in contact with enzymes in other parts of the plant and/or saliva and get broken down into their constituent compounds: hydrogen cyanide and sugar causing the animals to suffocate leading to death.

In India, due to increased population pressure and competing demand for food crops, it is not possible to increase the area under fodder crops. The only way to bridge the large gap between supply and demand of fodder is to maximize the fodder production per unit area and unit time within the existing farming systems and utilizing marginal and sub marginal dry lands and problematic soils for developing feed and fodder resources. With this aim the study was conducted to see the effect of different iron sources on the quality of the forage crops and its effectiveness in controlling iron induced deficiency.

Materials and Methods

The experiment was conducted in a pot house of Department of Soil Science and Agricultural Chemistry, Tamil Nadu Agricultural University. The treatment **comprised of** seven sources of iron (Fe glycinate citrate, Fe tartarate, Fe glutamate, FeSO₄, Fe-EDDHA, Fe malate) four levels (FeSO₄ - 0, 25, 37.5 and 50 kg ha⁻¹ and Fe chelates - 0, 1, 2.5 and 5 kg ha⁻¹) and two different soils (Calcareous and Non-calcareous). The treatment application was made through soil application of FeSO₄ and Fe chelates. **Planting was done** during May and harvested during **August** to determine the fresh and dry forage yield. The dried samples were ground to determine forage nutritive value. Crude protein was determined by the micro-Kjeldahl method (Pearson, 1970), whereas crude fiber was determined according to the method of AOAC (1984).

The method used for estimation of HCN content **was as suggested by** (Hogg and Ahlgren, 1942). A quantity of 0.15 g of green plant material was cut into short pieces with scissors and dropped in a test tube and **3 to 4 drops of chloroform were** added. A strip of moist filter paper saturated with sodium picrate solution was suspended above the moist sample. **Then,** saturated filter paper was held in place with a cork stopper which **served** to seal the test tube and it was incubated at room temperature (20°C) for 12 to 24 hours. The sodium picrate which prevailed in the filter paper was reduced to a reddish compound in proportion to the amount of hydrocyanic acid which evolved from the sample. The colour produced was dissolved by placing the filter paper in a clean test tube containing 10 ml of distilled water after which the colour of the sample was matched with colour standards and concentration of HCN content was measured in spectrophotometer and then expressed in ppm. The collected data were analyzed according to the statistical procedure described by Gomez and Gomez (1984).

Results and Discussion

Cyanogenic Glycoside content

Hydrocyanic acid (HCN) is an antinutritional factor which is potentially toxic to the animal which feeds on 30–35-day-old sorghum crops (Wheeler *et al.* 1990). The cyanogenic glycoside, dhurrin found in sorghum can be hydrolysed in the rumen and liberate deadly hydrocyanic acid. Hydrocyanic acid content in excess of 200 ppm (on wet weight basis) in the forage sorghum is toxic to the animal health.

The application of metal-amino acid chelates significantly affect the forage glycoside content, yield and the quality parameters like crude protein and crude fibre compared with control (no fertilizer applied) and non-chelated and synthetic chelated micro-nutrients. In soil 1, the HCN content was reduced when compared to soil 2 (Table 1) and within the treatment application the one applied with ferrous citrate recorded the lowest content of HCN irrespective of the soil. Increase in yield was noted with amino acid chelates of micro-nutrients as compared with control.

The statistical analysis obtained for the three-way ANOVA on the effects of sources, different levels and different soils on cyanogenic glucoside production in sorghum leaves is given in Table 1 where it can be seen that sources, different levels and different soils have all significantly influenced cyanogenic production in sorghum leaves. This showed that each factor had an important influence on cyanogenic glucoside production in sorghum, confirming that different sources, soil and the levels of their application significantly influence the cyanogen content in leaves. The cyanogen content varied based on the different sources, its corresponding levels and also showed significant difference between soils. Here it is seen that there is a significant reduction in the levels of HCN as the levels of different treatments are increased and it is seen that the effect of treatments on the soil is also significant. Calcareous and non-calcareous soils showed high response to the treatment application and even though the latter was found to bring out much more drastic reduction when compared to the former. This shows that there is influence of nutrients and mainly micronutrients have a major role in the suppression of anti-nutritional factors.

Table 1. Effect of different sources of iron and soils on the cyanogenic glycoside content

Fe sources	Non-calcareous soil					Calcareous soil				
	N0	N1	N2	N3	Mean	N0	N1	N2	N3	Mean
Fe glycinate	97.35	67.47	67.47	32.32	66.15	102.96	70.80	69.31	35.23	69.57
Fe citrate	95.79	28.59	28.59	26.79	44.94	102.40	58.87	56.88	33.56	62.93
Fe tartarate	94.89	47.74	47.74	30.08	55.11	99.77	56.63	50.79	31.32	59.63
Fe glutamate	95.89	42.84	42.84	30.36	52.98	105.16	59.05	54.35	36.36	63.73
FeSO ₄	95.42	62.55	62.55	55.05	68.89	106.30	78.51	67.23	57.57	77.40
Fe-EDDHA	96.78	50.77	50.77	42.37	60.18	104.91	59.54	58.62	45.96	67.26
Fe malate	96.31	63.93	53.21	50.67	66.03	103.33	67.76	64.61	57.35	73.26
Mean	96.06	51.99	50.45	38.23	59.18	103.55	64.45	60.26	42.48	67.68
			SEd				CD (0.05)			
	Fe		0.243				0.488**			
	N		0.184				0.369**			
	Soil		0.130				0.261**			
	Fe *N		0.487				0.977**			
	Fe*Soil		0.344				0.690**			
	N*Soil		0.260				0.522**			
	Fe*N*Soil		0.689				1.381**			

Levels of fe sources - FeSO₄ – N0-0, N1-25, N2-37.5 and N3-50 kg ha⁻¹ and Fe chelates – N0-0, N1-1, N2-2.5 and N3-5 kg ha⁻¹

Table 2. The effect of different sources of iron and soils on the crude fibre content

Fe sources	Non-calcareous soil					Calcareous soil				
	N0	N1	N2	N3	Mean	N0	N1	N2	N3	Mean
Fe glycinate	28.703	26.02	25.67	24.65	26.26	28.73	25.57	25.26	23.81	28.73
Fe citrate	27.905	26.02	23.02	22.08	24.76	28.03	22.89	22.04	21.23	28.03
Fe tartarate	29.107	28.96	28.37	27.61	28.51	29.36	29.19	28.20	27.14	29.36
Fe glutamate	27.216	26.90	26.74	26.47	26.83	28.78	26.93	26.58	26.14	28.78
FeSO ₄	29.530	29.18	28.78	29.19	29.17	28.99	29.57	29.01	29.24	28.99
Fe-EDDHA	28.880	28.84	28.36	28.63	28.68	29.57	29.41	28.14	28.51	29.57
Fe malate	29.875	29.17	28.20	28.67	28.98	29.79	30.09	26.89	28.66	29.79
Mean	28.75	27.87	27.02	26.76	27.60	29.04	27.66	26.59	26.39	28.75
			SEd				CD (0.05)			
	Fe		0.136				0.274**			
	N		0.103				0.207**			
	Soil		0.073				0.146**			
	Fe *N		0.273				0.548**			
	Fe*Soil		0.19373				0.388**			
	N*Soil		0.146				0.293**			
	Fe*N*Soil		0.387				0.776**			


Levels of fe sources - FeSO₄ – N0-0, N1-25, N2-37.5 and N3-50 kg ha⁻¹ and Fe chelates – N0-0, N1-1, N2-2.5 and N3-5 kg ha⁻¹

Crude protein

The crude protein content in dry matter is of utmost importance in forage crop as it determines the palatability and digestibility of forage crops. The application of different amino acid chelates has shown significant differences in the crude protein content (Fig 1). The

maximum crude protein (12%) was recorded by the treatment supplied with ferrous citrate irrespective of soil and was statistically superior to all the other treatments. The protein contents **increased** by the application of amino acid chelated micronutrients. It was also seen that there was proportionate increase in the protein content with respect to the different levels applied. The higher protein contents in dry matter ultimately will result higher protein yield on unit area. The significant differences in crude protein contents in dry matter have also been confirmed by Nabi *et al.* (2006), Filho *et al.* (2004) and Tauqir *et al.* (2009). Improved protein contents might be due to better uptake of nutrients which is an integral part of protein synthesis.

Fig 1. Response of Fe sources and soil on crude protein content




Levels of Fe sources - FeSO₄ – N0-0, N1-25, N2-37.5 and N3-50 kg ha⁻¹ and Fe chelates – N0- 0, N1-1, N2-2.5 and N3-5 kg ha⁻¹

Crude fibre

The crude fibre in forage material has adverse effect on forage quality as it affects the digestibility. The data presented in Table 2 show that the highest value for crude fibre was recorded by Ferrous sulphate (29.18 %). The lowest value of crude fibre was seen in dry matter produced by Ferrous citrate and it could be taken as good quality forage (Fig 2). The significant differences among sorghum genotypes have already been confirmed by studies conducted by Mahmud *et al.* (2003) and Nabi *et al.* (2006).

Fig 2. Response of Fe sources and soil on crude fibre content (%)



Levels of Fe sources - FeSO₄ – N0-0, N1-25, N2-37.5 and N3-50 kg ha⁻¹ and Fe chelates – N0-0, N1-1, N2-2.5 and N3-5 kg ha⁻¹

Conclusion:

The outcome of the experiment suggested application of micronutrients irrespective of soil has a greater potential in improving the quality of forage crop. Under the light of present study, the application of Fe Citrate is found to have brought about lower cyanogen content, crude

fibre, higher crude protein content and yield. The result of the improvement in quality is due to the effect of Fe which is a constructive component of different enzymes (hematian, cytochroms, propyryn and ferrichrome) that favourably **improve** the nutritional environment of crop and final yield. Irrespective of **soil**, there was improvement in the quality of sorghum for forage purpose.

References

- AOAC 1965. Official methods of analysis. 10th ed. Association of official Agricultural Chemicals. Washington, DC, USA.
- Ashemad, S.D. 2001. The chemistry of ferrous bis glycine complexes. *Arch. Latinoam. Nutr.*, 51: 7-12.
- Chaney, R.L., B.A. Coulombe, P.R. Bell and J.S. Angle. 1992. Detailed method to screen dicot cultivars for resistance to Fe-chlorosis using Fe DTPA and bicarbonate in nutrient solutions. *J. Plant. Nutr.*, 15: 2063-2083.
- Filho, C. J. D. A., M. C. D. Monterior, F. I. D. Oliverira, I. N. Tabosa, G. Q. Bastos, and O. Y. D. Reis (2004). Dry matter yield and contents of crude protein in genotypes of forage sorghum. *Agropecuaria Tecnia*. 25: 59-63.
- Gomez, K.A. and A.A. Gomez, (1984). Statistical procedures for agricultural research (2 ed.). John wiley and sons, NewYork, 680p.
- Hogg PG, Ahlgren HC. A rapid method of determining hydrocyanic acid content of single plant of Sudan grass. *J. American Society of Agron*. 1942; 43:199-200.
- Mahmud, K., I. Ahmed, and M. Ayub (2003). Effect of nitrogen and phosphorus on fodder yield and quality of two sorghum cultivars (*Sorghum bicolor* L.). *Int. J. Agric. Biol*. 5:61-63.
- Nabi, C. G., M. Riaz, and G. Ahmed (2006). Comparison of some advanced lines of Sorghum

bicolor L. Monech for green fodder/dry matter yield and morpho economical parameters. J. Agric. Res. 44: 191-196.

Nur FO, Siti AH, Umi KY. Comparative evaluation of organic and inorganic fertilizers on total phenolic, total flavonoid, antioxidant activity and cyanogenic glycosides in cassava (*Manihot esculenta*). African J Biotechnol. 2013;12(18):2414–21. doi: 10.5897/AJB12.1248.

Persson, J. A.; Wennerholm, M.; O'Halloran, S. Handbook for Kjeldahl Digestion; FOSS, Hilleroed, Denmark, 2008.

Schmidt, W. 1999. Mechanisms and regulation of reduction-based iron uptake in plants. *New Phytol.*, 141: 1-26.

Tagliavini, M. and A.D. Rombola. 2001. Iron deficiency and chlorosis in orchard and vineyard ecosystems. *Eur. J. Agron.*, 15: 71-92.

Tauqir, N. A., M. Sarwar, M. A. Jabbar, and S. Mahmood (2009). Nutritive value of jumbo grass (*Sorghum bicolor* sorghum sudanefe) silage in lactating nili-ravi buffaloes. Pakistan Vet. J. 29: 5-10.

Wheeler, J.L, Mulcahy C, Walcott JJ, Rapp GG Factors affecting the hydrogen cyanide potential of forage sorghum. Australian Journal of Agricultural Research. 1990; 41:1093-1100. [doi:10.1071/AR9901093](https://doi.org/10.1071/AR9901093).