

COMPARATIVE ANALYSIS OF THE PHYTOCHEMICALS, PROXIMATE AND MINERAL COMPOSITIONS OF SCENT LEAF (*Ocimumgratissimum*) AND BITTER LEAF (*Vernoniaamygdalina*) LEAVES.

ABSTRACT

Ocimumgratissimum(fam.Lamiaceae) ‘Scent leaf’ and *Vernoniaamygdalina*(fam.Asteraceae) ‘Bitter leaf’ are commonly consumed vegetables in Southern Nigeria. *O. gratissimum* is used by the ‘Igbos’ of South East Nigeria to flavor soups and stews while it is popular with the ‘Yorubas’ of South West Nigeria for treatment of stomach disorders. *V. amygdalina* is a well known vegetable in West Africa used as both food and for its pharmaceutical properties. The enormous use of these vegetables in the diet motivated the present study whereby the food compositions, mineral constituents and the phytochemicals of the two vegetables were determined. Qualitative analysis, of the aqueous extracts, revealed the presence of saponins, terpenoids,tannins, flavonoids, steroids, alkaloids, phenolics and anthraquinones in both vegetables.Generally *V.amygdalina*contained higherquantities of secondary metabolites, saponin which was reported as the highest amount was $5.71\pm 0.12\text{mg/g}$ and $3.52\pm 0.01\text{mg/g}$ followed by terpenoids $5.64\pm 0.11\text{mg/g}$ and $3.40\pm 0.11\text{mg/g}$ in *V.amygdalina* and *O. gratissimum* respectively. There is however a minute amount of anthraquinone in the two vegetables. Proximate analysis, and macro elements were determined using their various standard methods. The results of proximate analysis revealed the presence of moisture ($12.28 \pm 0.02\%$ and $10.01\pm 0.01\%$) protein ($35.37\pm 0.11\%$ and $22.20\pm 0.02\%$) and total ash ($6.00\pm 0.20\%$ and $5.75\pm 0.10\%$) each in *V.amygdalina* and *O.gratissimum* respectively. The following minerals were present calcium, magnesium, potassium, sodium, iron, copper and zinc, with zinc being in the highest concentration ($46.80\pm 0.03 \text{ mg/kg}$ and $58.00\pm 0.12 \text{ mg/kg}$) followed by copper ($9.45\pm 0.11 \text{ mg/kg}$ and $16.60 \pm 0.03\text{mg/kg}$) each in *V.amygdalina* and *O.gratissimum* respectively. There are higher values of phytochemicals and food components in *V.amygdalina* while *O.gratissimum* had higher mineral contents. These two vegetables are potential source of components for complementary medicine.

Keywords: Plant, Food, Bitter leaf, Scent leaf, nutrients.

INTRODUCTION

Medicinal plants have been identified and known throughout human history as plants whose roots, stems, leaves and seeds possess therapeutic, tonic, and other pharmacological potentials. They are used as drug, food additives and for their nutritive values. Medicinal values of some plants lie in the chemical substances that produce definite physiological actions in their body. Among the chemical substances are phytochemicals which are bioactive natural compounds found in plants (1). Many plants that are used in traditional medicine to alleviate symptoms of illnesses have been found to possess phytochemicals, examples such plants include; *Azadirachta indica* (Dogonyaro), *Vernonia amygdalina* (Bitter leaf), *Allium sativum* (Garlic), [*Zingiber officinale* (Ginger), *Moringa oleifera* (Moringa), *Ocimum gratissimum* (Scent leaf) etc. Chemical compounds present in these plants mediate their effects on human body through processes identical to those already well understood from chemical compounds in conventional drugs. Thus herbal medications do not differ greatly from conventional drugs in terms of their effects on the body. This enables herbal medicines to be effective as well as having the potentials to cause harmful side effects (2, 3). The use of herbs and search for drugs and dietary supplements derived from plants have accelerated in recent years because medicinal plants are known to contain some chemical substances which can be used for treatment purposes or to produce drugs (4). Medicinal plants play vital roles in the health of individuals, in fact most modern drugs are derived from them (5,6). Medicinal-plant-based drugs have the advantage of being simple, effective and have broad spectrum activity. An ethno-botanical and ubiquitous plant serves as rich resources of natural drugs for research development (7). **With the crises of covid-19 pandemic, researchers throughout the world are combing the earth searching for vaccines, remedies and possible breakthrough with the use of conventional drugs medicinal plants.**

Knowledge of the chemical constituents of plants is necessary for the discovery of the therapeutic agents in plants and for the purpose of disclosing new sources of economic materials like essential oils, gums, precursors etc for the synthesis of complex chemical substances (8). Pharmacologists, natural products and environmental chemists, botanists, microbiologists, and other researchers are combing the earth looking for phytochemicals from plants that could be further developed for treatment of various diseases and for new leads to development. The revival of interest in the use and importance of medicinal plants by the WHO and other developing countries has led to intensified efforts on the documentation of ethno-medical data of medicinal plants as traditional herbalists keep no records, their knowledge is passed on verbally from one generation to the other leading to missing gaps in knowledge. All these intrigued the researchers to this topic as much information are needed on the phytochemicals, nutritional value and mineral constituents of medicinal plants hence the study focused on *vernonia amygdalina* and *ocimum gratissimum* two of the most commonly used vegetables in Southern Nigeria.

Ocimumgratissimum (Scent leaf) is a scented shrub with lime green fuzzy leaves which are used to flavor soup and spice food due to their strong aromatic odour (9). *Ocimumgratissimum* is grown for the essential oil in its leaves and stems. The essential oil possesses antibacterial properties (10), and is also used as an insect repellent so also the leaves when dry and burnt (10). It is used in the traditional medicine for treatment of diarrhea (11), stomach ache, as a febrifuge, and components of anti-malaria remedies (12), general tonic, and antiseptic in wound dressing, skin infection, conjunctivitis and bronchitis. An infusion of the leaves called "Ocimum tea" is dispensed as a remedy for fever and diaphoresis. The roots are used as sedatives for children. The phytochemical screening of *Ocimumgratissimum* (scent leaf) shows that it is rich in alkaloids, tannins, phytates, flavonoids and oligosaccharides (13).

Vernoniaamygdalina (Bitter leaf) is a small evergreen shrub that grows well in tropical Africa. It is called bitter leaf because of its bitter taste. The bitterness is as a result of the presence of the sesquiterpenes lactones and steroid glycosides (1). *Vernoniaamygdalina* is the most widely cultivated of the genus *vernonia* which has about 1000 species of the shrub (14), an ever green shrub with a characteristics odour and bitter taste. It is well known as a medicinal plant for diabetes. In Nigeria the leaves are used as vegetables for cooking the popular bitter leaf soup as it stimulates the digestive system.

MATERIALS AND METHODS

SAMPLE COLLECTION AND PREPARATION: Samples of *Ocimumgratissimum* and *Vernoniaamygdalina* leaves were collected from a garden behind Federal Government College Okigwe in Imo State Nigeria, identified and authenticated in the Department of Crop Science, Micheal Okpara University of Agriculture Umudike Abia State Nigeria. The leaves of both samples were separately washed with distilled water, spread at room temperature, allowed to dry for two weeks and pulverized using Thomas Willey milling machine into fine powder and sieved through 2mm mesh sieve. The dried ground samples were separately stored in two different airtight containers at room temperature.

EXTRACTION: 100grams of the dried ground bitter leaf samples were put in a 1000ml beaker and 500ml of deionised water added and stirred vigorously after which the plant residue was filtered through a muslin cloth and the obtained filtrate further filtered using Whatman No 1. Filter paper. The filtrate was evaporated to dryness using a rotatory evaporator (1, 15, 16). The same procedure was carried out for *Ocimumgratissimum* (scent leaf).

PHYTOCHEMICAL ANALYSIS: The qualitative and quantitative phytochemical screening of the extracts of *Ocimumgratissimum* and *Vernoniaamygdalina* for secondary metabolites such as tannins, alkaloids, saponins, flavonoids, phenols, steroids, anthraquinones and terpenoids were carried out following the standard procedures according to the methods described by Abdullahi (17, 18).

PROXIMATE ANALYSIS: The dried ground samples of *O.gratissimum* and *V. amygdalina* were analyzed for percentage moisture, ash, fibre, fat, carbohydrate and protein using the methods of AOAC and Mgbemena (19, 20).

MINERAL ANALYSIS: The digested samples were analysed for minerals using their different methods of analysis. Flame photometer (Jenway Ltd, Dunmow Essex UK) was used to determine calcium, magnesium, potassium and sodium contents in the samples while Atomic Absorption Spectrometer (Bulk Scientific East Norwalk, USA) was used for the determination of iron, copper and zinc (20).

RESULTS AND DISCUSSION

The results of the qualitative and quantitative phytochemical analysis of the leaves extracts of *Vernoniaamygdalina* and *Ocimumgratissimum* were shown in Table 1.

Table 1: The results of the Qualitative and Quantitative Phytochemical analysis of the water extract of the leaves of *Vernoniaamygdalina* and *Ocimumgratissimum* (mg/100g)

Phytochemicals	<i>Vernonia amygdalin</i>		<i>Ocimumgratissimum</i>	
	Qualitative	Quantitative	Qualitative	Quantitative
Saponin	+++	5.71 ± 0.12	++	3.52 ± 0.01
Tannin	++	4.90 ± 0.23	+	2.48 ± 0.12
Flavonoids	++	4.60 ± 0.01	+	2.00 ± 0.03
Steroids	+	0.50 ± 0.02	+	0.48 ± 0.02
Terpenoids	+++	5.64 ± 0.11	++	3.40 ± 0.11
Anthroquinones	-	0.06 ± 0.14	-	0.07 ± 0.14
Alkaloids	++	3.16 ± 0.16	+	2.16 ± 0.02
Phenol	-	0.07 ± 0.06	+	0.73 ± 0.13

±

(+++)=highly present (++) = moderately present (+) =present (-) = absent




Figure 1: Phytochemical Compositions of Vernonia amygdalina and Ocimum gratissimum

The results of the qualitative phytochemical analysis of *Ocimum gratissimum* and *Vernonia amygdalina* revealed the presence of saponin, tannin, flavonoids, steroids, terpenoids and alkaloids. *O. gratissimum* also contained phenols but lacked anthroquinones while *V. amygdalina* contained both anthroquinones and phenols. However, the quantitative phytochemical analysis revealed the presence of all the secondary metabolites in varied quantities in both leaves extracts. Saponin was the highest among the phytochemicals determined in both leaves extracts with the value 5.71 ± 0.12 mg/g in *V. amygdalina* and 3.52 ± 0.01 mg/g in *O. gratissimum*, this is followed by terpenoids with the value 5.64 ± 0.11 mg/g in *V. amygdalina* and 3.40 ± 0.11 mg/g in *O. gratissimum*, tannin 4.90 ± 0.23 mg/g in *V. amygdalina* and 2.48 ± 0.03 mg/g in *O. gratissimum*, flavonoids 4.60 ± 0.01 mg/g in *V. amygdalina* and 2.00 ± 0.03 mg/g in *O. gratissimum*, alkaloids 3.16 ± 0.16 mg/g in *V. amygdalina* and 2.00 ± 0.02 mg/g in *O. gratissimum*, steroids 0.50 ± 0.02 mg/g in *V. amygdalina* and 0.48 ± 0.02 mg/g in *O. gratissimum*, phenol and anthroquinines were higher in *O. gratissimum* than in *V. amygdalina*. The results of phytochemical screening agreed with the works of Alexandra (21) and Udochukwu (22) that reported the abundance of bioactive components in these leaves. The presence of these metabolites showed the great potentials of these two vegetables as good sources of useful phytomedicines. The presence of saponins in plants is responsible for the various biological benefits like anti-inflammatory, anti-diabetic, anti-HIV and anti-atherosclerosis. It is very effective

in maintaining liver function, lowering blood cholesterol, preventing peptic ulcer. It was also found to reduce nutrient utilization and conversion efficiency as in ruminants (23, 24). The non sugar part of saponin has direct anti-oxidant activities which may result in other benefits such as reduced risk of cancer and heart diseases in man (25). Terpenoids which ranked second play a vital and diverse role in the field of foods, drugs(as it is exploited in the fight against cancer, malaria, inflammation and a variety of infectious diseases) , cosmetics, hormones ,vitamins etc (26,27). Many alkaloids are used as drugs some of which include; nicotine, quinine, caffeine, cocaine and morphine. They have wide range of pharmacological activities such as anti-malaria (quinine), anti-asthma (ephedrine), anti-cancer, analgesics, (morphine), caffeine in tea, coffee stimulate and balance the nervous system (28). Flavonoids are polyphenolic compounds which contribute to many other colours found in nature. They have been reported to have anti-viral and anti-allergic activities. Flavonoid, quercetin is known for its ability to relieve hay fever, eczema and asthma (29). The anti-oxidants present in flavonoids may act in synergy with other phytochemicals in *V.amygdalina* and *O.gratissimum* to produce the medical benefits inherent in their leaves. The body immune systems, DNA are made from steroids. The type of steroids used to treat diseases is called corticosteroids. The results of qualitative and quantitative phytochemical analysis of this study revealed the presence of enough secondary metabolite in *V.amygdalina* and *O.gratissimum*.

Table 2 Proximate analysis of *Vernoniaamygdalina* and *Ocimumgratissimum* (%)

Components	<i>Vernoniaamygdalina</i>	<i>Ocimumgratissimu</i>
Moisture	12.28±0.02	10.80±0.01
Crude Protein	35.37±0.11	22.20±0.02
Crude Fat	6.00±0.20	5.75±0.10
Crude Fibre	7.20±0.01	6.00±0.03
Total Ash	5.87±0.03	3.40±0.11
Carbohydrate	32.28±0.01	51.85±0.11




Fig 2: Proximate Compositions of Vernonia amygdalina and Ocimum gratissimum

Table 2 and figure 2 showed the result of proximate analysis of *V. amygdalina* and *O. gratissimum* leaves extracts, crude protein was $35.37 \pm 0.11\%$ in *V. amygdalina* and $22.20 \pm 0.02\%$ in *O. gratissimum*, crude protein was the highest of the food components in both leaves. *V. amygdalina* had higher protein than *O. gratissimum*. Proteins are important in the body for the production of hormones, enzymes and blood plasma. They are immune boosters and can help in cell division as well as in growth (30). Eating of *V. amygdalina* leaves are encouraged because of its high protein content. Carbohydrate was $32.28 \pm 0.01\%$ in *V. amygdalina* and $51.85 \pm 0.11\%$ in *O. gratissimum*, carbohydrate was higher in *O. gratissimum* than *V. amygdalina* and also the highest of all the food components analysed. The result of carbohydrate agreed with the work of Mgbemena et al (20) with the value of $51.7 \pm 0.13\%$ in the leaves of *Irvingia gabonensis* and $38.38 \pm 0.14\%$ in *Irvingia wombulu*. The high content of carbohydrates in the leaves of the two species makes them a good source of energy. Moisture content was $12.28 \pm 0.02\%$ in *V. amygdalina* and $10.80 \pm 0.01\%$ in *O. gratissimum*. *O. gratissimum* had lower moisture content than *V. amygdalina* (table 2). The moderate moisture contents showed that the leaves are less prone to deterioration since foods with high moisture are prone to deterioration (30). Moisture dissolves other substances and carries nutrients throughout the systems leading to effective performance of the organs (31). Crude Fibre was $7.21 \pm 0.01\%$ in *V. amygdalina* and $6.00 \pm 0.03\%$ in *O. gratissimum*. Generally dietary fibre helps in digestion and functions the body to slow down the rate of glucose absorption into the blood stream thereby reducing the risk of hyperglycemia, the levels of plasma cholesterol and hence preventing colon cancer and cardiovascular diseases (32). Crude fat was $6.00 \pm 0.20\%$ in *V.*

amygdalina and 5.75 ± 0.10 % in *O.gratissimum*. The value of crude fat was almost the same in the two species .Dietary fats are important not only because of their high energy value but also the fat-soluble vitamin and essential fatty acids contained in the fat of natural foods. Ash contents was 5.87 ± 0.03 % in *V.amygdalina* and 3.4 ± 0.03 % in *O. gratissimum* .Ash content was higher in *V. amygdalina* than *O. gratissimum*. Higher ash content in the leaves of *V. amygdalina* is an indication of high minerals preserved in it (33).

1.2

Table 3 Mineral compositions of *Vernoniaamygdalina* and *Ocimumgratissimum*leaves extracts (mg/kg)

Mineral	<i>Vernoniaamygdalina</i>	<i>Ocimumgratissimum</i>
Calcium	1.28 ± 0.03	1.84 ± 0.01
Magnesium	0.07 ± 0.01	0.08 ± 0.02
Potassium	4.00 ± 0.02	4.20 ± 0.01
Sodium	0.11 ± 0.12	0.14 ± 0.03
Iron.	2.50 ± 0.01	2.65 ± 0.02
Copper	9.45 ± 0.11	16.60 ± 0.03
Zinc	46.80 ± 0.03	58.00 ± 0.12




Fig 3: Mineral Compositions of *VernoniaAmygdalina* and *OcimumGratissimum*

Table 3 and figure 3 showed the mineral composition of *Vernoniaamygdalina* and *Ocimumgratissimum* in mg/kg, the result revealed very high concentration of zinc compared to the other elements. The level of zinc in *O. gratissimum* was 58.00 ± 0.12 mg/kg and higher than that in *V.amygdalina* with the level 46.80 ± 0.03 mg/kg. The result was not in agreement with the work of

Alexandra(29) that had low levels of zinc (0.20 ± 0.06 mg/kg) in *O.gratissimum* . The high level of zinc is necessary as zinc is required for optimal functioning of physiological and biochemical processes, however, adequate care should be taken as excessive intake of zinc and iron is capable of causing vomiting, dehydration, electrolytic imbalance and lack of muscular co-ordination (34). The concentration of copper in *O. gratissimum*(16.60 ± 0.03 mg/kg) was far higher than that of *V. amygdalina*(9.45 ± 0.11 mg/kg) and also the second highest among the minerals .The result was in agreement with the work of Olumideetal; (35)that had 17.60 for *O.gratissimum*and 10.55 for *V. amygdalina* . High concentration of copper is recommended as copper deficiency causes cardiovascular disorders as well as anemia. Cu is an essential element in the human body and exists as an integral part of Cu proteins ceruloplasmin, which is concerned with the release of iron from the cells into the plasma. The concentration of iron in *V. amygdalina*was 2.50 ± 0.01 mg/kg and 2.65 ± 0.02 mg/kg in *O. gratissimum*, the level of iron in *O.gratissimum* (2.65 ± 0.02 mg/kg) was higher than in *O.gratissimum*.The level of Fe in this study did not agree with the work of Mgbemenaetal; (20) that indicated low iron in the leaves of *I.gabonensis* and *I. wombulu*. The presence of Zn, Fe and Cu are necessary in leafy vegetables as they could serve as good sources of anti oxidants (36, 37), thus the high presence of many phytochemicals in the two leaves are very advantageous. The study also revealed the presence of macro- elements calcium, magnesium, potassium and sodium in varying quantities. The levels of these macro-elements were higher in *O.gratissimum*than *V.amygdalina*(Table 3). Potassium had the highest concentrations among the macro- elements with the value 4.20 ± 0.01 mg/kgin *O.gratissimum*and 4.00 ± 0.02 mg/kg in *V. amygdalina*. The result of potassium agreed with the work of Olumideetal; (35) that had 4.30 for *O.gratissimum* and 3.70 for *V. amygdalina*but did not agree with the work of Awe et al; (33) that had the value 2.28 in *O.gratissimum* and 0.08 in *V. amygdalina* . Potassium is actually needed in high amount as it is responsible for nerve action regulation and osmo- regulation in the body fluid. Potassium and sodium play important role in maintaining the water balance within cells (38). Calcium was 1.87 ± 0.01 mg/kg in *O.gratissimum* and 1.28 ± 0.03 mg/kg b in *V.amygdalina*. Calcium is required for strong bones and teeth. Magnesium and sodium had low values. Magnesium helps in maintaining a normal heart rhythm (38).

CONCLUSION

Quantitative phytochemical analysis of *Vernoniaamygdalina* and *Ocimumgratissimum* reveals the presence of phytochemicals such as saponins, tannins, flavonoids, steroids, terpenoids, anthroquinones and phenols in varied quantities while qualitative phytochemicals reveals the absence of anthroquinones in both leaves and phenol in *V.amygdalina*. Saponin was found in highest quantity followed by terpenoids. *V.amygdalina*contained higher quantities of the phytochemicals than *O.gratissimum*. Proximate analysis revealed the presence of moisture, protein, crude fat, crude fiber

ash and carbohydrates with protein and carbohydrate possessing higher percentage of all of them and protein higher in *V.amygdalina*. The use of *V.amygdalina* is highly recommended. Protein found higher in *V.amygdalina* is used to enhance the penetration of macromolecules through cell membrane and this confirms the need for the high use of *V.amygdalina*. The leaves of these two vegetables contained the following macro and micro minerals; calcium, magnesium, potassium, sodium, iron, copper and zinc. Potassium was the highest macro –elements while zinc was the highest micro elements. Zinc was the highest mineral found in the two leaves. *Ocimumgratissimum* contained higher mineral elements than *V.amygdalina*. The use of these two species of vegetables is therefore recommended because of their medicinal, mineral and food values.

REFERENCES

1. Kolawole, S., Obueh, H. and Ukachi, A. (2018). Antibacterial Effect of *VernoniaAmygdalina* and *Ocimumgratissimum* on some Bacteria that cause opportunistic Infections in HIV/AIDS Patients. *Journal of Nutritional Health.Food Science* 6(2) 1-5.
2. Tapsil, I.C., Hemphil, I. and Chobiac, L. (2006). Health benefits of herbs and spices, the past, present and future”. *Med. Journal, Australia* .185(4) 4-24.
3. Lai, P.K. and Roy, J.R. (2004). Antimicrobial and Chemopreventive properties of herbs and spices. *CurrMed.Chem* 11(11) 1451-1460.
4. Sofowora, A. (1999). *Medicinal plant and traditional medicine in Africa*. 2nd edition, Spectrum Books Limited.Ibadan, 178-188.
5. Odugbeni, T. (2006). *Outline of medicinal plants in Nigeria*, 1stEdition,University of Lagos Press. Nigeria 7780.
6. WHO (2004). *Guideline for the Assessment of Herbal Medicine*, Geneva 270
7. Effraim, K. D., Jacks, T. W. and Sopio, O. A. (2003). Histopathological Studies on the Toxicity of *Ocimumgratissimum* Leaves Extracts on some organs of Rabbit, *African Journal of Biomedical African Journal Biomedical Research* 6:21 – 28.
8. Erinle, O. D. (2012). Phytochemical Screening of Scent leaf (*Ocimumgratissimum*) and Bitter leaf (*Vernoniaamygdalina*) extracts ,N.D.Project ,FCFFT, New Bussa, Niger State 1-5
9. Oludosu-Ajahi, R.N., Dienye, H.E., Ajayi, C.T. and Erinle, O.D.(2017). Comparative Screening of Phytochemicals Compounds in Scent leaf *Ocimumgratissimum* Linn.(Family. Lamiaceae) and Bitter leaf *Vernoniaamygdalina* Dei. (Family: Asteraceae)Extracts *Advances in Resaerch* 6 21-25 *Zoology and Botany* 5(4) 50-54.
10. Health Facts NG (2015). *Health Facts*. Designed by iQubeLabs.www.healthfactsng 1-5.
11. Iwu, M.M. (1993). *Handbook of African Medicinal Plants*, Florida CRC Press Inc.BocaRation. 40.
12. DI Stasi, L.C., Oliveira, G.P., Carvalhaes, M.A., Queiroz, M. Jr., Tien,O. and Kkinmi. S. H. (2002). Medicinal plants popularly used, *J. of Physical and Chemical Sciences Volume 1* issue 1 ISSN: 2348-3278 in the Brazilian Tropical Atlantic Forest *Filoterapia* 73 69-91.
13. Ijeh, I.I., Njoku, O.U. and Ekenze, E. C.(2004). Medicinal Evaluation of *Xylopiiaethiopica* and *Ocimumgratissimum*; *Journal of Medicinal Aromatics Science*, 36(1) 44-49.
14. Muanya, C. (2015).Scent and Bitter leaves Retard Growth of Cancers, Boost Immune System *Guardian Newspaper Publication* of 3rd September 30.
15. Akujobi, C.O., Ogbulie, J.N. and Njoku, H.O. (2010). The extracts of *Ocimumgratissimum* on the dehydrogenase activities to clinical isolates of *Eschricha coli* and *Staphylococcus aureus*. *Journal of Agricultural Technology*, 6(1)57-65.
16. Audu, I., Shuaibu, B.S., Alona, C. L. ,Murtala, M.M. and Jiya, A.(2018). Phytochemical Analysis and Antimicrobial Activity of Bitter leaf (*Vernoniaamygdalina*) collected from Lapai, Niger State, Nigeria on some selected pathogenic micro-organism.
17. Abdullahi, M. (2012). Phytochemical constituents and antimicrobial and grain protectant activities of clove Basil (*Ocimumgratissimum* L) grown in Niger. *Journal of Plant Research*, 2(1)51-55.

18. Agholor, K., Lucy M. Yaki, Idris ,Abubakar, Lucy F. Olusoh and RakiyaZubairu (2018). Anti bacterial activity of *Ocimumgratissimum* (scent leaf) on some pathogenic gastrointestinal 17.
19. AOAC (Association of Official Analytical Chemists) (2016). Official Methods of Analysis, 20thEdWashington DC, USA: 120.
20. Mgbemena, N. M. ,Ilechukwu, I. ,Okwunodulu, F. U., Chukwurah, J.O. and Lucky, J.B. (2019).Chemical Composition, Proximate and Phytochemical analysis of *Irvingiagabonensis* and *Irvingiawombulu* peels, seed coat , leaves and seeds. *Ovidius University of Anals of Chemistry Science do Vol.30 (1)65-69*.
21. Alexander, Priscilla. (2016). Phytochemical Screening and Mineral Composition of the leaves of *Ocimumgratissimum* (Scent leaf)*Int.J.Appl.Sci.Biotechnolvol.(4) 2*.
22. Udochukwu, U., Omeje, F.I., Uloma,I.S. and Oseiwe ,F.D.(2015).Phytochemical Analysis of *Vernoniaamygdalina* and *Ocimumgratissimum* Extracts on their Antibacterial activity on some drug resistant bacteria .*Journal of Research communication.Vol. 3(5)225-235*.
23. Kashiwada, Y., Wang, H.K., Nagao, T., Kitanaka, S., Yasuda, I. and Fujioka, T. (2009). Anti-AIDS agents 30 Anti Activity of oleanolic acid pomolic acid and structurally related triterpenoids. *Journal of Natural Product, 61, 1090-1095*.
24. Banno, N., Akihia, T., Yasukawa, K., Higashihara, H. M., Ukiya and Watanabe, K.(2004). Triterpene acids from leaves of *Perillafrutescenes* and their anti- inflammatory and anti-tumour promoting effects. *Bioscience Biotechnology and Biochemistry 68, 85-90*.
25. Rhoades, D.F.(2009). Evolution of Plant Chemical Defence against Herbivores .In G.A.Rosenthal and D.H. Janzen; *Herbivores: Their Interaction with Secondary Plant Metabolites (Academic Press, New York) 41*.
26. Armella, T. Mbaveng and Victor Kuete (2014). Harmful and protective effects of Terpenoids from African Medicinal Plants *Toxicological Survey of African Medicinal Plants. Elsevier Inc.557-576*.
27. Perveen, S. (2018). Terpenes and Terpenoids. Introductory Chapter Intech Open <https://www.intechopen.com>.
28. Qui, S., Sun, H. , Zhang , A.H. , Xu, H.V., Van, G.I.and Han, Y. (2004). Natural alkaloids;basi aspects, biological roles and future perspectives roles and future. *Chin.J.Med. 12 (6)401-406*.
29. Prabhavathi, R.M., Prasad, M.P. and Jayaram ,M. (2016).Studies on Qualitative and Quantitative Phytochemical Analysis of *Cissusquadrangularis*. *Pelagria Research Library Advances in Applied Science Research .7(4): 11-17*.
30. Bouttwell, R.K.(1998). An over view of the role of Nutrition in carcinogenesis. In *Nutrition Growth and Cancer, Harms ,C.C. and P.A. Cerutti (Eds) Alan.R. LlssLondon.pp 387-418*.
31. Okeke, C.U. and Ekekwa , I.(2006). Proximate and preliminary phytochemicals analysis of avocado pear *Persea, gratissimaGaertn.f.(family. Lauraceae) Niger J.Bot.19 : 156-162*.
32. Ilodibia, C. V., Ugwu, U.R. , Okeke, C. U., Ezeabara, C. A., Okeke, N.F. ,Akachukwu, E.E. and Aziagba, B. O. (2014). Determination of proximate composition of various parts of Two *Dracaena specie.Int. J.Bot. 10(1) 37-41*.
33. Awe, S., Okeke, O.L. and Usman A.A.(2018). Comparative of proximate and mineral composition of selected fruits and vegetables .*FutaJournalof Research in Sciences, Vol.14(1) 67-74*.
34. 35. Olumide, M.A., Ajahi, O.A. and Olufunso, E.A. (2019). Comparative Study of proximate, mineral and phytochemical analysis of the leaves of *Ocimumgratissimum*, *Vernoniaamygdalina*and*Moringaoleifera**Journal Medical Plant Research 13(15) 301 – 356*.
35. Ogbodna, G.E. and Kabari L.G. (2015). Heavy Metal Content in Bitter leaf (*Vernoniaamygdalina*) grown Along Heavy Traffics Routes in PortHarcourt, *Agricultural Chemistry, Margarita Stoytcheva and Roumen 2(atev, Intech, 201 – 210*.
36. Tudor, R. ,Zalewski, P.D. and Ratnaike, R.N.(2005).Zinc in health and chronic disease. *J.Nutri.Health Aging 9: 45-50*
37. Okunlola, G .O.,Jumoh, M.A., Olatunji, O.A., RufiaAbdulfatai, B. andAdepeju,O.Omidiran (2018). Proximate analysis, mineral composition and antioxidant properties of bitter leaf and scent leaf, *International Journal of Vegetable Science Taylor and Francis 1-10*.

38. Hollenman, G., Arnold, F., Wiberg, G. and Egon, W.W. (1988). "Natrium" lehrbuch der Anorganischen Chemie (in German) (91-100 ed) Walter de Gruyter pp 931-943.

UNDER PEER REVIEW